

DIATOM BASED POLLUTION MONITORING IN URBAN WETLANDS OF COIMBATORE, TAMIL NADU



Karthick, B.
Alakananda, B.
Ramachandra, T. V.

ENVIS TECHNICAL REPORT – 31




February - 2009

ENVIRONMENTAL INFORMATION SYSTEM [ENVIS]
Centre for Ecological Sciences, Indian Institute of Science,
Bangalore - 560012, INDIA



Web: <http://ces.iisc.ernet.in/energy/>
<http://ces.iisc.ernet.in/biodiversity>
Email: cestvr@ces.iisc.ernet.in
energy@ces.iisc.ernet.in





Energy and Wetlands Research Group
Centre for Ecological Sciences, Indian Institute of Science,
Bangalore - 560012, INDIA

Web: <http://ces.iisc.ernet.in/energy/>
<http://ces.iisc.ernet.in/biodiversity>

Email: cestvr@ces.iisc.ernet.in; energy@ces.iisc.ernet.in

Telephone: 91-80-23600985/22932506/22933099

Fax: 91-80-23601428/23600085/23600683[CES-TVR]



Contents

Contents	1
List of Figures.....	Error! Bookmark not defined.
List of Tables.....	2
Summary	3
Introduction.....	4
Need to study Wetlands	5
Wetland monitoring	5
Biological monitoring	6
Diatoms	7
Diatoms as bio-indicators	8
Objectives.....	9
Study Area	9
Methods and Materials	11
Diatom sampling	11
Water sampling.....	11
Ecological Diversity and diatom indices	11
Results and Discussion	16
Diatom Diversity.....	16
Dominance.....	19
Evenness.....	19
Shannon index.....	20
Fisher alpha diversity Index.....	21
Berger-Parker Index	22
Diatom assemblages and trophic condition.....	23
Diatom Indices	24
Habitat preference	26
Conclusion	32
Acknowledgement	32
References	33
Annexure: I - List of Species.....	40
Annexure: II - Illustrations	44

List of Figures

Figure 1 Coimbatore city with the sampling points. (Maps Courtesy: Google)	10
Figure 2 Relative abundance of four most dominant genera plotted with sites arranged in order of increase in electrical conductivity (left) and pH (right).	18
Figure 3 Dominance Index across sites.....	19
Figure 4 Evenness Index across sites	20
Figure 5 Shannon Index across sites.....	21
Figure 6 Fisher alpha Index across sites	21
Figure 7 Berger- Parker Index across sites.....	22

List of Tables

Table 1 Diversity parameters and indices	12
Table 2 Diatom Indices.....	15
Table 3 Water Quality Variables of Coimbatore Wetlands.....	16
Table 4 Diversity indices for Coimbatore wetlands.....	17
Table 5 Diatom Indices values for the wetlands	24
Table 6 Class limit values for Diatom indices.....	25
Table 7 Trophic condition of the wetlands with dominant species.....	25
Table 8 Species list with their occurrence in three habitats	27

Diatom Based Pollution Monitoring in Urban Wetlands of Coimbatore, Tamil Nadu

Summary

Diatoms comprise a ubiquitous, photosynthetic and distinctive group of essentially unicellular algae. They are more specific in their preference and tolerance of environmental conditions than most other aquatic biota and have long been recognised as excellent indicators of ecological status of water bodies. This study documents the diatom flora of six urban wetlands of Coimbatore city, examines benthic diatom assemblages across different habitats and investigates pollution status based on diatom composition. 96 Species belonging to 34 genera were recorded and out of them 27 species were dominant. The dominant species that are cosmopolitan include *Cyclotella meneghiniana*, *Nitzschia sp.*, *Sellaphora pupula*, *Gomphonema parvulum* and *Navicula sp.* Singanallur wetland and Noyyal river stretches are characterised by pollution tolerant species with low diatom diversity. Diatom assemblages indicate wetlands; Vedapatti, Perur and Sundakamuthur are moderately polluted, while Pallapalayam, Noyyal River and Singanallur wetlands are heavily polluted.

Keywords: Urban wetlands, pollution indicators, diatom-indices, diatom assemblages, *Cyclotella*, Coimbatore

Diatom Based Pollution Monitoring in Urban Wetlands of Coimbatore, Tamil Nadu

Introduction

Wetlands are an essential part of human civilization, meeting many crucial needs for life on earth such as drinking water, energy, fodder, biodiversity, flood storage, transport, recreation, and climate stabilizers. They also aid in improving water quality by filtering sediments and nutrients from surface water. Wetlands play a major role in removing dissolved nutrients such as nitrogen and to some extent heavy metals (Ramachandra *et al.*, 2002). Hence, they are often described as “Kidneys of the landscape”. Wetlands encompass many different habitats including wetlands, marshes, swamps, flood plains, bogs, shallow ponds, littoral zones of larger water bodies and peatlands. All these share the fundamental feature of complex interactions among basic components such as soil, water, flora and fauna.

Wetlands are ecologically important in relation to stability and biodiversity in a region and also in terms of energy and material flow. Wetlands are “lands transitional between terrestrial and aquatic ecosystems where the water table is usually at or near the surface or the land is covered by shallow water” (Mitch and Gosselink, 1986). Hydrological conditions of a wetland modify or change chemical and physical properties such as nutrient availability, degree of substrate anoxia, soil salinity, sediment properties and pH, which in turn, influence the biotic integrity (Gosselink and Turner, 1978). Wetlands retain water during dry periods, thus keeping the water table high and stable. During floods they diminish floods intensity and biotic components trap suspended solids and attached nutrients. A healthy wetland retains a natural flow of water, minimising flooding in the catchment. Wetlands receive water deposited as groundwater, during dry seasons. Thus, a healthy wetland does the function of water recharge and discharge effectively, while meeting the human needs. However, humans have altered the natural

flow regime of wetlands either by altering the natural drains, changing the land cover drastically or letting the untreated sewage in urban areas in recent times. The removal of such wetland systems or letting untreated sewage has caused the deterioration of water quality and ecological degradation in catchment (Prasad *et al.*, 2003).

In India, wetlands are distributed in all the biogeographic regions occupying 58.2 million hectares, including areas under wet paddy cultivation (Directory of Indian Wetlands, 1990). They exhibit significant ecological diversity, primarily because of variability in climate, habitat and topography. Today, wetlands are one of the most threatened habitats in India. They have been converted for agriculture, industry or settlements and some are affected by industrial effluents, sewage, household wastes and sedimentation. Due to urbanization and lack of holistic approaches in land management, land and waterbodies in and closer to urban centres have been targeted. The water crisis, frequent flooding in urban areas has necessitated understanding the role of wetlands, and the need for integrated approaches to maintain the ecological balance, while meeting the demands of the growing population.

Need to study Wetlands

Rising water demand has exacerbated the impacts. Societies need to adopt improved strategies for integrated wetland management to ensure the quantity and quality of water is maintained for the ecosystem functions. In this regard, Ramsar Convention's Agenda 21 recommends the work towards better understanding of these threatened ecosystems through basic research, awareness and education, ecosystem and species conservation.

Wetland monitoring

Effective assessment tools are needed for consistent evaluation of the condition with stressors of wetland resources for solving problems. This entails inventorying and

regular monitoring of wetlands. Physical and chemical monitoring of water quality has been practiced for a long time. Standard techniques are used for measuring light penetration, turbidity, conductivity, dissolved oxygen, biological oxygen demand and nutrients like phosphates, nitrates, nitrites, ammonia, and so on (Chapman, 1992). These measurements even though provide us simple values, but don't provide overall health and condition of the ecosystem enabling both preventive as well as restorative measures. Many environmental factors vary on different spatial and temporal scales in complex ecosystems such as wetlands. These variables range from climate, landuse, and geomorphology of a watershed (eg, Richards *et al.*, 1996) to the physical, chemical and biological characteristics. In this context, monitoring involving biological communities of an ecosystem would help in assessing, as they can integrate and reflect the effects of chemical and physical disturbances that occur in short duration as well as over extended period of time.

Biological monitoring

Monitoring using organisms, to assess the ecosystem's condition is referred as biological monitoring or biomonitoring. Biological indicators based on organisms living from one day to several years provide an integrated assessment of environmental conditions in streams, rivers and wetlands that are spatially and temporally variable. An ideal biomonitoring should be useful for both long and short term monitoring. Current conditions may be linked to the past conditions very effectively, if the same biomonitors are used for both short and long-term monitoring (Dixit *et al.*, 1992). Biomonitoring consists of groups of species, each group with well defined habitats, so that they may reflect changes in a variety of habitats. Biological indicators are important parts of environment assessment because protection and management of these organisms are the objectives of most programs. Aquatic communities (like algae, fish, riparian vegetation, macro-invertebrates), integrate and reflect the effects of chemical and physical disturbances. A biota that undergoes change from dominance to gradual disappearance of a species is of ecological significance. The primary aim here is to detect changes in

abundance, structure and diversity of a target species assemblage as compared to the reference condition. Bio-indicators include organisms that are:

- close to the transfer of nutrient and energy in the food web;
- wider range of distribution;
- simple life-cycle stages, and identifiable to the species or even the morphotype level;
- sensitive to fine changes in the environment with a range of tolerance; and
- preference to environmental variables, so a change in the environment is reflected by a shift in species dominance.

Now, biological monitoring has begun to address the question of biological integrity of wetlands influenced by various anthropogenic land use activities.

Numerous methods have been developed in biomonitoring for an assessment of the integrity of aquatic systems. Most are based on the attributes of whole assemblages of organisms such as fish, algae or invertebrates. A variety of assemblages have been used in biological assessments ranging from macrophytes (Galatowitsch *et al.*, 1999, Gernes and Helgen, 1999) macroinvertebrates (Kerans and Karr, 1994 and Barbour *et al.*, 1996); amphibians (Micacchion, 2004); fish (Schulz *et al.*, 1999); birds (O'Connell *et al.*, 1998) and diatoms (Fore and Grafe, 2002).

Diatoms

Diatoms under Class *Bacillariophyceae* comprise a ubiquitous, photosynthetic and distinctive group of unicellular algae. Diatoms are made up of siliceous cell wall consisting of two valves; epivalve and hypovalve which fit together like a petri dish together known as frustules. In between two valves series of bands are present known as girdle bands. During cell division the new frustules are formed from the inside of the cell. The outer or older is the epivalve and inner or newly formed one is hypovalve forms one daughter cell where as outer or older hypovalve acts as epivalve and newly

formed valve will become hypovalve. This forms another daughter cell. During this process cell size goes on decreasing. The original size is attained by undergoing sexual reproduction by auxospore formation.

Diatoms as bio-indicators

Diatoms are more specific in their preference and tolerance of environmental conditions than most other aquatic biota. Diatoms were the first group of biota used for detecting organic pollution (e.g., the saprobial system by Kolkwitz and Marsson in 1909, cited in Stoermer and Smol, 2001). Diatoms respond directly and sensitive to many physical, chemical and biological changes such as temperature, nutrient concentration and herbivory. They are sensitive to many habitat conditions and show variability in biomass and species composition. At higher spatial and temporal levels effects of resources and stressors on diatom assemblages can be constrained by climatic, geology and land use. Diatoms are readily distinguished to species and subspecies level based on unique morphological features. Diatoms have one of the shortest generation times of all biological indicators. They reproduce and respond rapidly to environmental change and provide early warning indicators of both pollution increases and habitat restoration success. Frustules are preserved in sediments and record habitat history. Diatoms collection and methods are ease and low cost. Samples can be archived easily for long periods of time for future analysis and long term records.

Diatoms occur in all types of environment where ever moisture is present. A golden-brown mucilage film on the surface of substrata indicates the presence of benthic diatoms whereas free living in the water column is the planktonic diatoms. Data on diatoms as indicators of water quality reflecting pH, salinity and organic pollution in Europe, America, South Africa and Japan have been available for a long time (e.g. Patrick, 1986; Schoeman, 1973; Round, 1986, 1990; Cox, 1991). However, there is no information available on diatoms as indicator species of wetlands in India. The present study assesses six major wetlands in an urban ecosystem using diatoms as bioindicators.

Objectives

Objectives of this research are to:

- i. determine the pollution status of selected wetlands of Coimbatore by using diatoms
- ii. prepare an illustrated guide to the common diatom flora of wetlands of Coimbatore.

Study Area

Coimbatore also referred popularly as Manchester of India is an important industrial city, located in Tamil Nadu ($10^{\circ}55' - 11^{\circ}10' \text{ N}$, and $77^{\circ}10' - 76^{\circ}50' \text{ E}$) at an average altitude of 470m, ranking 11th in terms of population (Figure 1). There are more than 30,000 small, medium and large industries including textile mills and foundries in the city employing about 40% of the population. The growing industrial sector and ensuing immigration of people pose heavy burden on the city infrastructure that did not grow in proportion. The city does not have facilities for treatment of industrial, municipal and domestic wastes. Wetlands and Noyyal River have been used for disposal of wastes of the city. Natural drainage networks have been converted to storm water drains for letting the sewerage into wetlands without any treatment. In Coimbatore city there are 28 wetlands, mostly fed by the river Noyyal. The river, flowing through the city on its south, originates in the Vellingiri hills in Western Ghats, located on the south-western side of the city. Some of these wetlands are seasonal and have also been used as dumping yard for garbage and industrial wastes during dry period (Mohan Raj *et al.*, 2000). During the monsoon, with the inflow of water, this activity leads to contamination of groundwater sources. Six wetlands selected for biomonitoring (Figure 1) are: Vedapatti (VPP), Pallapalayam (PPL), Sundakamuthur (SMS), Perur (PRP), Noyyal (NLP) and Singanallur (SNP).




Figure 1 Coimbatore city with the sampling points. (Maps Courtesy: Google)

Methods and Materials

Diatom sampling

Diatom samples were collected (from cobbles, aquatic plants and sediment) and prepared using standard methods as per Taylor *et al.*, (2005) from selected wetlands. Diatom communities were then analysed by counting between 400 and 450 valves. During enumeration the dimensions of diatom valve characteristics, like its length, width and striae densities in 10 μm were measured. Identification of diatoms is carried out using taxonomic guides (Gandhi, 1957 1959a, 1959b, 1961, 1962, 1967, 1998; Lange-Bertalot, 2001; Krammer, 2002; Taylor, 2007; Karthick *et al.*, 2008).

Water sampling

Water samples were collected from all sites and physical variables like pH, temperature, Electric conductivity, Salinity and Total dissolved solids were measured using EXTECH combo probe.

Ecological Diversity and diatom indices

Ecological diversity was calculated for each sample using diversity indices given in Table 1.

Table 1 Diversity parameters and indices

Index	Equation	Remarks	References	Eq. No
Abundance	$\frac{\text{No. of Individuals of a species} \times 100}{\text{No. of Sampling units}}$			1
Shannon Weiner's (H')	$H' = - \sum_{i=1}^S p_i \ln p_i$	The value ranges between 1.5 and 3.5 and rarely surpasses 4.5	Ludwig and Renolds (1998); Legendre and Legendre 1998	2
	Pi: proportion of individuals of i th species			
Simpson's	$D = \frac{\sum_{i=1}^S n_i(n_i - 1)}{N(N - 1)}$	The value varies from 0 to 1. A value of 0 indicates the presence of only one species, while 1 means that all species are equally represented.	Ludwig and Reynolds (1998)	3
Dominance	1-Simpson index $D = \sum \left(\frac{n_i}{n} \right)^2$ Where ni is number of individuals of taxon i.	The occupancy of a species over an area. Ranges from 0 (all taxa are equally present) to 1 (one taxon dominates the community completely)		4

Evenness	$H' = - \sum_{i=1}^S p_i \ln p_i$	The measure of biodiversity which quantifies how equal the community	5
Fisher's alpha	$S = a * \ln \left(\frac{1 + n}{a} \right)$ Where S is number of taxa, n is number of individuals and <i>a</i> is the Fisher's alpha.	It is a mathematical model used to measure diversity	6
Berger-Parker	$d = \frac{N_{max}}{N}$ Where N _{max} is the number of individuals in the most abundant species and N is the total number of individuals in the sample.	The number of individuals in the dominant taxon relative to n, where n is the total number of species	Berger and Parker 1970; 7

Diatom specific indices like Generic Diatom Index or GDI (Coste and Ayphassorho, 1991), the Specific Pollution sensitivity Index or SPI (Coste in Cemagref, 1982), the Biological Diatom Index or BDI (Lenoir and Coste, 1996), the Artois-Picardie Diatom Index or APDI (Prygiel *et al.*, 1996), Sládeček's index or SLA (Sládeček, 1986), the Eutrophication/Pollution Index or EPI (Dell'Uomo, 1996), Rott's Index or ROT (Rott, 1991), Leclercq and Maquet's Index or LMI (Leclercq and Maquet, 1987), the Commission of Economical Community Index or CEC (Descy and Coste, 1991), Schiefele and Schreiner's index or SHE (Schiefele and Schreiner, 1991), the Trophic Diatom Index or TDI (Kelly and Whitton, 1995), and the Watanabe index or WAT (Watanabe *et al.*, 1986) were also computed as listed in Table 2. All the diatom indices were calculated using Equation 8 (Zelinka and Marvan, 1961) except for the CEC, SHE, TDI and WAT index and all of the above indices, except TDI (maximum value of 100), the maximum value of 20 indicates pristine water.

$$index = \frac{\sum_{j=1}^n a_j s_j v_j}{\sum_{j=1}^n a_j v_j} \quad (\text{Equation: 8})$$

Where a_j = abundance (proportion) of species j in sample, v_j = indicator value and s_j = pollution sensitivity of species j .

The performance of the indices depends on the values given to the constants s and v for each taxon and the values of the index ranges from 1 to an upper limit equal to the highest value of s . Each diatom species used in the calculation/equation is assigned two values; the first value reflects the tolerance or affinity of the diatom to a certain water quality (good or bad) while the second value indicates how strong (or weak) the relationship is. Abundance and weighted average were computed. This would indicate how many of the particular diatoms in the sample occur in relation to the total number counted.

Table 2 Diatom Indices

Abbreviation	Full name	Reference
IPS	Specific Pollution Sensitivity Metric	(Coste, 1987)
SLAD	Sládeček's pollution metric	(Sládeček, 1986)
DESCY	Descy's pollution metric	(Descy, 1979)
L&M	Leclercq and Maquet's pollution metric	(Leclercq and Maquet, 1987)
SHE	Steinberg and Schiefele trophic metric	(Steinberg and Schiefele, 1988)
WAT	Watanabe <i>et al.</i> , pollution metric	(Lecointe <i>et al.</i> , 2003)
TDI	Trophic Diatom metric	(Kelly and Whitton, 1995)
EPI-D	Pollution metric based on diatoms	(Dell'Uomo, 1996)
ROTT	Trophic metric	(Rott <i>et al.</i> , 1999)
IDG	Generic Diatom Metric	(Lecointe <i>et al.</i> , 2003)
CEE	Commission for Economical Community metric	(Descy and Coste, 1991)
IBD	Biological Diatom Metric	(Prygiel and Coste, 1999)
IDAP	Indice Diatomique Artois Picardie	(Lecointe <i>et al.</i> , 2003)
IDP	Pampean Diatom Index (IDP)	(Gómez and Licursi, 2001)

Results and Discussion

Water samples were collected from all sites and physical variables like pH, Temperature, Electric conductivity, Salinity and Total dissolved solids were measured and are listed in Table 3. pH of sampled wetlands range from 7.4 to 9 indicating neutral to alkaline conditions. Electric conductivity ranges from 280 (Vedapatti) - 2250 μ S/cm (Singanallur).

Table 3 Water Quality Variables of Coimbatore Wetlands

Sampling site	Conductivity (μ S/cm)	Water Temperature ($^{\circ}$ C)	pH	Total Dissolved Solids (mg/L)
Vedapatti	280	29.6	7.47	195
Sundakamuthur	283	32.4	9.06	198
Sundakamuthur	283	32.4	9.06	198
Perur	347	29	7.92	242
Pallapalayam	733	27.9	9.05	511
Pallapalayam	770	29.3	8.83	543
Noyyal River	1121	29.7	7.7	781
Singanallur	2250	29.3	8.53	1590

Diatom Diversity

Diatom samples were collected (from cobbles, aquatic plants and sediment) and prepared using standard methods from Vedapatti, Pallapalayam, Sundakamuthur, Perur, Noyyal and Singanallur wetlands in Coimbatore. Diatom communities were analysed as explained in methods section. 96 Species belonging to 34 Genera were recorded from these wetlands, which are provided in Appendix 1. Among these species, 27 species were dominant (i.e., occurring >5% of any given community). Appendix 2 gives the species-wise light microscopic illustrations. Table 3 lists the diversity indices, which show a significant difference in community structures across the sampled

wetlands. Higher values of Shannon, Simpson and evenness values are for Pallapalayam wetland compared to Singanallur wetland, where dominance index was relatively higher.

Table 4 Diversity indices for Coimbatore wetlands

	VP	PP	SM	PP	SN	PR	SM	NL
Number of species	23	22	29	26	10	28	30	14
Shannon Index	2.371	2.498	2.066	2.621	0.4135	2.366	2.538	1.472
Simpson	0.8526	0.8877	0.7276	0.8764	0.1402	0.8545	0.874	0.6768
Evenness	0.4654	0.5529	0.2723	0.5289	0.1512	0.3805	0.4217	0.3114
Margalef	3.649	3.453	4.66	4.161	1.496	4.56	4.973	2.016
Equitability	0.756	0.8083	0.6137	0.8045	0.1796	0.71	0.7461	0.5579
Fisher alpha	5.247	4.88	7.143	6.189	1.85	7.013	7.927	2.536
Berger-Parker	0.306	0.1986	0.4963	0.2604	0.9268	0.2547	0.2317	0.4596

Note: VP- Vedapatti (Epiphytic); PP- Pallapalayam (Epilithic); SM- Sundakamuthur (Episammic); PP-Pallapalayam (Epiphytic); SN-Singanallur (Epiphytic); PR-Perur (Epiphytic); SM- Sundakamuthur (Epiphytic); NL-Noyyal (Epiphytic)

Common diatoms genera namely *Cyclotella*, *Gomphonema*, *Nitzschia* and *Fragilaria* accounted for large proportion of the community in all sites. Figure 2, a plot of genera across pH and electrical conductivity ranges reveal that:

- *Cyclotella* – present in neutral to high alkaline, and high electrolytic;
- *Gomphonema*, *Nitzschia* – present in entire pH and conductivity ranges; and
- *Fragilaria* – prefer Neutral to alkaline, and moderate electrolytic water.




Figure 2 Relative abundance of four most dominant genera plotted with sites arranged in order of increase in electrical conductivity (left) and pH (right).

Dominance

Dominance is the degree to which different species in an ecological community predominate, ranging from 0 (all taxa are equally present) to 1 (a taxon dominates the community completely). Dominance is calculated (equation 4, Table 1) and is given in Figure 3. Singanallur wetland has 10 species with *Cyclotella meneghiniana* as dominant species (dominance: 0.85), while *Aulocosira granulata* (19.86 %,) dominated Pallapalayam (dominance: 0.11) and *Sellaphora pupula* (23.17%) and *Gomphonema parvulum* (18.48%) were prominent species in Sundakamuthur (dominance: 0.12) wetlands. Remaining sites showed dominance index value between 0.1- 0.4.




Figure 3 Dominance Index across sites

Note: VP- Vedapatti wetland; PP- Pallapalayam wetland; SM- Sundakamuthur wetland; PP-Pallapalayam wetland; SN-Singanallur wetland; PR-Perur wetland ; SM- Sundakamuthur wetland; NL-Noyyal River

Evenness

Evenness is a measure of biodiversity which quantifies how equal the community is numerically. Figure 4 depicts the evenness computed as per equation 5, Table 1.

Cyclotella meneghiniana constitute more than 90% of the total population accounted for low evenness in Singanallur wetland. In Pallapalayam wetland 22 species were recorded, among *Aulocosira granulata* and *Nitzschia obtuse* were represented by 19.86% and 14.61% abundance respectively.




Figure 4 Evenness Index across sites

Shannon index

Shannon diversity index (H') computed as per equation 2 (Table 1) takes into account the number of individuals as well as number of taxa. This varies from 0 for communities with only a single taxon to high values for communities with many taxa, each with few individuals. Low H' was recorded in Singanallur wetland (0.4135, *C. meneghiniana* representing 92%) and Noyyal River (1.472, *Nitzschia sp* representing 45.95 %). Species abundance in other sites ranges from 20 to 26 species that represent 15-50% of the total population.




Figure 5 Shannon Index across sites

Fisher alpha diversity Index

High Fisher's alpha diversity index computed (equation 6, Table 1) was noticed in Sundakamuthur (7.8), Pallapalayam (6.2) and Perur (7) wetlands. Singanallur wetland and Noyyal River with less number of taxa shows low index value.




Figure 6 Fisher alpha Index across sites

Berger-Parker Index

Berger-Parker is calculated (equation 7, Table 1) from the number of individuals in the dominant taxon relative to the total number of species. *Cyclotella meneghiniana* is the dominant species (with 92.68% abundance) showing a high index value in Singanallur wetland. *Diadesmis confervaceae* and *Gomphonema turris* was observed as abundant species (15-30%) in Vedapatti wetland. In Pallapalayam wetland *Aulocosira granulata*, *Cyclotella meneghiniana* and *Nitzschia obtuse* represents to 13-20% of the population from an epilithic habitat and *Aulocosira granulata*, *Cyclotella meneghiniana* represents 19-26% of the population from an epiphytic habitat. *Gomphonema parvulum* and *Cymbella turgid* represents 25.47 and 22.25% respectively in Perur wetland. *Nitzschia sp.* (45%) and *Navicula sp.* (32%) characterize the Noyyal River. In Sundakamuthur wetland, *Sellaphora pupula* 49.63 % and *Navicula rostellata* 12.04% in episammic habitat whereas 23.17% *Sellaphora pupula* and 18.48% *G. parvulum* being present in epiphytic habitat (Figure 6).




Figure 7 Berger- Parker Index across sites

Diatom assemblages and trophic condition

Distribution of diatoms reflects the average ecological conditions of water (Cholnoky 1968; Lowe 1974). In Vedapatti wetland, cosmopolitan extreme pollution resistant species *Diadomesmis confervaceae*, *Gomphonema gracile* and *G.turris* were dominant among 23 species highlighting eutrophic status of water with higher electrolyte. *Aulocosira granulata* and *Cyclotella meneghiniana* are dominant among 22 species in the epilithic substrata and 26 species in epiphytic substrata of Pallapalayam wetland. These species are cosmopolitan in distribution in both benthic and plankton representing electrolyte rich and brackish inland water.

Episammic sample from Sundakamuthur is dominated by *Sellaphora pupula* and *Navicula rostellata*, which are more tolerant to high levels of pollution. Epiphytic substratum sample is represented by *Gomphonema parvulum* and *G. affine*, which are tolerant to extreme pollution and occurs in water with elevated electrolyte. *Cyclotella meneghiniana* a cosmopolitan species, resistant to extreme pollution with wide range of distribution including eutrophic, electrolyte rich water, accounts for more than 90% of 10 species in Singanallur wetland. Perur wetland with 28 species has *Cymbella turgida*, *Gomphonema parvulum*, *Nitzschia clausii* and *N. obtuse* as dominant species. *Gomphonema parvulum* and *Nitzschia sp.* survive even extreme pollution in wetlands where as *Cymbella turgida* thrive in mesotrophic to eutrophic condition. The assemblages of Noyyal river is similar as of Perur wetland, where this site is represented by *Aulocosira granulata*, *Craticula ambigua*, *Gomphonema parvulum* and *Nitzschia sp.* more in number. *Aulocosira granulata* and *Craticula ambigua* thrive in mesotrophic to eutrophic conditions and *Gomphonema* and *Nitzschia sp.* are capable of surviving even in extreme conditions of Pollution.

A wide range of diatoms distribution is observed in all sampled wetlands of Coimbatore, which include *Gomphonema sp.* and *Nitzschia sp.* *Aulocosira granulata*, *Cyclotella meneghiniana* and *Sellaphora pupula* were dominant in Noyyal River, Pallapalayam and

Sundakamuthur wetlands. These wetlands receives untreated sewage and are either eutrophic to mesotrophic evident from diatom assemblages.

Diatom Indices

Variants of diatom indices have been used across the globe. Table 2 lists most commonly used diatom indices for representing the degree of pollution suitable for tropical conditions. Diatom indices listed in Table 3 were computed for all sampled wetlands to evaluate water quality. This diatom index score is expressed as water quality optima (i.e. the tolerate limits of diatoms to water quality variables) of the sample, based on the diatom taxa 'i' weighted by the abundance of each taxon.

Table 5 Diatom Indices values for the wetlands

SITES	IPS	SLAD	DESCY	L&M	SHE	WAT	EPI-D	ROTT	IDG	CEE	IBD	IDAP	TDI
VP	7.7	13.2	17.3	11.1	14	8.5	10.9	16.3	12.8	11.6	1	11.6	84.3
PP	7.3	10.3	11.3	9.6	13.4	10.6	7.6	8.3	10.1	4.6	6.1	7.2	92.5
SM	9.3	10.6	9.8	9.3	13	6.2	8.1	11.3	10	8.4	8.1	7.2	76
PP	7.6	10.1	11.1	9.3	13.4	10.2	7.9	12.7	11.9	3.7	6.6	7.2	90.3
SN	5.9	7.8	10.4	8.2	8.9	1.7	8.1	NA	13.5	3.3	6.5	5.8	99.9
PR	13.5	10.1	11.7	9.1	6.1	13.6	8.3	10.7	12.3	6.3	15.8	6.7	77.9
SM	9.8	10.5	9.9	8.5	8.6	10.7	8.3	11	12.2	8.2	7.7	4.9	86.3
NL	8	9.2	9.8	7.8	9.6	10.9	8.9	3.8	6	5.2	3.9	7.2	81.3

Note: VP- Vedapatti wetland; PP- Pallapalayam wetland; SM- Sundakamuthur wetland; PP-Pallapalayam wetland; SN-Singanallur wetland; PP-Perur wetland ; SM- Sundakamuthur wetland; NL-Noyyal River. Refer Table 2 for details about the diatom indices

IPS and GDI Indices attributing to trophic status are listed in Table 6 (adopted from Eloranta & Soininen, 2002, Taylor, 2004). Based on this, scores listed in Table 5, indicate an increasing level of pollution or eutrophication.

Table 6 Class limit values for Diatom indices (Eloranta & Soininen, 2002)

Index score	Class	Trophy
>17	High quality	Oligotrophy
15 to 17	Good quality	oligo-mesotrophy
12 to 15	Moderate quality	Mesotrophy
9 to 12	Poor quality	meso-eutrophy
<9	Bad quality	Eutrophy

Diatom assemblages along with water quality class and trophic conditions of the wetlands are listed in Table 7.

Table 7 Trophic condition of the wetlands with dominant species

Site name	Dominant Species	Substrata	Class	Water quality	Trophic conditions
Vedapatti wetland	<i>Diadsmis confervaceae</i> , <i>Gomphonema turris</i> , <i>G. gracile</i>	Aquatic plant	3-4	Moderate to poor quality	Meso-eutrophic to mesotrophic
Pallapalayam wetland	<i>Aulocosiera granulata</i> , <i>Nitzschia</i> sp., <i>Cyclotella meneghiniana</i>	Stone	3-5	Moderate to bad quality	Mesotrophic to eutrophic
Sundakamuthur wetland	<i>Sellaphora pupula</i> , <i>Navicula rostellata</i>	Sediment	4-5	Bad quality	Eutrophic
Pallapalayam wetland	<i>Cyclotella meneghiniana</i> ,	Aquatic plant	3-5	Moderate to bad	Mesotrophic to Eutrophic

	<i>Aulocosira</i>			quality	
	<i>granulata</i>				
Singanallur wetland	<i>Cyclotella meneghiniana</i>	Aquatic plant	5	Bad quality	Eutrophic
Sundakamuthur wetland	<i>Sellaphora pupula</i> , <i>Gomphonema parvulum</i> , <i>Gomphonema sp.</i>	Aquatic plant	4-5	Bad to poor quality	Eutrophic
Perur wetland	<i>Gomphonema parvulum</i> , <i>Cymbella turgida</i> , <i>Nitzschia obtuse</i> , <i>Nitzschia clausii</i>	Aquatic plant	4	Moderate to Poor quality	Meso-eutrophic
Noyyal River	<i>Nitzschia sp.</i> <i>Navicula sp.</i>	Aquatic plant	4-5	Bad to poor quality	Meso-eutrophic

Habitat preference

Diatom community structure varied very distinctly across the habitats. Epiphytic, Epilithic and Episammic habitats contained 50%, 10.4%, and 7.2% of taxa unique to that habitat. In all these habitats, *Gomphonema affine*, *G.parvulum*, *Aulocosira granulata* and *Navicula roetellata* were common, while *G. parvulum* and *A. granulata* were abundant.

Table 8 lists species with their habitats, shows that majority of the diatom species are epiphytes. Diatoms specific to epilithic habitats are *Fragilaria ungeriana*, *Thalassiosira duostra*, *Navicula anthracis*, *Eolimna subminuscula*, *Amphora veneta*, *Navicula veneta* and *Nitzschia sigma*. Epilithic habitat supports both centric and pennate diatoms. Episammic habitat supported 10 species which includes *Navicula viridula*, *Aulacoseira muzzanensis*,

Gomphonema pseudoaugar, *Hantzschia* sp., *Anomoeoneis sphaerophora*, *Pinnularia microstauron*, *P.graciloides*, *P.interrupta*, *Caloneis bacillum* and *Rhopalodia* sp.

Cyclotella meneghiniana and *Nitzschia obtuse* were most abundant and specific to epiphytic and epilithic substrata. Similarly, species with average dominance were restricted to only epiphytic and episammic habitats. However, diatom community specific to both epilithic and episammic were absent.

Table 8 Species list with their occurrence in three habitats

Species	Epiphytic	Epilithic	Episammic
<i>Gomphonema affine</i> Kutzing	+	+	+
<i>Gomphonema parvulum</i> Kutzing var. <i>parvulum</i>	+	+	+
<i>Aulacoseira granulata</i> (Ehr.) Simonsen	+	+	+
<i>Navicula rostellata</i> Kutzing	+	+	+
<i>Cyclotella meneghiniana</i> Kutzing	+	+	-
<i>Craticula accomoda</i> (Hustedt) Mann	+	+	-
<i>Nitzschia obtusa</i> W.M.Smith	+	+	-
<i>Nitzschia frustulum</i> (Kutzing) Grunow var. <i>frustulum</i>	+	+	-
<i>Eunotia mesiana</i> Chohnoky	+	+	-
<i>Fragilaria biceps</i> (Kutzing) Lange- Bertalot	+	+	-
<i>Navicula erifuga</i> Lange-Bertalot	+	+	-
<i>Fragilaria ulna</i> var. <i>acus</i> (Kutz.)Lange- Bertalot	+	+	-
<i>Nitzschia</i> sp.	+	+	-
<i>Seminavis</i> D.G. Mann	+	+	-
<i>Navicula symmetrica</i> Patrick	+	+	-

<i>Tryblionella calida</i> (Grunow in Cl. & Grun.	+	-	+
<i>Gomphonema</i> sp.	+	-	+
<i>Sellaphora laevis</i> (Kutzing) D.G. Mann	+	-	+
<i>Fallacia pygmaea</i> (Kützing) Stickle & Mann	+	-	+
<i>Surirella tenera</i> Gregory	+	-	+
<i>Sellaphora pupula</i> (Kutzing) Mereschkowksy	+	-	+
<i>Luticola acidoclinata</i> Lange-Bertalot	+	-	+
<i>Pinnularia acrospheria</i> Rabenhorst	+	-	+
<i>Nupela</i> sp.	+	-	+
<i>Nitzschia palea</i> (Kutzing) W.Smith	+	-	+
<i>Placoneis</i> sp.	+	-	+
<i>Navicula gregaria</i> Donkin	+	-	+
<i>Pinnularia</i> sp.	+	-	+
<i>Craticula ambigua</i> (Ehrenberg) Mann	+	-	+
<i>Amphora copulata</i> (Kutz) Schoeman & Archibald	+	-	+
<i>Caloneis molaris</i> (Grunow) Krammer	+	-	-
<i>Nitzschia umbonata</i> (Ehrenberg) Lange-Bertalot	+	-	-
<i>Navicula trivialis</i> Lange-Bertalot var. <i>trivialis</i>	+	-	-
<i>Aulacoseira ambigua</i> (Grun.) Simonsen	+	-	-
<i>Bacillaria paradoxa</i> Gmelin	+	-	-
<i>Navicula zannoni</i> Hustedt	+	-	-

<i>Nitzschia pumila</i> Hustedt	+	-	-
<i>Craticula</i> sp.	+	-	-
<i>Cymbella turgidula</i> Grunow in A. Schmidt & al.	+	-	-
<i>Navicula germainii</i> Wallace	+	-	-
<i>Cocconeis</i> sp.	+	-	-
<i>Cocconeis placentula</i> Ehrenberg var. <i>placentula</i>	+	-	-
<i>Nitzschia liebetruthii</i> Rabenhorst var. <i>liebetruthii</i>	+	-	-
<i>Surirella angusta</i> Kutzing	+	-	-
<i>Rhopalodia gibba</i> (Ehr.) O.Muller var. <i>gibba</i>	+	-	-
<i>Pinnularia viridiformis</i> Krammer	+	-	-
<i>Surirella</i> sp.	+	-	-
<i>Amphora montana</i> Krasske	+	-	-
<i>Actinocyclus normanii</i> (Greg. ex Grev.) Hustedt	+	-	-
<i>Pleurosigma salinarum</i> (Grunow) Cleve & Grunow	+	-	-
<i>Aulacoseira distans</i> (Ehr.) Simonsen	+	-	-
<i>Pinnularia species</i>	+	-	-
<i>Nitzschia supralitorea</i> Lange-Bertalot	+	-	-
<i>Planothidium rostratum</i> (Oestrup) Lange-Bertalot	+	-	-
<i>Planothidium robustum</i> (Hustedt) Lange-Bertalot	+	-	-
<i>Gomphonema turris</i> Ehr.	+	-	-

<i>Gomphonema gracile</i> Ehrenberg	+	-	-
<i>Geissleria decussis</i> (Ostrup) Lange-Bertalot	+	-	-
<i>Diploneis ovalis</i> (Hilse) Cleve	+	-	-
<i>Cyclotella woltereckii</i> Hustedt	+	-	-
<i>Diadomesis confervacea</i> Kützing	+	-	-
<i>Gomphonema species</i>	+	-	-
<i>Encyonema mesianum</i> (Cholnoky) D.G. Mann	+	-	-
<i>Eunotia sp.</i>	+	-	-
<i>Eolimna sp.</i>	+	-	-
<i>Fragilaria ulna</i> (Nitzsch.) Lange-Bertalot	+	-	-
<i>Diploneis puella</i> (Schumann) Cleve	+	-	-
<i>Encyonema minutum</i> (Hilse in Rabh.) D.G. Mann	+	-	-
<i>Eunotia minor</i> (Kützing) Grunow in Van Heurck	+	-	-
<i>Lemnicola hungarica</i> (Grunow) Round & Basson	+	-	-
<i>Cymbella turgida</i> Gregory	+	-	-
<i>Nitzschia clausii</i> Hantzsch	+	-	-
<i>Nitzschia amphibia</i> Grunow f.amphibia	+	-	-
<i>Navicula sp.</i>	+	-	-
<i>Cymbella tumida</i> (Brebisson)Van Heurck	+	-	-
<i>Placoneis sp.</i>	+	-	-
<i>Navicula sp.</i>	+	-	-
<i>Nitzschia capitellata</i> Hustedt in A.Schmidt & al.	+	-	-

<i>Fragilaria ungeriana</i> Grunow	-	+	-
<i>Thalassiosira duostra</i> Pienaar	-	+	-
<i>Navicula anthracis</i> Cleve et Brun	-	+	-
<i>Eolimna subminuscula</i> (Manguin) Moser	-	+	-
Lange- -Bertalot & Metzeltin			
<i>Amphora veneta</i> Kutzing	-	+	-
<i>Navicula veneta</i> Kutzing	-	+	-
<i>Nitzschia sigma</i> (Kutzing)W.M.Smith	-	+	-
<i>Navicula viridula</i> (Kutzing) Ehrenberg	-	-	+
<i>Aulacoseira muzzanensis</i> (Meister)	-	-	+
Krammer			
<i>Gomphonema pseudoaugur</i> Lange- Bertalot	-	-	+
<i>Hantzschia</i> sp.	-	-	+
<i>Anomoeoneis sphaerophora</i> (Ehr.) Pfitzer	-	-	+
<i>Caloneis bacillum</i> (Grunow) Cleve	-	-	+
<i>Pinnularia microstauron</i> (Ehr.) Cleve	-	-	+
<i>Pinnularia graciloides</i> Hustedt	-	-	+
<i>Pinnularia interrupta</i> W.M.Smith	-	-	+
<i>Rhopalodia</i> sp.	-	-	+

Conclusion

Samples collected from six wetlands of Coimbatore, records 27 dominant species of 96 Species belonging to 34 genera. Dominant species that are cosmopolitan include *Cyclotella meneghiniana*, *Nitzschia* sp., *Sellaphora pupula*, *Gomphonema parvulum* and *Navicula* sp. Singanallur wetland and Noyyal river stretches are characterised by pollution tolerant species with low diatom diversity. Diatom assemblages indicate that Vedapatti, Perur and Sundakamuthur wetlands are moderately polluted, while Pallapalayam, Noyyal River and Singanallur wetlands are heavily polluted. In these wetlands distribution of *Cyclotella* was determined by high electrolyte conductivity and *Gomphonema* and *Nitzschia* were distributed in all pH and conductivity ranges, where *Fragilaria* is restricted to neutral alkaline pH and moderate electrolytic waters. With respect to habitat preference epiphytic, epilithic and episammic habitats contained 50%, 10.4%, and 7.2% of taxa unique to that habitat. Diatom indices reveal that water quality of the sampled wetlands are moderate (mesotrophic) to heavily polluted (eutrophic).

Acknowledgement

We thank the Ministry of Environment and Forests, Government of India and Indian Institute of Science for the infrastructure support and sustained financial support for ecological research. We are also grateful to Dr. Jonathan Taylor and Dr. J.P. Kociolek for confirming the identity of species. Dr. P Pramod and Mr. Joseph Reginald from SACON, Coimbatore provided useful tips during field investigations. We thank Institute Nanoscience Initiative (INI) for permitting us to use SEM facility and INI staff for their help and support.

References

1. Atazadeh, I. Sharifi, M. Kelly, M. G., 2007. Evaluation of the Trophic Diatom Index for assessing water Quality in River Gharasou, western Iran. *Hydrobiologia* 589:165–173.
2. Barbour, M.T. Gerritsen, J. and White, J.S., 1996. Development of the Stream Condition Index (SCI) for Florida. A Report to the Florida Department of Environmental Protection, Stormwater and Nonpoint Source Management Section, Tetra Tech, Inc., Owing Mills, Maryland, USA .
3. Berger, W.H. Parker, F.L., 1970. Diversity of planktonic foraminifera in deep-sea.
4. Cemagref, 1982. Etude des methods biologiques d'appré- ciation quantitative de la qualitédes eaux. Rapport Q. E. Lyon, Agence de l'eau Rhône-Me'diterrane'e-Corse-Cemagref, Lyon, France.
5. Cholnoky, B. J., 1968. Die Ökologie der Diatomeen in Binnengewässern, p. 699. J. Cramer, Braunschweig.
6. Coste, M. and Ayphassorho, H., 1991. Étude de la qualité deseaux du Bassin Artois-Picardie àl'aide des communautés de diatomées benthiques (application des indices diatomiques). Rapport Cemagref. Bordeaux-Agence de l'Eau Artois-Picardie, Douai.
7. Cox, E. J, 1991. What is the basis for using diatoms as monitors of river quality? In B .A. Whitton, E. Rott & G.Friedrich (eds), *Use of algae for monitoring rivers*. E. Rott, Innsbruck: 33-40.
8. Dakshini, K. M. M. and Soni, J. K., 2006. Diatom distribution and status of organic pollution in sewage drains. *Hydrobiologia* 87, 205-209.
9. Dell'Uomo, A., 1996. Assessment of water quality of an Apennine river as a pilot study. In Whitton, B. A. & E. Rott (eds), *Use of Algae for Monitoring Rivers II*. Institut fu" r Botanik, Universität Innsbruck, 65–73.

10. Descy, J.P. and Coste, M., 1991. A test of methods for assessing water quality based on diatoms. *Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie* 24: 2112–2116.
11. Directory of Indian Wetlands, 1993. Compiled by the World Wide Fund for Nature (WWF), India in collaboration with the Asian Wetland Bureau.
12. Dixit, S.S. Smol, J.P. Kingston, J.C. and Charles, D.F., 1992. Diatoms: Powerful indicators of environmental change. *Environmental Science and Technology*. 26 23-33.
13. Fore, L. S. and Grafe, C., 2002. Using diatoms to assess the biological condition of large rivers in Idaho (U.S.A.). *Freshwater Biology*. 47, 2015–2037.
14. Galatowitsch¹, S. M. Whited, D.C. and Tester, J.R., 1999. Development of community metrics to evaluate recovery of Minnesota wetlands. *Journal of Aquatic Ecosystem Stress and Recovery* 6: 217–234, 1999.
15. Gandhi, H.P., 1957. A contribution to our knowledge of the diatom genus *Pinnularia*. *Journal of the Bombay Natural History Society* 54: 845–853.
16. Gandhi, H.P., 1959. Freshwater diatoms from Sagar in the Mysore State. *Journal of the Indian Botanical Society* 38: 305–331.
17. Gandhi, H.P., 1959. Notes on the Diatomaceae from Ahmedabad and its environs-II. On the diatom flora of fountain reservoirs of the Victoria Gardens. *Hydrobiologia* 14: 130–146.
18. Gandhi, H.P., 1961. Notes on the Diatomaceae of Ahmedabad and its environs. *Hydrobiologia* 17: 218–236.
19. Gandhi, H.P., 1962. Notes on the Diatomaceae from Ahmedabad and its environs-IV -The diatom communities of some freshwater pools and ditches along Sarkhej Road. *Phykos* 1: 115–127.
20. Gandhi, H.P., 1964. The diatom flora of Chandola and Kankaria Lakes. *Nova Hedwigia* 8: 347–402.

21. Gandhi, H.P., 1967. Notes on Diatomaceae from Ahmedabad and its environs. VI. On some diatoms from fountain reservoirs of Seth Sarabhai's Garden. *Hydrobiologia* 30: 248–272.
22. Gandhi, H. P., 1998. Freshwater Diatoms of Central Gujarat. Bishen Singh Mahendra Pal Singh. Dehra Dun.
23. Gernes, M.C. and Helgen, J.C., 1999. Indexes of biotic integrity for wetlands, section B: wetland vegetation IBI for depressional wetlands. Final Report to the United States Environmental Protection Agency Assistance Number CD995525-01, April 1999. Minnesota Pollution Control Agency, St. Paul, Minnesota, USA.
24. Gómez, N. and Licursi, M., 2001. The Pampean Diatom Index (IDP) for assessment of rivers and streams in Argentina. *Aquatic Ecology* 35: 173–181.
25. Gopal, B., 2005. Does inland aquatic biodiversity have a future in Asian developing countries?. *Hydrobiologia* 542:69–75.
26. Gosselink, J.C. and Turner, R.E., 1978. The role of hydrology in freshwater wetland ecosystems. Pp. 63-78 in freshwater wetlands. Ecological processes and management potential. New York. Academic press.
27. John, J., 1993. The use of diatoms in monitoring the development of created wetlands at a sandmining site in Western Australia. *Hydrobiologia* 269/270: 427-436.
28. Karthick, B. Krithika, H. and Alakananda, B., 2008. Short Guide to common freshwater Diatom Genera (Poster). Energy and Wetlands Research Group, CES, IISc, Bangalore.
29. Kelly, M. G. and Whitton, B. A., 1995. The Trophic Diatom Index: a new index for monitoring Eutrophication in rivers. *Journal of Applied Phycology* 7:433-444.
30. Kelly, M. G. Cazaubon, A. Coring, E. Dell'Uomo, A. Ector, L. Goldsmith, B. Guasch, H. Hürlimann, J. Jarlman, A. Kawecka, B. Kwandrans, J. Laugaste, R. Lindstrøm, E.A. Leitao, M. Marvan, P. Padisák, J. Pipp, E. Prygiel, J. Rott, E. Sabater, S. Dam, V .H. and Vizinet, J., 1998. Recommendations for the routine

- sampling of diatoms for water quality assessments in Europe. *Journal of Applied Phycology* 10: 215-224, 1998.
31. Kelly, M. G. Penny, C. J. And Whitton, B. A., 1995. Comparative performance of benthic diatom indices used to assess river water quality. *Hydrobiologia* 302: 179-188.
 32. Kerans, B.L. and Karr, J.R., 1994. A benthic index of biotic integrity (B-IBI) for rivers in the Tennessee valley, *Ecol. Appl.* 4 (4), pp. 768–785.
 33. Kolkwitz, R. and Marsson, M., 1908. *Ökologie der pflanzliche Saprobien*. *Ber. Deutsche Bot. Gesellsch.* 26: 505-5019.
 34. Lavoie, I. Somers, K .M. Paterson, A. M. and Dillon, P .J., 2000. Assessing scales of variability in benthic diatom community structure. *Journal of Applied Phycology* 17: 509–513.
 35. Leclercq, L. and Maquet, B., 1987. Deux nouveaux indices chimique et diatomique de qualite´ d’eau courante. Application au Samson et à ses affluents (bassin de la Meuse belge). Comparaison avec d’autres indices chimiques, bioce´notiques et diatomiques. Institut Royal des Sciences Naturelles de Belgique, document de travail 28.
 36. Lecoingte, C., Coste M. and Prygiel, J., 1993. “Omnidia”: Software for taxonomy, calculation of diatom indices and inventories management. *Hydrobiology* 269/270: 509–513.
 37. Legendre, P. and Legendre, L., 1998. *Numerical Ecology*. 2nd English edition. Elsevier, Amsterdam.
 38. Lenoir, A. and Coste, M., 1996. Development of a practical diatom index of overall water quality applicable to the French National Water Board network. In Whitton, B. A. & E. Rott (eds), *Use of Algae for Monitoring Rivers II*. Institut für Botanik. Universität Innsbruck, 29–43.
 39. Lowe, R. L., 1974. Environmental requirements and pollution tolerance of freshwater diatoms. *Environmental Monitoring Series*, National Environmental Research Center, Cincinnati, Ohio.

40. Ludwig, John A. and Reynolds, J.F., 1988. Statistical ecology: a primer of methods and computing. Wiley Press, New York, New York. 337 pp.
41. Micacchion, M., 2004. Integrated Wetland Assessment Program. Part 7: Amphibian Index of Biotic Integrity (AmphIBI) for Ohio Wetlands. Ohio EPA Technical Report WET/2004-7. Ohio Environmental Protection Agency, Wetland Ecology Group, Division of Surface Water, Columbus, Ohio.
42. Mitsch, W.J. and Gosselink, J.G., 1986. Wetlands. Van Nostrand Reinhold. New York. Pp.539.
43. O'Connell, T. J. Jackson, L. E. Brooks, R. P., 2008. A bird community index of biotic integrity for the Mid-Atlantic Highlands. Environmental monitoring and Assessment 51:145-156.
44. Panini, D., 2007. The Ramsar Convention and National Laws and Policies for Wetlands in India. Technical Consultation on Designing Methodologies to Review Laws and Institutions Relevant to Wetlands.
45. Patrick, R., 1986. Diatoms as indicators of changes in water quality. In M. Ricard (ed.), Proceedings of the 8th International Diatom Symposium. Koeltz Scientific Books, Koenigstein: 759-766.
46. Prasad, S.N. Sengupta, T. Alok Kumar Vijayan VS, Vijayan, L. Ramachandra, T.V. Ahalya, N. And Tiwari, A.K., 2003. Wetlands of India. Natural Aquatic Ecosystems of India. 6-25.
47. Prygiel, J. and Coste, M., 1993. The assessment of water quality in the Artois-Picardie water basin (France) by the use of diatom indices. Hydrobiologia. 269/279 343-349.
48. Prygiel, J. Leveque, L. and Iserentant, R., 1996. Un nouvel indice diatomique pratique pour l'évaluation de la qualité des eaux en réseau de surveillance. Rev. Sci. Eau 1: 97-113.
49. Ramachandra.T.V., Kiran Rajashekariah and Ahalya.N., 2002. Conservation and Mangement of Wetlands, Allied Publishers, Mumbai, India

50. Reiss, K.C., 2006 Florida Wetland Condition Index for depression forested wetlands. *Ecological Indicators*. Volume 6, Issue 2, Pages 337-352.
51. Richards, C., L.B. Johnson, and Host, G.E., 1996. Landscape scale influences on stream habitats and biota. *Canadian Journal of Fisheries and Aquatic Science*. 53 (suppl. 1). 295-311.
52. Rott, E., 1991. Methodological aspects and perspectives in the use of periphyton for monitoring and protecting rivers. In Whitton, B. A., E. Rott & G. Friedrich (eds), *Use of Algae for Monitoring Rivers*. Institut für Botanik, Univ. Innsbruck, 9-16.
53. Round, F. E., 1991. Diatoms in River water monitoring studies. *Journal of Applied Phycology* 3: 129-145.
54. Schiefele, S. and Schreiner, C., 1991. Use of diatoms for monitoring nutrient enrichment acidification and impact salts in Germany and Austria. In Whitton, B. A., E. Rott and G. Friedrich (eds), *Use of Algae for Monitoring Rivers*. Institut für Botanik, Univ. Innsbruck.
55. Schoeman, F. R. and Archibald, R. E. M., 1976-1980. *The Diatom Flora of Southern Africa*. National Institute for Water Research, Pretoria.
56. Schulz, E.J. Hoyer, M.V. Canfield, D.E. 1999. An index of Biotic Integrity: A Test with Limnological and Fish Data from Sixty Florida Lakes. *Transactions of the American Fisheries Society* 128:564-577.
57. Sládeček, V., 1986. Diatoms as indicators of organic pollution. *Acta Hydrochimica et Hydrobiologica* 14: 555-566.
58. Soininen, J. Könönen, K., 2004. Comparative study of monitoring South-Finnish Rivers and streams using macroinvertebrate and benthic diatom community structure. *Aquatic ecology* 38: 63-75.
59. Springe, G. Sandin, L. Briede, A. and Skuja, A., 2006. Biological quality metrics: their variability and appropriate scale for assessing streams. *Hydrobiologia* 566:153-172.

60. Stoermer, E.F. and Smol, J.P., 2001. The Diatoms: Applications for the Environmental and Earth Science. Cambridge University Press. Cambridge
61. Taylor, J.C., Harding W. R and Archibald, C.G.M., 2007. An Illustrated Guide to Some Common Diatom Species from South Africa. WRC Report TT 282/07. Water Research Commission. Pretoria
62. Watanabe, T. Asai, K. and Houki, A., 1986. Numerical estimation of organic pollution of flowing waters by using the epilithic diatom assemblage – Diatom Assemblage Index (DIApo). Science of the Total Environment 55: 209–218.
63. Winter J.G. Duthie H.C., 2000. Stream epilithic, epipelic and epiphytic diatoms: habitat fidelity and use in biomonitoring. Aquatic Ecology 34: 345-353.
64. Wu, J. A., 1999. Generic index of diatom assemblages as bioindicator of pollution in the Keelung River of Taiwan. Hydrobiologia 397: 79–87.
65. Zelinka, M. and Marvan, P., 1961. Zur Präzisierung der biologischen Klassifikation der Reinheit fliessender Gewässer. Arch.. Hydrobiol. 57 389-407.

Annexure: I - List of Species

<i>Achnanthidium exiguum</i> (Grunow) Czarn.	AEHE
<i>Actinocyclus normanii</i> (Greg. ex Grev.) Hustedt morphotype <i>normanii</i>	ANMN
<i>Amphora copulata</i> (Kutz) Schoeman & Archibald	ACOP
<i>Amphora montana</i> Krasske	AMMO
<i>Amphora veneta</i> Kutzing	AVEN
<i>Anomoeoneis sphaerophora</i> (Ehr.) Pfitzer	ASPH
<i>Aulacoseira ambigua</i> (Grun.) Simonsen	AAMB
<i>Aulacoseira distans</i> (Ehr.) Simonsen	AUDI
<i>Aulacoseira granulata</i> (Ehr.) Simonsen	AUGR
<i>Aulacoseira muzzanensis</i> (Meister) Krammer	AMUZ
<i>Bacillaria paradoxa</i> Gmelin	BPAR
<i>Caloneis bacillum</i> (Grunow) Cleve	CBAC
<i>Caloneis molaris</i> (Grunow) Krammer	CMOL
<i>Carticula</i> sp.	CRAT
<i>Cocconeis placentula</i> Ehrenberg var. <i>placentula</i>	CPLA
<i>Cocconeis</i> sp.	COCS
<i>Craticula accomoda</i> (Hustedt) Mann	CRAC
<i>Craticula ambigua</i> (Ehrenberg) Mann	CAMB
<i>Cyclotella meneghiniana</i> Kutzing	CMEN
<i>Cyclotella woltereckii</i> Hustedt	CWOL
<i>Cymbella tumida</i> (Brebisson) Van Heurck	CTUM
<i>Cymbella turgida</i> Gregory	CTUR
<i>Cymbella turgidula</i> Grunow in A.Schmidt & al.	CTGL
<i>Diadsmis confervaceae</i> Kützing	DCOF
<i>Diploneis ovalis</i> (Hilse) Cleve	DOVA
<i>Diploneis puella</i> (Schumann) Cleve	DPUE
<i>Encyonema mesianum</i> (Cholnoky) D.G. Mann	ENME

<i>Encyonema minutum</i> (Hilse in Rabh.) D.G. Mann	ENMI
<i>Eolimna</i> sp.	EOLI
<i>Eolimna subminuscula</i> (Manguin) Moser Lange-Bertalot & Metzeltin	ESBM
<i>Eunotia mesiana</i> Cholnoky	EMES
<i>Eunotia minor</i> (Kutzing) Grunow in Van Heurck	EMIN
<i>Eunotia</i> sp.	EUNO
<i>Fallacia pygmaea</i> (Kützing) Stickle & Mann	FPYG
<i>Fragilaria biceps</i> (Kutzing) Lange-Bertalot	FBCP
<i>Fragilaria ulna</i> (Nitzsch.) Lange-Bertalot <i>var. ulna</i>	FULN
<i>Fragilaria ulna</i> <i>var. acus</i> (Kutz.) Lange-Bertalot fo. <i>teratogene</i>	FUAT
<i>Fragilaria ungeriana</i> Grunow	FUNG
<i>Geissleria decussis</i> (Ostrup) Lange-Bertalot & Metzeltin	GDEC
<i>Gomphonema affine</i> Kutzing	GAFF
<i>Gomphonema gracile</i> Ehrenberg	GGRA
<i>Gomphonema parvulum</i> Kutzing <i>var. parvulum f. parvulum</i>	GPAR
<i>Gomphonema pseudoaugur</i> Lange-Bertalot	GPSA
<i>Gomphonema</i> sp.1	GOMS
<i>Gomphonema</i> sp.2	GOMS
<i>Gomphonema turris</i> Ehr.	GTUR
<i>Hantzschia</i> sp.1	HAN1
<i>Lemnicola hungarica</i> (Grunow) Round & Basson	LHUN
<i>Luticola acidoclinata</i> Lange-Bertalot	LACD
<i>Navicula anthracis</i> Cleve et Brun	NANT
<i>Navicula erifuga</i> Lange-Bertalot	NERI
<i>Navicula germainii</i> Wallace	NGER
<i>Navicula gregaria</i> Donkin	NGRE
<i>Navicula rostellata</i> Kutzing	NROS
<i>Navicula</i> sp.1	NASP

<i>Navicula</i> sp.2	NAVI
<i>Navicula symmetrica</i> Patrick	NSYM
<i>Navicula trivialis</i> Lange-Bertalot <i>var. trivialis</i>	NTRV
<i>Navicula veneta</i> Kutzing	NVEN
<i>Navicula viridula</i> (Kutzing) Ehrenberg	NVIR
<i>Navicula zanonii</i> Hustedt	NZAN
<i>Nitzschia amphibia</i> Grunow f.amphibia	NAMP
<i>Nitzschia capitellata</i> Hustedt in A.Schmidt & al.	NCPL
<i>Nitzschia clausii</i> Hantzsch	NCLA
<i>Nitzschia frustulum</i> (Kutzing) Grunow <i>var. frustulum</i>	NIFR
<i>Nitzschia liebetruthii</i> Rabenhorst <i>var. liebetruthii</i>	NLBT
<i>Nitzschia obtusa</i> W.M.Smith	NOBT
<i>Nitzschia palea</i> (Kutzing) W.Smith	NPAL
<i>Nitzschia pumila</i> Hustedt	NPML
<i>Nitzschia sigma</i> (Kutzing)W.M.Smith	NSIG
<i>Nitzschia</i> sp.	NZSS
<i>Nitzschia supralitorea</i> Lange-Bertalot	NZSU
<i>Nitzschia umbonata</i> (Ehrenberg)Lange-Bertalot	NUMB
<i>Nupela</i> sp.	NUPE
<i>Pinnularia acrospheria</i> Rabenhorst	PACR
<i>Pinnularia graciloides</i> Hustedt	PGRO
<i>Pinnularia interrupta</i> W.M.Smith	PINT
<i>Pinnularia microstauron</i> (Ehr.) Cleve	PMIC
<i>Pinnularia</i> sp.	PINS
<i>Pinnularia</i> sp.1	PIN1
<i>Pinnularia viridiformis</i> Krammer	PVIF
<i>Placonesi</i> sp.1	PLAS
<i>Placonesi</i> sp.2	PLAS

<i>Planothidium robustum</i> (Hustedt) Lange-Bertalot	PLRO
<i>Planothidium rostratum</i> (Oestrup) Lange-Bertalot	PRST
<i>Pleurosigma salinarum</i> (Grunow) Cleve & Grunow	PSAL
<i>Rhopalodia gibba</i> (Ehr.) O.Muller var.gibba	RGIB
<i>Rhopalodia</i> sp.	RHOS
<i>Sellaphora laevissima</i> (Kutzing) D.G. Mann	SELA
<i>Sellaphora pupula</i> (Kutzing) Mereschkowksy	SPUP
<i>Seminavis</i> sp.	SMNA
<i>Surirella angusta</i> Kutzing	SANG
<i>Surirella</i> sp.	SURS
<i>Surirella tenera</i> Gregory	SUTE
<i>Thalassiosira duostra</i> Pienaar	TDUO
<i>Tryblionella calida</i> (grunow in Cl. & Grun.) D.G. Mann	TCAL

Annexure: II - Illustrations

This report is based on one time sampling and may not reflect the entire diatom flora of Coimbatore. This report is written keeping in mind the requirement of beginner's who have started the journey with diatoms. It is important not to adopt "nearest match" approaches in identification of diatom flora. Photographs included here were taken using a camera attached with bright field microscope (scale bars are equal to 10 μm) and some are using scanning Electron Microscope (SEM) available at **Institute Nanoscience Initiative (INI), Indian Institute of Science**. Identification of diatom taxa and ecological information provided in this report are based on the following literatures:


1. Cox, E.J., 1996 Identification of freshwater Diatoms from Live material. Chapman & Hall. London.UK.
2. Krammer, K. and Lange-Bertalot, H., 1986. Süßwasserflora von Mitteleuropa. Band 2. *Bacillariophyceae*. Teil 1. *Naviculaceae*. Gustav Fisher Verlag, Stuttgart. Germany.
3. Krammer, K. and Lange-Bertalot, H., 1988. Süßwasserflora von Mitteleuropa. Band 2. *Bacillariophyceae*. Teil 2. *Bacillariaceae, Epithemiaceae, Surirellaceae*. Gustav Fisher Verlag, Stuttgart. Germany.
4. Krammer, K. and Lange-Bertalot, H., 1991. Süßwasserflora von Mitteleuropa. Band 2. *Bacillariophyceae*. Teil 3. *Centrales, Fragilariaceae, Eunotiaceae*. Gustav Fisher Verlag, Stuttgart. Germany.
5. Krammer, K. and Lange-Bertalot, H., 1991. Süßwasserflora von Mitteleuropa. Band 2. *Bacillariophyceae*. Teil 4. *Achnanthaceae, Kritische Ergänzungen zu Navicula (Lineolatae) and Gomphonema*. Gustav Fisher Verlag, Stuttgart. Germany.
6. Krammer, K., 2000. The genus *Pinnularia*. Diatoms of Europe, Volume 1. Edited by H. Lange-Bertalot. A.R.G. Gantner verlag K.G.Germany.
7. Krammer, K., 2002. *Cymbella*. Diatoms of Europe, Volume 3. Edited by H. Lange-Bertalot. A.R.G. Gantner verlag K.G.Germany.

8. Krammer, K., 2003. *Cymboplectra, Delicata, Navicymbella, Gomphocymbellipsis, Afrocyymbella*. Diatoms of Europe, Volume 1. Edited by H. Lange-Bertalot. A.R.G. Gantner verlag K.G.Germany.
9. Lange-Bertalot H., 2001. The genus *Navicula* sensu stricto 10 Genera Separated from *Navicula* sensu lato *Frustulia*. Diatoms of Europe, Volume 2. Edited by H. Lange-Bertalot. A.R.G. Gantner verlag K.G.Germany.
10. Lange-Bertalot, H. and Krammer, K., 1987. Bacillariaceae, Epithemiaceae, Surirellaceae. Bibliotheca Diatomologica 15. J. Cramer, Stuttgart.
11. Mann, D.G. Thomas, S.J. and Evans, K.M., 2008. Revision of the diatom genus *Sellaphora*: a first account of the larger species in the British Isles. Fottea, Olomouc. 8(1): 15-78.
12. Taylor, J.C., Harding W. R and Archibald, C.G.M., 2007. An Illustrated Guide to Some Common Diatom Species from South Africa. WRC Report TT 282/07. Water Research Commission. Pretoria
13. Vuuren, S.J. Taylor, J., Gerber, A. and Ginkel, C., 2006. Easy identification of the most common Freshwater Algae. A guide for the identification of microscopic algae in South African freshwaters.




NOTE: This document does not provide comprehensive list of all diatom species. If particular taxon is not found in this report, researchers are advised to refer diatom floras listed above. If this is not possible, it is appropriate to leave the specimen catalogued as “unidentified” with illustrations or photographs for future references.

PLATE -01

Aulacoseira granulata (Ehrenberg) Simonsen



SEM



Dimensions:

Valve diameter: 5-15.5 μm


Valve mantle depth: 2.5-8 μm

Striae density: 4-7/10 μm

Ecology:

PLATE -02

Aulacoseira muzzanensis (Meister) Krammer



Dimensions:

Valve diameter: 8 μ m


Valve mantle depth: 7 μ m

Striae density: 8-21/10 μ m

Ecology:

A Planktonic and benthic species found in eutrophic waters.

Aulacoseira ambigua (Grunow) Simonsen



Dimensions:

Valve diameter: 8 μ m

Valve mantle depth: μ m






Striae density: μ m

Ecology:

A Planktonic and benthic species found in eutrophic waters.

PLATE -03

Cyclotella meneghiniana Kutzing



SEM

Dimensions:

Valve diameter: 5-28 μ m

Striae density: 6-10/10 μ m

Ecology:

This taxon has a cosmopolitan distribution in the benthos and plankton of eutrophic, electrolyte rich lakes, rivers and streams.

PLATE -04

***Discotella pseudostelligera* (Hustedt) Houk & Klee** **Syn. *Cyclotella pseudostelligera* (Hustedt)**

**Dimensions:**

Valve diameter: 7-8 μm

Striae density: 7/10 μm

Ecology:

Found in freshwater in the plankton of inland rivers and lakes.

***Discotella woltrerekii* (Hustedt) Houk & Klee** **Syn. *Cyclotella woltrerekii* Hustedt**

**Dimensions:**

Valve diameter: 6-6.5 μm


Valve mantle depth: 4-8 μm

Striae density: 8-21/10 μm

Ecology:

Found in freshwater in the plankton of inland rivers and lakes.

***Thalassiosira duostra* Pienaar**


**Dimensions:**

Valve diameter: 10-11 μm

Ecology: Unknown

PLATE -05

Actinocyclus normanii Ehrenberg



Dimensions:


Valve diameter: 22-27 μm

Ecology: Planktonic, freshwater taxa.

Ecology Unknown

PLATE -06

Fragilaria ulna (Nitzsch) Lang-Bertalot
Syn. *Synedra ulna* (Nitzsch) Ehrenberg



SEM

Dimensions:

Valve length: 100-365 μm

Valve breadth: 7-8.5 μm

Striae density: 8-10/10 μm



Ecology:

This cosmopolitan taxon is found in the benthos of rivers and lakes and is easily suspended in the plankton due to its relatively large surface area. Often found in mesotrophic to eutrophic, alkaline waters.

PLATE -07

***Fragilaria ulna* var. *acus* (Kutzing) Lang-Bertalot**

Syn. *Synedra ulna* Kutzing



Dimensions:

Valve length: 140-210 μm

Valve breadth: 5.5-7 μm


Striae density: 8-10/10 μm

Ecology:

This cosmopolitan taxon is found in the benthos of rivers and lakes and is easily suspended in the plankton due to its relatively large surface area. Found in mesotrophic to eutrophic, alkaline freshwaters. Living cells are usually apically attached to a substratum.

PLATE -08

***Fragilaria biceps* (Kutzing) Lang-Bertalot**
Syn. *Synedra ulna* v *biceps* (Kutzing) Kirchner in Cohn



Dimensions:

Valve length: 280-320 μm

Valve breadth: 6-7 μm

Striae density: 7-9/10 μm

Ecology:

This cosmopolitan taxon is found in the benthos of rivers and lakes and is easily suspended in the plankton due to its relatively large surface area. Often found in mesotrophic to eutrophic waters together with *F.ulna*. Living cells are usually apically attached to a substratum by a mucilage pad or free living.

PLATE -09

Fragilaria ungeriana Grunow

Dimensions:

Valve length: 75-90 μm


Valve breadth: 8-9 μm

Striae density: 9-10/10 μm

Ecology:

Cells attached face to face. Found in tropical and sub-tropical, weakly alkaline, oxygen-rich waters.

Eunotia minor (Kutzing) Grunow



Dimensions:

Valve length: 50-58 μm

Valve breadth: 8-9 μm

Striae density: 8-9/10 μm

Ecology:

Occurs in circumneutral waters, in pools and springs.



PLATE -10

Eunotia pectinalis v *gibbulosus*

Dimensions:

Valve length: 95-140 μm

Valve breadth: 7.5-9.5 μm

Striae density: 9/10 μm

Ecology:

Unknown

Achnanthes exigua

Dimensions:

Valve length: 7 μm

Valve breadth: 4 μm

Striae density: 24-30/10 μm RV
: 20/10 μm RLV



Ecology:

This cosmopolitan species has very wide ecological amplitude and is found in many different types of water including industrial and other wastewater. It is also able to grow under very low light and can tolerate temperatures of up to 40°C. The optimum growth conditions for this taxa are alkaline water with moderate to elevated electrolyte content.





PLATE -11

***Lemnicola hungarica* (Grunow) Round & Basson** **Syn. *Achnanthes hungarica* Grunow**



Dimensions:

Valve length: 23-25 μm


Valve breadth: 5.5 μm

Striae density: 16-23/10 μm

Ecology:

Occurs in weakly alkaline waters with moderate to elevated electrolyte content and may also occur in critically polluted waters. Found mostly as an epiphyte, commonly found associated with *Lemna* spp. (duckweed).

***Planothidium rostratum* (Oestrup) Round & Bukhityarova** **Syn. *Achnanthes lanceolata* spp. *Frequentissima* var. *rostrata* (Oestrup) Hustedt**



Dimensions:

Valve length: 15-25 μm

Valve breadth: 6.5-7 μm

Striae density: 12-16/10 μm

Ecology:

Occurring in circumneutral to alkaline waters with low to moderate electrolyte content. More often attached to plants than stones.

PLATE -12

Cocconeis placentula Ehrenberg



Dimensions:

Valve length: 13-21/33 μm

Valve breadth: 8-13/18 μm

Straie density: 24-26/10 μm RV
: 20-23/10 μm RLV

Ecology:

Occurring in meso-to eutrophic flowing and standing waters. Found in abundance on plants, wood and stone.

Cocconeis sp.



Dimensions:

Valve length: 33 μm

Valve breadth: 18 μm

Straie density: /10 μm

Ecology:

A very widespread and common genus occurring in all waters. The frustules are epiphytic on filamentous algae or on aquatic plants, sometimes occurring so abundantly as to form a coating over the host surface. Cells of *Cocconeis* may also be found attached to stones and various other substrata.

PLATE -13

Pleurosigma salinarum Grunow

**Dimensions:**

Valve length: 85-88 μm

Valve breadth: 11.5-12.5 μm

Striae density: 22-25/10 μm

Ecology:

Cosmopolitan, found in brackish and saline inland waters.

Diploneis oblongella (Naegeli) Cleve-Euler

Dimensions:

Valve length: 23 μm

Valve breadth: 9-10 μm

Striae density: 10-11/10 μm

Ecology:

Found in well-aerated clean or mildly polluted water with moderate electrolyte content.





PLATE -14

Diploneis ovalis (Hilse) Cleve

**Dimensions:**

Valve length: 20-22 μm


Valve breadth: 10 μm

Striae density: 10-11/10 μm

Ecology:

Unknown

Craticula sp.

**Dimensions:**

Valve length: 25 μm

Valve breadth: 8 μm


Striae density: /10 μm

Ecology:

Craticula species tend to be associated with fresh to brackish waters and some species are very tolerant to elevated levels of organic pollution. This taxa occurs as single cells in the benthos, but may be washed into the plankton.

PLATE -15

***Craticula accomoda* (Hustedt) DG Mann**
Syn. *Navicula accomoda* Hustedt



Dimensions:

Valve length: 20-23

Valve breadth: 5-6


Striae density: 20-28/10 μm

Ecology:

A common characteristic indicator species for high levels of pollution. Found in strongly organically polluted waters, in particular effluent from sewage treatment works. It has a scattered occurrence in oligo- to eutrophic waters.

PLATE -16

Craticula ambigua (Ehrenberg) DG Mann
Syn. *Navicula ambigua* Ehrenberg




Dimensions:

Valve length: 70-98 μm

Valve breadth: 15-26 μm

Striae density: 18-20/10 μm




Ecology:

A cosmopolitan usually Epipelagic species found in moderately to very electrolyte rich, eutrophic waters, resistant to critical and strong levels of pollution.

PLATE -17

Luticola acidoclinata Lange- Bertalot



Dimensions:

Valve length: 16/18 μm

Valve breadth: 5.5/6 μm

Striae density: 20-24/10 μm

Ecology:

A cosmopolitan species found in oligo- to dystrophic circumneutral to slightly acidic, electrolyte poor waters.

Sellaphora pupula (Kutzing) Mereschowsky sensu lato *Syn. Navicula pupula* Kutzing








PLATE -18

***Sellaphora pupula* (Kutzing) Mereschowsky sensu lato**
Syn. *Navicula pupula* Kutzing




Dimensions:

Valve length: 15-32 μm

Valve breadth: 6-10 μm

Striae density: 7-14/10 μm




Ecology:

A cosmopolitan species found in a broad spectrum of electrolyte rich waters with some populations found under strongly polluted conditions.

PLATE -19

***Sellaphora laevissima* (Kütz.) D. G. Mann**



Dimensions:

Valve length: 25-29 μm

Valve breadth: 9-10 μm

Striae density: 20/10 μm

Ecology:

Unknown

PLATE -20

Eolimna sp.





Dimensions:

Valve length: 4-6 μm
Valve breadth: 3 μm
Striae density: /10 μm

Ecology:

Cosmopolitan, found in a wide range of waters.
More ecology is Unknown.

Fallacia pygmaea (Kutzing) Sickle & Mann
Syn. Navicula pygmaea Kutzing



Dimensions:



Valve length: 17-25 μm
Valve breadth: 9-10 μm
Striae density: 20-25/10 μm

Ecology:

A cosmopolitan Epipelagic species occurring in waters with elevated electrolyte content. Tolerant to critical levels of pollution.

PLATE -21

***Diadesmis confervacea* (Kutzing) DG Mann**
***Syn. Navicula confervacea* Kutzing**




Dimensions:

Valve length: 15-22 μm

Valve breadth: 5.5-7 μm


Striae density: 18-20/10 μm



Ecology:

A cosmopolitan species found in range of waters, including eutrophic, electrolyte rich and extremely polluted waters.

***Nupela* sp.**



Dimensions:

Valve length: μm

Valve breadth: μm

Striae density: /10 μm

Ecology:

Unknown

PLATE -22

***Geissleria decussis* (Hustedt) Lange-Bertalot**

***Syn. Navicula decussis* Østrup**



Dimensions:

Valve length: 23-25 μm

Valve breadth: 7 μm


Striae density: 20-22/10 μm

Ecology:

A cosmopolitan species found in eutrophic, unpolluted or moderately polluted waters with average or slightly elevated electrolyte content.

PLATE -23

Placoneis sp.



Dimensions:

Valve length: 19-28 μm

Valve breadth: 9-10 μm


Striae density: 10-16/10 μm

Ecology:

A cosmopolitan species found in wide range of waters, in unpolluted to slightly polluted waters with moderate to high electrolyte content.

PLATE -24

Anomoeneis sphaerophora (Ehrenberg) Pfitzer



Dimensions:

Valve length: 58-60 μm

Valve breadth: 17.5-18 μm


Striae density: 6-8/10 μm

Ecology:

A common littoral diatom species occurring in water with moderate to high electrolyte content, extending to brackish coastal waters and saline inland biotopes. Tolerant to critical levels of pollution.

PLATE -25

Navicula cf zanonii Hustedt

**Dimensions:**

Valve length: 28-32 μm


Valve breadth: 66.5 μm

Striae density: 14/10 μm

Ecology:

A tropical to sub-tropical species, found commonly in alkaline waters.

Navicula cf antonii Lange-Bertalot

**Dimensions:**

Valve length: 17.5-20 μm

Valve breadth: 5.5-7 μm

Striae density: 9-11/10 μm

Ecology:

Cosmopolitan, found in eutrophic to hypereutrophic waters with moderate to high electrolyte content. Tolerant to strongly polluted conditions. A good indicator for such anthropogenically impacted waters.

PLATE -26

Navicula veneta Kützing

**Dimensions:**

Valve length: 20-22 μm


Valve breadth: 5-6 μm

Striae density: 10-13/10 μm

Ecology:

Cosmopolitan, common in heavily eutrophied, electrolyte-rich to brackish water. Very pollution tolerant, often the dominant species in industrially impacted waters.

Navicula sp.

**Dimensions:**

Valve length: 28-32 μm

Valve breadth: 6-9 μm


Striae density: 12-14/10 μm

Ecology:

Navicula is found in all types of waters ranging from oligotrophic to eutrophic. Cells inhabit the plankton or benthos. In benthic habitats the cells may occur singly, in films on submerged substrates and sediments, or as colonies within a mucilage tube (e.g., *N. recens*). *Navicula* like many other raphe bearing diatoms secretes mucilage from the raphe to enable the cells to glide along a substratum.

PLATE -27

Navicula trivialis Lange-Bert.



Dimensions:

Valve length: 23.5-25 μm

Valve breadth: 6-7 μm


Striae density: 14-16/10 μm

Ecology:

A cosmopolitan Epipelagic species. Occurs in eutrophic waters with moderate electrolyte content. Tolerant of dessication and strongly polluted conditions.

Navicula rostellata Kützing

Navicula viridula var. *rostellata* (Kützing) Cleve



Dimensions:

Valve length: 29-52 μm

Valve breadth: 6.5-11 μm


Striae density: 9-15/10 μm

Ecology:

A cosmopolitan eutrophic species. Tolerant to critical levels of pollution.

PLATE -28

***Navicula erifuga* (OF Muller) Bory**
***Navicula cinctaeformis* Hustedt sensu Cholnoky**



Dimensions:

Valve length: 28-35 μm

Valve breadth: 6.5-7 μm


Striae density: 13-14/10 μm

Ecology:

A cosmopolitan species found in eutrophic, brackish waters or those with very high electrolyte content. Tolerant to critical levels of pollution.

PLATE -29

Navicula viridula (Kütz.) Ehrenb.



Dimensions:

Valve length: 40-100 μm

Valve breadth: 10-15 μm


Striae density: 8-11/10 μm

Ecology:

Cosmopolitan but infrequent. Epilithic, Epipellic, as well as on detritus and macrophytes. Occurring in eutrophic waters and tolerant of critical levels of pollution.

PLATE -30

Navicula gregaria Donkin



Dimensions:

Valve length: 21-26 μm

Valve breadth: 5.5-7 μm


Striae density: 9-10/10 μm

Ecology:

Cosmopolitan, very common in eutrophic to hypereutrophic fresh waters with moderate to highly electrolyte content. Also found in brackish waters. Tolerant of strongly polluted conditions. A good indicator species for these conditions.

Navicula symmetrica Patrick

Syn. Navicula schroeteri var. *symmetrica* (Patrick) Lange-Bertalot



Dimensions:

Valve length: 28-32 μm

Valve breadth: 6-7 μm


Striae density: 14-16/10 μm

Ecology:

Cosmopolitan in eutrophic, electrolyte-rich waters. Tolerant of strongly polluted conditions.

PLATE -31

Navicula germainii J. H. Wallace



Dimensions:

Valve length: 26-40 μm

Valve breadth: 5-8 μm

Striae density: 13-15/10 μm

Ecology:

Cosmopolitan, found in eutrophic waters, tolerant to critical levels of pollution.

PLATE -32

Caloneis bacillum (Grunow) Cleve

**Dimensions:**

Valve length: 23-25 μm

Valve breadth: 8 μm

Striae density: 20-22/10 μm

Ecology:

A cosmopolitan littoral species found in a range of waters with moderate electrolyte content, as well as in damp mosses.

Caloneis molaris (Grunow) Krammer

**Dimensions:**

Valve length: 27-36 μm

Valve breadth: 5.5-6.5 μm


Striae density: 18-20/10 μm

Ecology:

Cosmopolitan, ecology uncertain

PLATE -33

Pinnularia acrosphaeria W Smith



Dimensions:

Valve length: 45-52 μm

Valve breadth: 7-9 μm


Striae density: 10-14/10 μm

Ecology:

Cosmopolitan, but mostly in the tropics. Epipelagic in circumneutral waters with a moderate electrolyte content.

PLATE -34

Pinnularia viridiformis Krammer

**Dimensions:**

Valve length: 70-93 μm

Valve breadth: 13-15 μm

Striae density: 10-11/10 μm

Ecology:

Cosmopolitan, common *Pinnularia* in oligo- to mesotrophic waters with low to moderate electrolyte content.

Pinnularia latarea Krammer

Dimensions:

Valve length: 34-46 μm

Valve breadth: 6-8.5 μm

Striae density: 10-12/10 μm

Ecology:

This taxon is wide spread.






PLATE -35

Pinnularia cf. latevittata Cleve

**Dimensions:**

Valve length: 48-50 μm

Valve breadth: 9 μm

Striae density: 12/10 μm

Ecology:

Unknown

Pinnularia interrupta W. Sm.

Dimensions:

Valve length: 42-46 μm

Valve breadth: 8-8.5 μm

Striae density: 8/10 μm

Ecology:

Cosmopolitan in distribution




PLATE -36

Pinnularia graciloides Hust.

Dimensions:


Valve length: 79-80 μm

Valve breadth: 14 μm

Striae density: 8/10 μm

Ecology:

Unknown



Pinnularia conica Gandhi

Dimensions:

Valve length: 48 μm

Valve breadth: 9-9.5 μm

Striae density: 9/10 μm

Ecology:

unknown





PLATE -37

Pinnularia sp.





Ecology:

A very common benthic genus living on stones and sediment mainly in freshwater, they are often abundant especially in acid waters. Cells of Pinnularia may also be washed into the plankton.

PLATE -38

***Amphora copulata* (Kutzing) Schoeman & Archibald**
Syn. *Amphora libyca* Ehrenberg



Dimensions:

Valve length: 18-40 μm

Valve breadth: 12-24 μm

Striae density: 10-13/10 μm

Ecology:

A cosmopolitan species found in waters with moderate electrolyte content, sometimes occurring in brackish habitat.

PLATE -39

Amphora veneta Kützing

**Dimensions:**

Valve length: 21 μm

Valve breadth: 4 μm

Striae density: 10-12/10 μm

Ecology:

A cosmopolitan species found in waters with elevated electrolyte content, tolerating critical to very heavy pollution.

Amphora montana Krasske *Syn. Amphora submontana* Hustedt

**Dimensions:**

Valve length: 12.5-25 μm

Valve breadth: 8-11 μm

Striae density: 27-36



(40 near apices)/ 10 μm

Ecology:

A cosmopolitan species found in alkaline waters, rarely becoming dominant.

PLATE -40

Cymbella tumida (Brébisson) Van Heurck



Dimensions:

Valve length: 55-60 μm

Valve breadth: 18-18.5 μm

Striae density: 11-12/10 μm

Ecology:

A cosmopolitan species found in oligo- to mesotrophic waters with moderate electrolyte content.

Occurs in the littoral zone of standing and flowing waters.

Cymbella descripta (Hust.) Krammer & Lange-Bert.

Dimensions:

Valve length: 11-13 μm

Valve breadth: 2-3.5 μm

Striae density: 8-12/10 μm

Ecology:

Unknown





PLATE -41

Cymbella turgidula Grunow



Dimensions:

Valve length: 30-40 μm

Valve breadth: 10-13 μm


Striae density: 10/10 μm

Ecology:

A cosmopolitan species found in oligo- to mesotrophic, alkaline waters with moderate electrolyte content.

PLATE -42

***Encyonema mesianum* (Cholnoky) DG Mann**
***Syn. Cymbella mesiana* Cholnoky**



Dimensions:

Valve length: 36-38 μm

Valve breadth: 10-11 μm


Striae density: 8-9/10 μm

Ecology:

A cosmopolitan montane species found in weakly acidic waters.

PLATE -43

***Encyonema minutum* (Hilse) DG Mann**
***Syn. Cymbella minuta* Hilse**



Dimensions:

Valve length: 22-24 μm



Valve breadth: 7 μm

Striae density: 12/10 μm

Ecology:

A cosmopolitan species found in oligotrophic waters with moderate electrolyte content.

***Seminavis strigosa* (Hust.) Danielidis et D.G.Mann**



Dimensions:

Valve length: 21-23 μm

Valve breadth: 5-5.5 μm


Striae density: 8-12/10 μm

Ecology:

Unknown

PLATE 44

***Gomphonema affine* Kützing**



Dimensions:

Valve length: 36-53 μm

Valve breadth: 10-12 μm


Striae density: 10-12/10 μm

Ecology:


A tropical/ subtropical species tolerant of elevated electrolyte concentrations.

PLATE -45

Gomphonema parvulum (Kützing) Kützing sensu stricto



SEM



Dimensions:

Valve length: 20-33 μm

Valve breadth: 6-8 μm

Striae density: 10-16/10 μm

Ecology:

A cosmopolitan species which is very wide spread in a range of waters, from small pools to lakes and rivers and generally considered to be tolerant to extremely polluted conditions.

PLATE -46

Gomphonema pseudoaugur Krammer



Dimensions:

Valve length: 35-37.5 μm

Valve breadth: 10 μm


Striae density: 11-12/10 μm

Ecology:

A cosmopolitan species found in oligo- to mesotrophic waters but not tolerant to critical levels of pollution. Attached to the substratum by dichotomous mucilage stalks.

PLATE -47

Gomphonema cf gracile Ehrenberg sensu stricto



Dimensions:

Valve length: 40-70 μm

Valve breadth: 8.5-11 μm





Striae density: 9-12/10 μm

Ecology:

This taxon is able to tolerate extremely polluted conditions and is found in abundance in mining effluent.

PLATE -48

Gomphonema turris Ehrenberg



Dimensions:

Valve length: 52-56 μm

Valve breadth: 11-12.5 μm

Striae density: 11/10 μm

Ecology:

A cosmopolitan species found in circumneutral waters with moderate electrolyte content.



PLATE -49

Gomphonema sp.







PLATE -50

Gomphonema sp.



Ecology:

Species of Gomphonema are very common epiphytes found attached to substrata by branched, gelatinous stalks originating at their narrow, or base, end. They occur in a wide range of mostly fresh waters including those enriched with sewage.

PLATE -51

Rhopalodia gibba (Ehrenberg) O Müller



Dimensions:

Valve length: 29 μm

Valve breadth: 11 μm

Striae density: 14/10 μm

Ecology:

A cosmopolitan species found in standing and slow flowing waters, especially springs, of moderate to high electrolyte content.

Rhopalodia sp.



Dimensions:

Valve length: 32-34 μm

Valve breadth: 7-7.5 μm


Striae density: 14 /10 μm

Ecology:

The genus may be found in fresh waters as well as brackish waters. Species of this genus are found mostly in the benthos.

PLATE -52

Bacillaria paradoxa Gmelin

**Dimensions:**

Valve length: 66 μm


Valve breadth: 6-6.5 μm

Striae density: 5-9/10 μm

Ecology:

A cosmopolitan species widespread in very electrolyte-rich and brackish waters particularly near the coast. The cells form a unique type of mobile colony, in which individual cells slide to and fro with respect to each other. They are held together by interlocking grooves on the raphe sterna. The colony extends to form a linear array, where only the poles of the cells are touching then retracts to form a tabular array, with all the cells side by side.

Hantzschia amphioxys (Ehrenberg) Grunow

**Dimensions:**

Valve length: 28 μm

Valve breadth: 4 μm


Striae density: 4-11/10 μm

Ecology:

A cosmopolitan species favouring periodically dry habitat, including soils and rock crevices. Widespread in a range of rivers, but probably washed in from soils.

PLATE -53

Hantzschia sp.



Dimensions:

Valve length: 90-110 μm

Valve breadth: 6-7.5 μm

Striae density: 16/10 μm


Fibulae density: 8/10 μm

Ecology:

A cosmopolitan species found in waters with very high electrolyte content.

PLATE -54

***Tryblionella calida* (Grunow) DG Mann**
Syn. *Nitzschia calida* Grunow



Dimensions:

Valve length: μm

Valve breadth: μm

Striae density: 9-12/10 μm

Fibulae density: 7-10/10 μm


Ecology:

A cosmopolitan species commonly occurring in eutrophic waters with elevated electrolyte content.




Favours standing waters.

PLATE -55

Nitzschia obtusa var. *kurzii* Rabenhorst



SEM



Dimensions:

Valve length: 65-110 μm

Valve breadth: 6.5-8 μm

Striae density: >40/10 μm


Fibulae density: 6-8/10 μm

Ecology:

A commonly occurring species found in brackish coastal and inland waters.

PLATE -56

Nitzschia clausii Hantzsch



Dimensions:

Valve length: 31.5-35 μm

Valve breadth: 4-5 μm

Striae density: (32) 38-42/10 μm

Fibulae density: 10-13/10 μm

Ecology:

A cosmopolitan species found in brackish coastal waters as well as in electrolyte-rich inland waters. In large rivers systems this species may be associated with industrial effluents and is tolerant of strongly polluted conditions.

Nitzschia sigma (Kutzing) W Smith

Dimensions:

Valve length: 35-1000 μm

Valve breadth: 4-15(26) μm

Striae density: (15) 19-38/10 μm

Fibulae density: (3)7-12/10 μm


Ecology:

A cosmopolitan species found in eutrophic, electrolyte-rich inland waters and extending into brackish estuarine and coastal biotopes.



PLATE -57

***Nitzschia umbonata* (Ehrenberg) Lange-Bertalot**
Syn. *Nitzschia thermalis* (Ehrenberg) Auerswald



Dimensions:

Valve length: 15-45 μm

Valve breadth: 2.5-7 μm

Striae density: (16) 17- 19(20) /10 μm /10 μm




Fibulae density: 3.56(7)/10 μm

Ecology:

A common species in eutrophic electrolyte rich waters and tolerating extremely polluted conditions.

PLATE -58

Nitzschia palea (Kützing) W Smith



Dimensions:

Valve length: 15-45 μm

Valve breadth: 2.5-7 μm

Striae density: 28-40/10 μm

Fibulae density: 9-17/ 10 μm

Ecology:

A cosmopolitan and very commonly occurring species.

Nitzschia siliqua Archibald

Dimensions:

Valve length: 11-30 μm


Valve breadth: 2-4 μm

Striae density: 16-18/10 μm

Fibulae density: 9-10/10 μm

Ecology:

Unknown



SEM

PLATE -59

Nitzschia intermedia Hantzsch

Dimensions:

Valve length: 60-100 μm

Valve breadth: 4-7 μm

Striae density: 20-33/10 μm

Fibulae density: 7-13/10 μm

Ecology:

Found in the littoral zone of large eutrophic rivers and lakes with moderate to high electrolyte content. This species does not tolerate more than critical levels of pollution.

Nitzschia capitellata Hust.

Dimensions:

Valve length: 40 μm

Valve breadth: 4.5 μm

Striae density: 23-40/10 μm

Fibulae density: 12/10 μm

Ecology:

A wide spread species found in electrolyte rich and brackish waters. Tolerant to extremely polluted conditions.





PLATE -60

Nitzschia frustulum (Kützing) Grunow

**Dimensions:**

Valve length: 11-30 μm


Valve breadth: 2-4 μm

Striae density: 16-18/10 μm

Fibulae density: 9-10/10 μm

Ecology:

A cosmopolitan species found in electrolyte rich and brackish waters. Tolerant to fluctuations in osmotic pressure and of critical levels of pollution.



Nitzschia liebertruthii Rabenhorst

Dimensions:

Valve length: 30-60

Valve breadth: 2.5-3.5

Striae density: 16/10 μm


Fibulae density: 8/10 μm

Ecology:

A cosmopolitan species occurring in very electrolyte-rich to brackish waters.

PLATE -61

***Nitzschia supralitorea* Lange-Bertalot** **Syn. *Nitzschia fonticola* sensu Cholnoky**




Dimensions:

Valve length: 35 μm
Valve breadth: 3.5 μm
Striae density: 16/10 μm
Fibulae density: 14/10 μm

Ecology:

A cosmopolitan species found in eutrophic waters with moderate to moderately high electrolyte content. Also occurring supralittoral sites, tolerant to osmotic fluctuations and strongly polluted conditions.

***Nitzschia amphibia* Grunow**



Dimensions:


Valve length: 25-30 μm
Valve breadth: 4.5 μm
Striae density: 12/10 μm
Fibulae density: 6/10 μm

Ecology:

Found in eutrophic waters over a range from electrolyte-poor to electrolyte-rich waters.

PLATE -62


Nitzschia sp.



Ecology:

Nitzschia is a large, diverse and ecologically versatile Genus. They are mostly benthic but include Planktonic taxa (e.g. *N. acicularis* (Kutzing) W Smith). The genus can yield much ecological information since several taxa are indicative of nutrient enrichment (eutrophication), while other are useful indicators of elevated salinities. Nitzschia may also occur in oligotrophic waters, but they are rarely a major component of these assemblages.

PLATE -63



Nitzschia gracilis Hantzsch

Dimensions:

Valve length: 30-110 μm

Valve breadth: 2.5-4 μm


Striae density: 38-42/10 μm

Fibulae density: 12-18/10 μm

Ecology:

A cosmopolitan species found in eutrophic, electrolyte-rich waters but not tolerating more than moderately polluted conditions.

SEM



Nitzschia pumila Hustedt

Dimensions:

Valve length: 62-66 μm

Valve breadth: 4.5-5 μm

Striae density: 14/10 μm

Fibulae density: 8/10 μm

Ecology:

Little is known of this species except that it occurs in alkaline lakes.




PLATE -64

Surirella angusta Kützing

Dimensions:

Valve length: 33.5 μm


Valve breadth: 10 μm

Striae density: 6/10 μm

Fibulae density: 5.5-8/10 μm

Ecology:

A cosmopolitan species found in eutrophic waters with moderate electrolyte content.





Surirella species

Ecology:

It is a large and common freshwater to marine genus. Cells found in the benthos of hard waters, are rarely planktonic, and seldom abundant. Some species may tolerate very nutrient-rich conditions.

PLATE -65

Surirella tenera



Dimensions:

Valve length: 65-120 μm

Valve breadth: 27-35 μm

Striae density: 3/10 μm

Ecology:

PLATE -66

Surirella species

