


May 1990

Technical Report No 61

To appear in :
Social Biology of wasps (eds) K. G. Ross and R. W. Matthews,
Cornell University Press, in press

RAGHAVENDRA GADAGKAR
Centre for Ecological Sciences
And
Centre for Theoretical Studies
Indian Institute of Science
Bangalore 560 012 INDIA

Belonogaster, Mischocyttarus, Parapolybia,
and Independent - founding Ropalidia



INDIAN INSTITUTE OF SCIENCE
Centre for Ecological Sciences



Table of Contents

Distribution and Systematics.....6

Nest Architecture.....11

Nesting Cycle.....17

Seasonal Nesting Cycle.....26

Aseasonal Determinate Nesting Cycle.....27

Indeterminate Nesting Cycle.....34

Enemies and Colony Defense.....38

Food and Feeding Habits.....47

Mating Behavior.....49

Social Organization.....51

Dominance Hierarchies.....53

Role of the Egg Layer.....55

Behavioral Caste Differentiation.....56

Caste66

Origin of social life : a perspective from studying Independent-
founding polistines.....70

A possible route to eusociality.....77

Concluding Remarks80

Acknowledgements.....82

References.....83

Appendix 5.1.....96

Appendix 5.2.....99

Animals that live in colonies of individuals of more than one generation, cooperate in brood care, and relegate reproduction to one or a small number of colony members are said to represent that pinnacle of social evolution, eusociality (Michener 1969; Wilson 1971). For the naked mole-rat, which lives in underground tunnels in Africa (Jarvis 1981), eusociality has been achieved only by ants and termites and by some bees and wasps. With the exception of one or a few species of sphecids (Mathews 1968a, this volume), all eusocial wasps belong to the family Vespidae. This family has traditionally been divided into three subfamilies, namely Stenogastrinae, Polistinae, and Vespinae (Richards 1962, 1978a,b). Carpenter's (1982) recent classification recognizes three additional subfamilies within the family Vespidae - namely, Eumeninae, Masarinae, Euparigiinae - but eusociality is restricted to the three previously mentioned subfamilies.

Stenogastrines have sometimes been thought to be rather different from the polistines and vespines (Spradbery 1975) and their position within the Vespidae has been questioned (Vecht 1977a), although Carpenter's (1982) cladogram indicates that the Stenogastrinae form a sister group to Polistinae + Vespinae (see Carpenter, this volume). Stenogastrines are regarded as primitively eusocial (see Cowan, this volume [Table 2.1] for definitions of social levels) or, perhaps in some instances, presocial (Turillazzi, this volume). Vespines represent a uniformly highly eusocial group (Matsura, Greene, this volume). That leaves the Polistinae, a rather large group consisting

of 29 genera and about 800 species. Richards (1962) subdivides the Polistinae into three tribes: Ropalidiini (consisting only of the genus *Ropalidia*), Polistini (consisting of *Polistes* and "*Sulcopolistes*") and Polybini (consisting of about 26 genera). However, there are problems with this classification (see Distribution and Systematics). In any case, further subdivision of the Polistinae based on phylogenetic considerations may be less rewarding to students of social evolution than subdivision based on behavior and nest architecture. An elegant example of the latter has been provided by Jeanne (1980a) who distinguishes two subgroups, the *independent-founding* and the *swarm-founding* Polistinae (see also Jeanne, this volume).

Independent-founding Polistinae live in relatively small colonies (rarely more than 100 adult wasps) and construct small, simple, unenveloped combs that are normally suspended by a narrow pedicel. Queens initiate new colonies either singly or in small groups, but without the aid of workers (that is, independently). Swarm founding Polistinae live in more populous colonies (often with 1000 or more adults) and have correspondingly large nests that sometimes have several tiers of combs covered by an envelope. New colonies are always initiated by queens in the company of workers (that is, by swarms). This classification on the basis of behavioral and architectural features is supported by concomitant morphological specializations.

Queens of the independent-founding Polistinae, with their small colonies, do not normally use a pheromonal means of control

over their nestmates; overt physical dominance is the rule. Queens in swarm-founding groups, however, must find it quite impossible to subjugate large numbers of nestmates by physical dominance and have, perhaps for this reason, evolved a pheromonal means of control. Independent-founding species have a special problem with ants. Direct physical resistance to marauding ants is difficult enough with small numbers of adult wasps, but when the nest and its brood have to be left completely unguarded while a single foundress is away foraging it is impossible. Not surprisingly, independent-founding species have evolved a chemical defense against ants. Independent-founding species have a well-developed gland on the sixth (terminal) gastral sternum named van der Vecht's gland, from which an ant repellent substance is secreted. The gland is associated with a tuft of hairs that presumably serves as an applicator brush (Jeanne et al. 1983). Swarm-founding species never leave their nests unattended and usually have a substantial adult population to physically rid the nest of scouting ants. Also, they typically build enveloped nests with only one or a small number of entrance holes. Most swarm-founding species studied appear not to have the specialized ant repellent-producing van der Vecht's gland. Instead, at least some of them have a well-developed Richard's gland on the fifth gastral sternum that is often used to lay an odor trail to guide the members of a swarm to their new nesting site (Jeanne 1981a; Jeanne et al. 1983; see also Downing, this volume).

Swarm founding is characteristic of 24 Polistine genera, as

well as some species of *Ropalidia*. Independent founding is seen in five genera: *Belonogaster*, *Mischocyttarus*, *Parapolybia*, *Polistes*, and some *Ropalidia*. It is easy to argue that the independent-founding polistines provide perfect model systems for studies on social behavior. Small colonies make it possible to mark and study every adult wasp, and the open combs hide nothing from the observer. The nests of many species are quite abundant and often are built in remarkably accessible places. A relatively primitive level of eusociality, characterized by a lack of morphological caste differentiation, an essentially behavioral mechanism of queen control, and considerable flexibility in social roles of the adult wasps make the independent-founding polistines especially attractive subjects. Indeed, they have received increasing attention since the 1970's and have contributed more than any other group of social wasps to the formulation and testing of ideas concerning the forces that mold the evolution of group living and sterile worker castes.

The genus, *Polistes*, unrivaled in its wide distribution and in the attention it has received, is the subject of the previous chapter (Reeve, this volume). The present chapter deals with the remaining four genera of independent-founding Polistinae. Some earlier information regarding these wasps may be obtained from several general reviews (Richards 1971; Spradbery 1973a; Iwata 1976; Akre 1982).

Mischocyttarus is one of the largest genera among all the social vespines. In spite of its taxonomic diversity (206 species), it is essentially restricted to tropical South America, although two species extend into the southern and western portions of North America. The most authoritative account of the systematics of this genus is that of Richards (1978a). A few species have been studied in considerable detail. Notable among these are *M. drewseni* in Brazil (Jeanne 1972), *M. labiatus* in Colombia (Litte 1981), *M. flavitarsis* in Arizona (Litte 1979),

1986; Kojima and Keeping 1985). Several modern studies have been initiated on *B. juncea* and *B. petiolata* (Richards 1969; Keeping and Crewe 1983; Keeping et al. Pardi 1977; Pardi and Marino Piccioli 1970, 1981). More recently, has been studied in detail (Marino Piccioli and Pardi 1970, 1978; study of *B. juncea*, the biology of only one species, *B. grisea*, also Hensen and Blommers 1987). Since Roubaud's (1916) classic taxonomic revision of the genus is that of Richards (1982; see northern Africa, Arabia, and India. The latest comprehensive Africa south of the Sahara, with scattered populations in *Belonogaster*, comprising 79 species, is widely distributed in of each extend into neighboring temperate latitudes. best represented in the tropics, although at least a few species the New World (Veitch 1965, 1967) (Fig 5.1). All four genera are the Old World, whereas *Mischocyttarus* is found exclusively in *Belonogaster*, *Parapolybia*, and *Ropalidia* are restricted to

M. mexicanus in Florida (Little 1977) and in Georgia (Hermann and Chao 1984a). Preliminary information has been published on *M. angulatus* and *M. basimacula* in Panama (Itô 1984b).

Parapolybia is a refreshingly small genus with only five species distributed from Iran in the west, to Japan, the Philippines, and New Guinea in the east (Vecht 1966). Only three of the five species are known to any extent. Yamane (1980, 1984, 1985) has provided a detailed account of *P. varia* in Taiwan along with fragmentary information on *P. nodosa*, while *P. indica* in Japan has been studied by Sugiyura et al. (1983a,b) and Sekijima et al. (1980).

Ropalidia, another large genus, with about 136 known species, occurs in tropical Africa, southern Asia, Australia, and Okinawa. The Indo-Australian species have been revised by Vecht (1941, 1962), the Australian and New Guinea species by Richards (1978b), the Philippine species by Kojima (1984a), the Nepalese species by Yamane and Yamane (1979), and the Indian species by Das and Gupta (1989). Although Vecht (1962) recognized three subgenera, it is now common to recognize six (Richards, 1978b). Of these, the subgenera *Anthrenoida*, *Icariola*, *Polistratus*, and *Ropalidia* can be characterized as independent-founding, although at least a few species in the subgenus *Icariola* appear to be swarm founders (R. Jeanne, unpubl.). *Ropalidia marginata* (Gadgil and Mahabai 1974; Gadagkar 1980; Belavadi and Govindan 1981; Gadagkar et al. 1982a,b; Gadagkar and Joshi 1982a, 1983), *R. cyathiformis* (Gadagkar and Joshi 1982a,b, 1984, 1985), *R. variegata* (Yamane 1986), *R. fasciata* (Itô, 1983, 1985b, 1986b;

Fig. 5.1. Distributions of *Belonogaster*, *Parapolybia*, *Ropalidia* and *Mischocyttarus*, (Redrawn from Vecht 1967, courtesy of Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen Afdeling Natuurkunde.)




Fig. 5.1

The phylogeny of social wasps is now being vigorously investigated (Carpenter 1982, 1987 a,b, 1988a, this volume; Rasnitsyn 1988), particularly with regard to the relationships within the diverse subfamily Polistinae. The main problem with Richards' (1978a) subdivision of Polistinae into tribes is that the the four Old World polistine genera (including the swarm-founding *Polybioides*) appear to have affinities that unite them as a natural group, so that Richards's inclusion of the Old World genera *Parapolybia*, *Polybioides*, and *Belonogaster* with the New World swarm-founding genera in the tribe "Polybiini" is inappropriate. A fascinating behavioral trait linking the Old World polistine genera is their meconium extraction behavior. Usually in social wasps, the larval meconium (fecal pellet) is left at the bottom of the brood cell, but *Belonogaster*, *Polybioides*, *Parapolybia*, and *Ropalidia* have evolved an elaborate behavior of chewing a small hole at the bottom of the cells and removing the meconium (Marino Piccioli 1968; Marino Piccioli and Pardi 1970; Kojima 1983c; Gadagkar, unpubl.). Although differences in the details of this behavior and the means by which the holes are closed are evident, all four Old World genera have windows at the bottoms of cells from which adults have emerged, while such features have not been seen in any of the remaining 25 genera of polistines. This fact alone

to the subgenus *Icarola*.
 founding species studied in any detail, and all of these belong (1980), and *R. cincta* (Darchen 1976a) are the only independent-
 Ito et al. 1985; Kojima 1983a,b,c, 1984b,c; Suzuki and Murai

The suspending of the comb by a long and narrow pedicel coated with an ant repellent is clearly a major defense of these wasps against predation by ants (Jeanne 1970a), although this form of defense appears to be compromised in many nests of *Ropalidia* where the pedicel is short and many secondary pedicels

5.3).

and Evans 1982; Itô 1987a; *M. punctatus*: Richards 1978a) (Fig 1966a,b; Yamane 1986; Gadagkar, unpubl.; *R. revolutionalis*: Hook combs with two columns of cells (e.g., *R. variegata*: Davis which it is constructed. Some species invariably build long irregular or may have sharp corners to suit the substrate on The comb is usually circular or oval, although it may be the cells and brood facing downward (held by a vertical pedicel). by a horizontal pedicel) or oriented in a horizontal plane with rule, and this may be either oriented in a vertical plane (held architecture within the group (Fig.5.2). A single comb is the combs, so that there is relatively little variation in nest but the independent-founding polistines all build simple architecture (Jeanne 1975a; Vecht 1977b; Wenzel, this volume), but the independent-founding polistines all build simple stelecyttarus (suspended by a pedicel), gymnodomus (unenclosed) combs, so that there is relatively little variation in nest architecture within the group (Fig.5.2). A single comb is the rule, and this may be either oriented in a vertical plane (held by a horizontal pedicel) or oriented in a horizontal plane with the cells and brood facing downward (held by a vertical pedicel). The comb is usually circular or oval, although it may be irregular or may have sharp corners to suit the substrate on which it is constructed. Some species invariably build long combs with two columns of cells (e.g., *R. variegata*: Davis 1966a,b; Yamane 1986; Gadagkar, unpubl.; *R. revolutionalis*: Hook and Evans 1982; Itô 1987a; *M. punctatus*: Richards 1978a) (Fig 5.3).


Social wasps show considerable variation in nest architecture (Jeanne 1975a; Vecht 1977b; Wenzel, this volume), but the independent-founding polistines all build simple combs, so that there is relatively little variation in nest architecture within the group (Fig.5.2). A single comb is the rule, and this may be either oriented in a vertical plane (held by a horizontal pedicel) or oriented in a horizontal plane with the cells and brood facing downward (held by a vertical pedicel). The comb is usually circular or oval, although it may be irregular or may have sharp corners to suit the substrate on which it is constructed. Some species invariably build long combs with two columns of cells (e.g., *R. variegata*: Davis 1966a,b; Yamane 1986; Gadagkar, unpubl.; *R. revolutionalis*: Hook and Evans 1982; Itô 1987a; *M. punctatus*: Richards 1978a) (Fig 5.3).

Nest Architecture


volume) confirms.

genera (Jeanne 1980a), as Carpenter's cladistic analysis (this strongly suggests a monophyletic origin for the four Old World


Fig. 5.2. Mature nest of (a) *Belonogaster grisea* (Drawing based on photograph from Marino Piccioli and Pardi 1970, courtesy of *Monitore Zoologico Italiano.*), (b) *Mischocyttarus drewseni* (Drawing based on photograph from Jeanne 1972, courtesy of Museum of Comparative Zoology, Harvard University.), (c) *Parapolybia varia* (Drawing based on photograph from Yamane 1984, courtesy of VEB Gustav Fischer Verlag.) and (d) *Ropalidia marginata* (Drawing based on photograph by Gadagkar). (All drawings by Amy B. Wright). All views are from the side except (d), in which, the comb face is viewed from the front. This comb (d) is perpendicular to the ground and attached to the substratum by a horizontal pedicel which is not visible.



b



p



c



a

Fig 52

Fig. 5.3. A typical long nest of *Ropalidia variegata* with two columns of cells.




Fig 5.3

Belonogaster grisea and *P. varia* show the greatest degree of deviation from the nest architecture typical of the independent-founding polistines. Nests of *B. grisea* (Marino Piccoli and Pardi 1978) are single-combed and are always suspended by means of a single eccentric pedicel attached to the first cell. As the nest grows, it takes on a unique hammocklike appearance, with a highly convex ventral surface and a concave dorsal surface (see Fig. 5.2a). Another atypical characteristic of *B. grisea* nests is that cells are not reused but are often wholly or partly destroyed after the first use. In one population of *P. varia* in Taiwan (Yamane 1984), nests are always

Kojima 1984b).

of nest construction (S. Chandran and Gadagkar, unpubl.; see also opportunities for many females to simultaneously perform the task nest start construction of several combs, thereby providing after the original nest is destroyed, females working on a new construction are for a rather different reason. When re-nesting (see Jeanne 1979a). At least some instances of multiple-comb conceivably minimize damage from certain predators or parasites of this are not entirely clear, although this strategy might 1984b; Herre et al. 1986; Itô 1986b) (Fig. 5.4). The advantages function as a single colony (Gadagkar and Joshi 1982b; Kojima Sometimes several combs are built close to one another and

Parapolybia.

are constructed (see Enemies and Colony Defense). Furthermore, such a means of chemical defense remains to be demonstrated in

New nests are initiated independently (without workers) by one inseminated female (haplometrosis) or a small group of them (pleometrosis) (Table 5.1). In single-foundress colonies, the lone female forages for building material and for food for her larvae, builds the nest, lays eggs, and defends the nest until her first progeny become adults. Foundresses in multiple-foundress colonies, who are often sisters or other close relatives (e.g., Jeanne 1972; Little 1977) engage in aggressive

The Nesting Cycle

Description of nest architecture is by necessity based on studies of a few isolated species, so we still have little appreciation of the relative extent of variation within and among species (see for e.g., Carpenter and Wenzel 1988). *Ropalidia* colonies in India show considerable variation in nest architecture and clearly vary the shapes and sizes of their nests to suit the available nesting site (Davis 1966a; Gadagkar, unpubl.). *Ropalidia marginata* is capable of at least sometimes constructing multicombed nests similar to those of *P. varia* in

Taiwan (see Figs. 5.5, 5.2c).

are not reused.

These subsequent combs are all without pedicels, and their cells additional combs are constructed close to the original one. two or more lateral lobes are added, and later up to ten cells may be reused. As more brood-rearing space is required, multicombed (Fig. 5.2c). The first comb is petiolate, and its

Fig. 5.4. Multiple combs of a *Ropalidia* nest from India.




Fig. 54

Fig. 5.5. An unusual multilobed nest of *Ropalidia marginata* from Bangalore, India. At the time of collection only one of the combs was occupied.




Fig. 5.5

Table 5.1. Single- and multiple-foundress colonies of *Belonogaster*, *Mischocyttarus*, *Parapolybia*, and independent-founding *Ropalidia*

Species	Single-foundress colonies No. (%)	Multiple-foundress colonies No. (%)	Max. no. of foundresses	Time of nest founding	Locality	Reference
<i>Belonogaster petiolata</i>	38 (47)	43 (53)	16	?	Transvaal, S. Africa	Keeping and Crewe 1987
<i>Mischocyttarus diversus</i>	20 (69)	9 (11)	8	Throughout the year	Lower Amazon, Brazil	Daane 1972
<i>M. flavitarsis</i>	131 (97.8)	3 (2.2)	2	March-June	Arizona, USA	Little 1979
<i>M. labiatus</i>	14 (77.8)	4 (22.2)	9	Throughout the year	Colombia, S. America	Little 1979, 1981
<i>M. mexicanus</i>	114 (66.7)	57 (33.3)	20	Throughout the year	Southern Florida, USA	Little 1977, 1979
<i>Parapolybia indica</i>	49 (100)	0	1	May	Southwestern Japan	Sugitara et al. 1983b
<i>P. indica</i>	108 (100)	0	1	May	Southern Japan	Sekijima et al. 1980
<i>P. varia</i>	2 (100) ^a	0 ^a	1	March-May	Central Taiwan	Yamane 1980, 1985
<i>P. varia</i>	1 (7)	13 (93)	22	December-February	Southern Taiwan	Yamane 1985

Although only two nests were studied in detail, the author has anecdotal evidence to suggest that single foundress nests may be the rule.

Species	Single- foundress colonies No. (%)	Multiple- foundress colonies No. (%)	Max. No. of foundresses	Time of nest founding	Locality	Reference
<i>Ropalidia fasciata</i>	331 (75)	109 (25)	13	February- April	Okinaawa, Japan	Suzuki and Murai 1980
<i>R. fasciata</i>	27 (46.6)	31 (53.4)	22	February- April	Okinaawa, Japan	Ito 1985b
<i>R. marginata</i>	8 (29.6)	19 (70.4)	20	Throughout the year	Pune and Bangalore, India	Gadagkar et al. 1982a
<i>R. variegata jacobsoni</i>	37 (63.8)	21 (36.2)	4	Throughout the year?	Sumatra	Yamane 1986

Table 5.1 cont'd

Subordinate foundresses usually begin to die or to disappear from the nest at about the time of the emergence of adult progeny. This first batch of offspring, nearly always females, become workers and take on the duties of foraging, nest building, feeding larvae, and other tasks involved in nest maintenance and defense. This is called the *worker production substage* (*worker phase* of Reeve, this volume) or, sometimes, the *ergonomic substage* (Oster and Wilson 1978). At this point, new cells are

called the *preemergence phase* (*founding phase* of Reeve, this volume). Initiation to the emergence of the first adult offspring is called the *pupal substage*. The entire period from nest spinning of the first cocoon to the emergence of the first adult is referred to as the *larval substage*, and the period from the hatching of the first egg to the spinning of the first cocoon is on their cells and undergo metamorphosis. The period from the foundress(es) start feeding the larvae. Mature larvae spin a cap called *egg substage*. When the eggs begin to hatch, the eggs. Foraging is mostly for building material during this so-increase in the number of cells, all of which are filled with Immediately after nest initiation there is usually a rapid

or more of the subordinate foundresses. other duties of nest and brood maintenance are performed by one occasionally forage for building material. Foraging for food and monopolizes all egg laying and does little else except hierarchy. One foundress at the top of this hierarchy often interactions that lead to the establishment of a dominance

usually added to the nest. Later, males and nonworker females (reproductives or gynes) are produced. In some species, this male and nonworker production stage (reproductive phase of Reeve, this volume) is well separated from the worker production stage. After the reproductives emerge, there is a considerable amount of brood destruction during the so-called *declining substage*, after which the colony usually is abandoned. The time from the emergence of the first worker to nest abandonment is called the *postemergence phase*.

The preemergence phase and the postemergence phase together constitute what might be called a colony cycle. If the nest is abandoned at the end of one such cycle, the colony cycle and the nesting cycle become equivalent and species which show this pattern may be called (after Jeanne, chapt. 6, this volume) *determinate nesting cycle species*. The nesting cycles of the determinate nesting cycle species are sometimes markedly seasonal and fairly synchronous in a population (see Table 5.1). In other cases, nests are initiated asynchronously at all times of the year but the nesting cycle is still determinate because, every nest goes through the six substages of a colony cycle, after which the nest is abandoned. Finally, several colony cycles may be repeated on the same nest because the nest is not necessarily abandoned at the end of each declining substage. Species which show this pattern may be called (after Jeanne, chapt. 6, this volume) *indeterminate nesting cycle species*.

Most species occurring at temperate and subtropical latitudes show a markedly seasonal nesting cycle. Examples include *B. juncea* and *B. petiolata* in southern Africa (Keeping and Crewe 1983), *M. flavitarsis* in Arizona (Litte 1979), *M. mexicanus* in Georgia (Hermann and Chao 1984b), *P. indica* in Japan (Sugiyama et al. 1983b), *P. varia* in Taiwan (Yamane 1980), and *R. fasciata* in the Okinawa (Suzuki and Murai 1980; Ito 1983; Ito et al. 1985). Nests are typically initiated by inseminated, overwintered females early in the spring (see Fig. 5.6). Females that emerge early in the colony cycle become workers, at least partly because there are no males at this time of the year with whom they can mate. However, males of *R. fasciata*, are produced at all stages of the colony cycle, and females of this species emerging from the first brood can mate and found their own nests (Ito and Yamane 1985). Colonies that show a seasonal cycle begin to decline in the fall; wasps that emerge in late summer remain on the nest for a while as nonworking individuals but later leave the nest and mate. The males die and the females hibernate, either on their natal nest (e.g., *R. fasciata*: Ito et al. 1985) or away from the nest in crevices of rocks or under the bark of trees.

The proximate causes of colony decline during the fall are not clear. The accumulation of nonworking males and females at this time may be as important as declines in the food supply. However, seasonal changes in temperature, daylength, and humidity

An excellent example of an aseasonal determinate cycle is provided by *M. drewseni* in Brazil (Fig. 5.8). Nests are initiated throughout the year, go through the typical cycle including the declining substage, and are always abandoned after about six months (Jeanne 1972). *Mischocyttarus mexicanus* in Florida seems to follow an identical pattern (Litte 1977). Because nests are asynchronously initiated, the population is expected to contain males throughout the year. It is thus quite common in these species for inseminated daughters to replace their mothers as queens or to usurp other colonies. Nevertheless, there seems to be a pronounced tendency to produce typical workers in early broods and nonworking females (potential reproductives) in later broods. What is the cause of this tendency, and why does the colony decline so regularly after six months of existence, independently of the season? Jeanne (1972)

Aseasonal Determinate Nesting Cycle

M. mexicanus cubicola: Hermann and Chao 1984b).
 another. (*M. drewseni drewseni*: Dantas de Araújo 1982; within species two examples are known in which populations follow seasonal cycles in one environment and aseasonal cycles in another. Indeed, throughout the year and does not hibernate (Fig. 5.7).
 extremes, *P. indica* in Japan nests for only about three-and-a-half months, whereas *P. varia* in southern Taiwan nests duration of the nesting period and local climate. At the the colony cycle, as evidenced by the relationship between are likely to have a profound effect on the form and duration of

Fig. 5.6. Features of the colony cycle of *Parapolybia indica* in Japan illustrating the seasonal nesting cycle. a, first hatching of larva; b, first appearance of capped cell; c, emergence of adult; d, death of critical number of workers; e, first male emergence; f, death of foundress. The uppermost line indicates total number of cells. (Redrawn from Sugiyama et al. 1983b, courtesy of Bulletin of the Faculty of Agriculture, Mie University.)




Fig. 5.6

Fig. 5.7. Comparison of life cycles of *Parapolybia indica* in warm temperate southwestern Japan, *P. varia* in warm temperate central Taiwan, and *P. varia* in sub-tropical southern Taiwan. H, hibernating phase; P, preemergence phase; M, matrifilial (worker production substage); R, reproductive (male and non-worker production substage). (Redrawn from Sugitara et al. 1983b and Yamane 1980, courtesy of Bulletin of the Faculty of Agriculture, Mie University and Soichi Yamane.)

Fig. 5.7

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Species

Locality and climate

Parapolybia indica



Southwestern Japan
Warm temperate

31

Parapolybia varia



Central Taiwan
Warm temperate

Parapolybia varia



Southern Taiwan
Subtropical

Fig. 5.8. Growth of a typical colony of *Mischocyttarus drewseni*. Colonies may be initiated at any time of the year. A, total cells; B, empty cells; L, first larva ecdosed; P, first larva spun its cocoon to pupate; W, first adult worker emerged; M, first adult male emerged. (Redrawn from Jeanne 1972, courtesy of Museum of Comparative Zoology, Harvard University).




Fig. 5.8

Why are colony decline and abandonment not complete? One

be used for a series of such cycles. to start another colony cycle in the same nest. A given nest may complete in the latter, so that a small number of females remain that dispersal of adults during the declining substage is not and the indeterminate nesting cycle of the *R. marginata* type is between the determinate nesting cycle of the *M. dresseri* type substage involving brood abortion (Fig. 5.9). The main difference stages of the typical colony cycle, including a declining colonies, but successful colonies appear to go through all the pleometrically. Rates of colony failure are very high in young initiated throughout the year either haplotometrically or Gadagkar et al. 1982a,b; Chandrashekara et al. 1990). Nests are independent-founding polistines (Gadgil and Mahabai 1974; clearly documented case of an indeterminate nesting cycle among *Ropalidia marginata* in peninsular India provides the only

Indeterminate nesting cycle

abortion and adult dispersal. nonworkers and males to workers, all go hungry, leading to brood become nonworkers. With a sharp increase in the ratio of colony grows larger more females escape such domination and necessary for a female to assume the role of worker, and as the Jeanne's hypothesis is that physical domination by the queen is conditions but intrinsic factors that regulate the nesting cycle. has argued rather convincingly that it is not environmental

Future studies of such indeterminate nesting cycles are bound to provide many interesting new facts regarding the regulation of colony cycles in these wasps. Because daughters may leave their natal nests to initiate new nests more-or-less continuously, indeterminate nesting cycles may make the composition of a founding group quite heterogeneous in terms of age and social status. We know of at least one clear case in *R. cyathiformis* where a group of individuals who already had well-defined social roles on their natal nest left and initiated a new nest, largely retaining their previous roles (Gadagkar and Joshi

periodically before all is lost to *Vespa tropica*. Large colonies of *R. marginata* might find it adaptive to issue "swarms" of dispersing wasps to found new nests predation by *Vespa tropica*, which is especially relentless on determinate nesting cycle, may have evolved in response to cycle which appears to encompass several repeats of the the colony cycle anew. Alternatively, the indeterminate nesting and that the individual who wins in this competition may start substage is really a period of intense reproductive competition place? It is conceivable that what appears as the declining daughters, then why is there a declining substage in the first queen, in fact, does not become incapable of dominating her control other than physical domination is important. If the throughout the colony cycle or because some mechanism of queen because many of the daughters leave their parental nests the queen is completely unable to dominate her daughters either possibility is that these colonies never reach the stage where

Fig. 5.9. Development of a *Ropalidia marginata* colony transplanted into a cage from which free foraging was allowed. This colony reached a maximum standing crop of 265 cells, 103 eggs, 96 larvae, 76 pupae, 21 adult males, and 47 adult female wasps. After considerable brood and cell destruction, the colony declined in late April to a standing crop of 164 cells, 4 eggs, 0 larvae, 0 pupae, 0 adult males and 17 adult female wasps, only to continue a second colony cycle on the same comb.




Fig. 5.9

There is also no specific defense against parasites, although the alertness of the adults and their efforts to chase away parasites undoubtedly reduce the extent of parasitism. The most common response to parasitized brood is simply to ignore those cells until the parasite emerges, and then to reuse the

1982a; Hermann and Chao 1984a,c).

Adult wasps in the genera considered here seem to have few serious enemies, but a variety of parasites and predators occasionally maraud their nests (Appendixes 5.1 and 5.2). Birds, lizards, and bats sometimes plunder the nests, and tachinid and phorid flies, ichneumonid and Torymid wasps, and pyralid moths may seriously affect productivity by parasitizing the brood. The wasps appear to have no particular defense against vertebrate predators, with the exception of *Mischocyttarus immarginatus*, which nests in association with the much more aggressive wasp *Polybia occidentalis* and thereby presumably derives some protection (Windsor 1972; Gorton 1978). It has been suggested that the choice of nesting site as well as coloration of the nests may help in avoiding predators (Fitzgerald 1950; Kojima 1982a; Hermann and Chao 1984a,c).

Enemies and Colony Defense

distinction between independent founding and swarm founding. species will offer many surprises and may even obliterate the of the indeterminate-cycles in independent founding *Ropalidia* founding. It therefore seems likely that more detailed studies 1984, 1985). This is not very different in principle from swarm

In the Neotropics, army ants must pose some threat to wasp colonies, although their overall impact on wasp populations is difficult to estimate. Several species of *Mischocyttarus* have presumably been under sufficient selection pressure from army ants to evolve nesting associations with "army ant-resistant" ant species. Herre et al. (1986) found that 29 out of 31 active

stay while others leave. brood on the same nest. Often the queen and some of the workers brood and abandon the nest, or sometimes continue to produce more cells with great agitation, and either cannibalize any remaining predator departs. They then return to the nest, inspect the usually sit around the nest completely helpless until the volume). Adults of the prey species are untouched, and they brood of large and conspicuous prey colonies (See Matsura, this systematically search for, locate, and consume almost the entire (Yamane 1980) is the hornet *Vespa tropica*, whose workers southern India (Gadagkar, unpubl.) and *Parapolybia* in Taiwan Perhaps the most serious natural enemy of *Ropalidia* in

(Litt 1981). begins construction of a new nest in roughly the original spot found to be consumed by ants - brood, parasites, and all - and queen simply cuts the pedicel and lets the whole nest fall to the attack by a phorid moth can be so devastating to a nest that the *Mischocyttarus labiatus* is a clear exception to this rule. An to remove or destroy the affected larvae or pupae. cells. Somewhat surprisingly, the wasps seldom make any attempt

The first line of defense appears to be the single narrow pedicel suspending the comb, which restricts routes of access by ants. As a second line of defense, the pedicel is usually coated with an ant repellent produced from the van der Vecht's gland. *Mischocyttarus drewsini* is the prototype for these two forms of defense (Fig. 5.10). The nest pedicel is very narrow and

and devoured by ants within minutes.

pedicel and drop the nest to the ground; it will be discovered only to break down such defenses: remove the adults or break the appreciate the overriding importance of ant predation one has species have evolved fairly effective defenses against them. To the importance of such predators is not obvious is that most wasp workers upon finding a valuable source of food. The reason that out by individual scouts who can quickly recruit additional here is to a variety of species of ants where foraging is carried decided the course of evolution of these wasps. The reference and 5.2, which may tell us little about those enemies that have conundrum is best illustrated by an examination of Appendixes 5.1 have found it necessary to evolve effective defenses? This enemies seen in action today or to those against whom the wasps Should this distinction be given to the most devastating of the What is the most important enemy of these social wasps?

avoid such ant-bearing plants.

derive significant protection from army ants, which consistently *Pheidole* ants. It is clear that by such association the wasps *Mischocyttarus*, were built on plants occupied by *Allomerus* and social wasp nests, including those of ten species of

The third line of defense is the constant guarding of the nest by the adults, who become very agitated when ants are moving nearby. This is, of course, unavailable to a single foundress away foraging, but it appears that larger nests of *Ropalidia* depend primarily on the third line of defense. It is in

defense in *Ropalidia*.

probably diminishes the importance of the first two lines of appear to rub their abdomens on the secondary pedicels. All this substrate (contrast Fig. 5.11 with Fig. 5.2b), and the wasps do not several points to strengthen the attachment of the nest to the pedicel. Besides, secondary pedicels are often constructed at that most ants can probably get to the nest without crossing the unpublish.). On the other hand, the pedicels are often so short repellent has been demonstrated (Kojima 1982b, 1983a; Gadagkar, gland and brush are well developed, and the efficacy of the ant *Ropalidia* females rub their abdomens on the pedicels, the

role of these secretions in chemical defense in this genus. pedicels. Several recent observations (Keeping 1990) support the secretions from their van der Vecht's glands onto their nest those of *B. petiolata* (Keeping 1990) have also been seen to rub Females of *B. grisea* (Marino Piccioli and Pardi 1970) and

1970a).

efficiency of the ant repellent has been demonstrated (Jeanne The gland and the brush are well developed in this genus and the ants but also making it easy to keep coated with ant repellent. long in this species, not only physically restricting access by

Fig. 5.10. Female *Mischocyttarus drewseni* applying ant-repellent secretion by rubbing the tuft of hair on the terminal gastral sternum against the nest pedicel. (From Jeanne 1972, courtesy of Museum of Comparative Zoology, Harvard University.)




Fig. 5.10

Fig. 5.11. A large comb of *Ropalidia marginata* viewed from the side. Notice the large number of pedicels and the narrow distance between the nest and the substrate.






Fig 511

A similar conundrum over the relative importance of historical versus current predation pressures concerns the function of the sting and venom. The most striking feature of wasps to anyone who comes into casual contact with them is their often formidable sting. But detailed studies show no profound role for the sting in the defensive biology of most independent-founding species. Deliberate attention to the role of the sting brings the verdict that "the venom apparatus appears to fall

(1979b)

(Vecht 1967; Jeanne 1975a), especially in the tropics (Jeanne emphasized that predation pressure from ants has been a very important factor in the course of evolution of eusocial wasps defenses are quite effective. It has therefore been rightly quite serious in the absence of defenses but that existing true for all species that predation pressure from ants can be adults, is most important in this species too. It is probably clearly the third line of defense, namely, active defense by nest by adding several additional combs that have no pedicels; wasp doing it once! More important, these wasps enlarge their their abdomens, although Kojima (1983b) reports that he saw one never once saw *P. varia* females rubbing the nest pedicel with In a 20-month study and involving 113 nests, Yamane (1980)

recognition of the importance of the third line of defense that multiple-foundress associations are often considered an antipredator adaptation (see Origin of Social Life : a perspective from studying independent-founding polistines).

This section is intended to highlight our remarkable ignorance of the food and feeding habits of the independent-founding polistines. Barring Jeanne's (1972) study of *M. drewseni*, we have little to go by. Adult wasps forage for two classes of food: arthropods and nectar. It is a safe bet that there is no great specificity about which arthropod species are taken or where the nectar is collected from; easy availability is probably the deciding factor. These wasps are known to hunt lepidopteran larvae (Belavadi and Govindan 1981); spiders (K. Chandrashekara and Gadagkar, unpubl.); and moths, ants, spider eggs, hemipteran nymphs, and tettigoniid grasshoppers (Jeanne 1972). *Mischocyttarus* wasps not only seek out insects caught in spider webs but are actually able to land and walk along the strands of the web, presumably on account of a special adaptation of the tarsal lobes of the mid- and hindlegs (Jeanne 1972). Foragers returning to the nest with masticated solid food

Food and Feeding Habits

short as a defensive mechanism against their chief vertebrate predators" (Hermann and Chao 1984c:339; see also Starr 1985b, 1989b; Kukuk et al. 1989). Here again we must imagine that the concentrated resource that large nests with brood represent was exploited by a large number of vertebrate and invertebrate predators during the evolutionary history of the wasps, and that the sting evolved as a means of deterring these former enemies. Only a few specialized predators have been able to overcome this defense to become important present-day enemies.

Larvae regularly produce droplets of a liquid imbibed by the adults, who often enter larval cells and solicit the secretion by mouthing the larvae. Such larval-adult trophallaxis is a rather striking feature of wasp societies but remains very poorly studied. It is clear from the studies of Hunt *et al.* (1982) that wasp larval secretion has enormous nutritive value for the adults, and it almost certainly is a glandular secretion rather than simply a regurgitation of food given to the larva. A recent attempt to better understand the significance of this form of trophallaxis is that of Hunt (1988, this volume), who seems to have shown that *Mischocyttarus* larvae in preemergence colonies

times of food scarcity.

the adults. Such nectar storage is usually more frequent during larval but for later removal, distribution, and consumption by or young larvae. This is not for direct consumption by the is occasionally stored in empty cells or in cells containing eggs brought to the nest and shared with nestmates and larvae. Nectar honeydew collected from mealybugs or other homopterans, is also Nectar collected from a variety of flowers, and sometimes

regurgitate.

larvae (e.g., Pardi 1977) but have never been observed to masticated. Males may also occasionally offer solid food to lumps as well as by regurgitation after the lumps have been juices are extracted and imbibed. Larvae are fed with the solid food is further masticated by the adult wasps, during which routinely share it with their nestmates, including males. The

Most studies of presocial and primitively eusocial insects aim either to unravel the mechanism of social organization and integration or to understand the origins of cooperative behavior. Quite understandably, males, which appear peripheral to these questions, have largely been neglected. Typically, colonies first produce several worker females and later produce males and nonworking (reproductive) females. Some species, such as *M. drewsi* (Jeanne 1972) and *P. indica* (Sugiyama et al. 1983b), produce males only during a relatively short period toward the end of the colony cycle. Consequently, males of species with a seasonal nesting cycle, such as *P. indica*, are present during only a small part of the year and opportunities for mating are restricted. The aseasonal and asynchronous nesting cycle of *M. drewsi*, however, ensures the presence of males in the population throughout the year.

Mating Behavior

are more likely to surrender saliva than are those in postemergence colonies. His hypothesis to explain this differential behavior is that larvae in preemergence colonies face a greater risk of being cannibalized by the adults, who are themselves likely to be undernourished during this phase of the colony cycle. On the other hand, the better nourished adults in postemergence colonies are less likely to cannibalize larvae. In other words, surrendering of saliva is hypothesized to be a kind of appeasement behavior by larvae, meant to forestall their own deaths by cannibalism.

Mischocyttarus drewseni males in Brazil patrol areas where females are likely to forage, pouncing on conspecific females as well as conspecific males and the similar-looking females of *Polybia sericea* (Jeanne and Castellón Bermúdez 1980). These males make no attempt to mark or defend territories. Males of *M. labiatus* in Colombia similarly patrol areas that are likely to be frequented by females, but drag their abdomens and appear to apply some glandular secretions to their perch sites. The

observed to mate on their natal nests. every day (presumably to attempt to mate). Males have never been entire lives on their natal nests, leaving only for several hours *Cyathiformis* (Gadagkar and Joshi 1984), appear to spend their Males of other species, such as *M. labiatus* (Litte 1981) and *R.* the remaining several days of their lives attempting to mate. nests for a few days after they emerge but then leave to spend *marginata*, *M. drewseni*, and *M. flavitarsis*, remain on their natal males of some species, such as *P. varia* (Yamane 1980), *R.* The behavior of adult males appears is quite diverse. The

in *Polistes* see Reeve, this volume). or usurp the egg-laying position of their mothers (for examples first brood can potentially mate and establish their own colonies consequences for social organization, as females reared from the opportunities to mate are more widespread. This has important fairly long period, so that even if the nesting cycle is seasonal *Cyathiformis* (Gadagkar and Joshi 1984) males are produced over a *Ropalidia fasciata* (Itô and Yamane, 1985) and *R.*

Any society, be it insect, avian, or mammalian should have a set of rules that govern division of labor and access to resources if it is to function efficiently. Without doubt, insect societies deserve to be studied with a view to discover these rules, and, indeed, honey bee and ant societies have been used as model systems with great profit (Oster and Wilson 1978; Seeley 1985; Winston 1987). The study of social organization in

Social Organization

Patrolling males appear to orient visually to anything crudely resembling a conspecific female. Subsequent recognition and release of copulatory behavior is clearly mediated by chemical cues from the female's venom gland and possibly also head and thorax (Litté 1979; Keeping et al. 1986).

(Post and Jeanne 1982b).
 have such well-developed glands and lack the applicator brush that do not conspicuously drag their abdomens generally do not (see Downing, this volume). Males of other species in the genus dense brush of hairs, presumably aiding in pheromone application well-developed exocrine glands on their sterna, which open onto a nests as perch sites (Litté 1979). Males of this species possess other males, sometimes using abandoned or active conspecific behavior. They mark their perch sites and defend them against 1981). Males of *M. flavitarsis* in Arizona exhibit lekking territories and exclude other males as to attract females (Litté function of this application seems not to be as much to mark

All independent-founding polistines, including *Polistes* (Reeve, this volume), have a fundamentally similar social organization. When colonies are founded singly there is, of course, no society until the first progeny emerge, but in multiple-foundress colonies social organization and division of labor are crucial issues in both the preemergence and postemergence phases. In species with seasonal nesting cycles, especially in temperate regions where hibernation is required, foundresses are always of the same generation (even-age cohorts). But with the emergence of the first workers foundresses other than the queen begin to die or disappear, so that the postemergence phase consists primarily of a mother queen and her daughters (see Yamane, 1985). Social organization during the pre- and postemergence phases may thus be expected to be different, as has been emphasized in studies of temperate-zone species (e.g., Yamane 1985). In less-seasonal environments, where asynchronous nesting cycles occur, foundresses may not necessarily belong to the same generation; thus differences in social organization between pre- and postemergence colonies are

group living. certain to shed light on the forces that mold the evolution of eusociality itself. The study of social organization is thus organization is crucial for an understanding of the evolution of foundress colonies are still possible, an appreciation of social not obligately social, at least to the extent that single-even more to offer. Because these primitively eusocial wasps are insect societies such as those considered in this chapter has

The most common such ritualized interaction in *R. marginata* and *R. cyathiformis* consists of one animal, dominant by definition, climbing on top of another and attempting to reach

sees highly ritualized mock fights. Leading to injury and even death. More frequently, however, one fights may involve grappling, biting, and stinging, occasionally and Pardi 1970; Gadagkar 1980; Kojima 1984c; Itô 1985a). Severe frequency of such interactions vary widely (e.g., Marino Piccioli interactions are seen in all species, but the intensity and 1980; Gadagkar and Joshi 1982b; Röseler, this volume). Dominance the establishment of a dominance hierarchy (see also Gadagkar determined largely by means of aggressive interactions leading to Pardi (1948) showed in a classic study, role differentiation is will become the queen and who will take on the worker role. As when a group of wasps nest together it is of interest to know who there is considerable flexibility in the social roles of adults. lack of morphological caste differentiation and consequently Independent-founding polistine wasps are characterized by a

Dominance Hierarchies

when striking differences are evident. mention pre- and postemergence phases of the colony cycle only I shall therefore consider social organization in general and indeterminate nesting cycle in *R. marginata* (Gadagkar, unpubl.). been noticed between pre- and postemergence phases of the less pronounced. For instance, no consistent differences have

Frequencies of dominance interactions vary greatly. Many colonies of *R. marginata* may be observed for days without evidence of aggressive interactions, while other colonies of the same species may show dozens of aggressive interactions per hour of observation. Itô (1983, 1984b, 1985a, 1986a,b) has attempted to discern some pattern in this variation, suggesting that increased levels of dominance interactions in postemergence as opposed to preemergence colonies constitute evidence of maternal manipulation, and that the inherently low levels of dominance interactions characteristic of some *Ropalidia* species may have facilitated the evolution of group living. Considering that so few quantitative data are available, the observed variation in dominance interactions of species such as *R. marginata* and *R.*

but this has not yet been demonstrated.

are species-specific differences in dominance-related behaviors, be radically different from these. It seems probable that there for *P. varia* and by Jeanne (1972) for *M. drewseni* appear not to subordinate. Dominance interactions described by Yamane (1985) wasp nibbling, chasing, sitting on, or holding a wing or leg of a *R. cyathiformis* (Gadagkar and Joshi 1982b), such as a dominant other less frequent forms of dominance interaction, especially in have a very similar form of dominance interaction. There are also 1976a) and *B. grisea* (Marino Piccioli and Pardi 1970) appear to 1980; Gadagkar and Joshi 1982b, 1983). *Ropalidia cincta* (Darchen as compact as possible and its mouthparts inaccessible (Gadagkar subordinate by definition, becomes motionless, keeping its body out and bite its mouthparts. The animal being so treated,

The behavioral repertoire of the queen varies considerably from species to species. At one extreme is the queen of *R. marginata*, who does almost nothing other than lay eggs. In most colonies she rarely indulges in overt dominance interactions with her nestmates, let alone performs any other tasks, although her superior status is obvious - nestmates simply withdraw from her

Without exception, the most dominant female (queen) is the principal if not sole egg layer. Subordinates may lay no eggs (e.g., *R. marginata*: Gadagkar et al. 1990a; premergence colonies of *P. varia*: Yamane 1985; *P. indica*: Sugitara et al. 1983a) or may lay some eggs, most of which are eaten by the queen (e.g., *B. grisea*: Marino Piccioli and Pardi 1970). In at least some cases, however, subordinates lay a substantial proportion of the eggs, which are not necessarily eaten. (e.g., *M. mexicanus* fall nests: Litte 1977; *R. cyathiformis*: Gadagkar and Joshi 1982b).

Role of the Egg Layer

But what makes some females dominant and others subordinate? Although crucial, this question is almost entirely unanswered in this group of wasps. Body size, age, and hormone levels suggest themselves as factors that might influence an animal's probability of becoming dominant over another (Röseler, this volume), but only future work can tell us more.

cyathiformis leads one to suspect that intraspecific variation may often equal or exceed interspecific variation.

The foregoing discussion is based on an emphasis of a queen-worker dichotomy modeled after the highly eusocial insects, but this is arguably an insufficient if not misleading analogy for primitively eusocial species. Variability among workers of the latter is often large, and the range of behaviors or other traits

Behavioral Caste Differentiation

other nest-maintenance activities. In all species studied, subordinates (workers) have the primary responsibilities for tasks other than oviposition, such as foraging, brood care, nest building, and "rarely left the nest and monopolized most ovipositions by physically disturbing the subordinates' attempts to oviposit" (Litte 1981:11). *Parapolybia varia* queens are said to have "frequent biters and solicitors of nestmates" described as "the primary but not the sole egg-layers", "the only food to larvae and nestmates; and even leave the nest to forage building material, and nectar from returning foragers; distribute postemergence colonies, elongate cell walls; solicit food, prepare them to receive fresh eggs; initiate new cells and *drewseni* queens, who chew the silk caps off vacated cells and nest (Gadagkar and Joshi 1984). Even more active are *M. cyathiformis*, on the other hand, is often the most aggressive and active individual in her colony, running about, physically attacking and challenging her nestmates, yet never leaving the presence (Gadagkar and Joshi 1983). The queen of *R.*

of workers may sometimes overlap that of the queen. More significantly, in primitively eusocial insects, caste is at least partly determined in the adult stage, and workers therefore have not completely lost their capacity for direct reproduction. Challenging and replacing the queen, laying eggs in the presence of the queen, or leaving the nest to found their own nest are various ways by which workers may realize direct reproductive success. One might therefore expect workers to adopt a variety of strategies to maximize their chances of direct reproduction. Consequently, variability in worker behavior must hold important clues to the nature of reproductive competition, a fact that has rarely received the attention it deserves.

As an example, Little (1981) distinguished three types of females in preemergence colonies of *M. labiatus* in Colombia; queens, dominant cofoundresses and foraging cofoundresses. Dominant cofoundresses performed fewer tasks than even queens and sometimes laid eggs, while the foraging cofoundresses had poor ovarian development and did most of the foraging. Most daughters became foragers on their natal nests but were nevertheless capable of founding new nests and were therefore considered potential reproductives.

Investigations of *R. marginata* and *R. cyathiformis* in India have been explicitly concerned with the elucidation of interindividual variability and its potential for understanding social organization, reproductive competition, and, ultimately, the evolution of eusocial behavior (Gadagkar and Joshi 1982a,

The position of the queen among these clusters is of particular interest in view of the fact that data on egg laying, the traditional criterion defining the queen-worker dichotomy, were not used in the cluster analysis. In 13 out of 14 R. *marginata* colonies studied the queens were Sitters (Gadagkar and Joshi 1983; Chandrashekhar and Gadagkar 1990). An obvious interpretation of this is that the queens are programmed to spend little time and energy doing anything other than laying eggs. Since there is normally only one queen per colony, the remaining

of bringing food and building material back to the nest. From the nest a great deal of the time and show high frequencies of high frequencies of dominance behaviors. Foragers are absent deal of their time sitting with raised antennae, and also show of their time sitting and self-grooming. Fighters spend a great Sitters are those wasps that spend a relatively large proportion "castes" as "Sitters," "Fighters," and "Foragers" (Fig. 5.12). analysis on data for frequencies of traditionally studied individuals or behavioral castes. Superimposing this cluster differences show that each colony has three distinct kinds of techniques used to examine patterns of interindividual not been the focus of much attention. Multivariate statistical "sitting with raised antennae," behaviors that traditionally have documented include seemingly trivial ones, such as "sitting" or identified wasps using all common activities. The behaviors time activity budgets have been constructed for individually 1983, 1984, 1985). Departing somewhat from typical approaches,


Why do queens of *R. marginata* and *R. cyathiformis* differ in this way? One possibility is that a queen of *R. marginata* is

sitting with raised antennae. Interactions with her nestmates and spending much of her time the colony, routinely taking part in aggressive dominance *marginata* queens, she is one of the most active individuals in does more than just lay eggs. In fact, in contrast to *R. the fighter caste. This means that the queen of R. cyathiformis that in all R. cyathiformis colonies studied, queens belong to two species is very similar, with an important difference being peninsular India, has revealed that social organization in these *marginata* and *R. cyathiformis*, both of which are common in comparative and experimental methods. A comparative study of *R. behavioral caste differentiation can be evaluated using both These hypotheses regarding the evolutionary significance of**

(Chandrashekhara and Gadagkar 1990). significantly better developed ovaries than foragers becoming queens. Besides, both sitters and fighters have gathering food and building material, have the least chance of Foragers, who leave the nest to perform the risky tasks of might be expected, but among themselves. It seems likely that show the highest frequency of fighting, not with foragers as may also be hopeful queens is suggested by the fact that they colony active and guarding it against parasites. That fighters fighters are probably performing the function of keeping the attempting to maximize their chances for future reproduction. Sitters who are not queens may be thought of as hopeful queens

Fig. 5.12. Behavioral profiles of 20 individually identified females (numbered) of two colonies of *Ropalidia marginata* analyzed by principal components analysis. Each point represents one wasp plotted in the coordinate space of the first two principal components. The points fall into 3 clusters (referred to as behavioral castes) by the nearest-centroid (circled dot) criterion. (From Gadagkar and Joshi 1983, courtesy of *Animal Behavior*.)

Fig. 5:12



Evolutionary hypotheses concerning behavioral castes are also amenable to experimental testing (Gadagkar 1987). The approach here has been to examine behavioral caste differentiation in a colony, experimentally remove the queen, and study the consequent changes in social organization. Every *R. cyathiformis* colony seems to include one or two wasps who behave much like the queen (Fig. 5.13). These potential queens, who take over the role of the queen when she is removed, always belong to the fighter caste, as does the queen in this species. Upon removal of the queen, not only does one of the other fighters become the queen, but one of the sitters may change her behavioral profile rather drastically to become a fighter and,


rather than fighting). done on the basis of time activity budgets of behaviors other classified as fighters, because the delineation of a fighter is competition (Notice that even single foundresses can be *cyathiformis* is also a sitter when she faces no reproductive rather than fighter caste. In other words, a queen of *R. cyathiformis*, the queen or egg layer belongs to the sitter caste single egg layer. Second, in single-foundress colonies of *R. cyathiformis* colonies sometimes have multiple egg layers (sitters and foragers also occasionally lay eggs in this species), whereas *R. marginata* colonies usually have only a competition. This idea is supported by several facts. First, *R. cyathiformis* is a fighter because she faces relatively high levels of such competition from her nestmates, while a queen of *R. cyathiformis* a sitter because she faces relatively little reproductive

Yet another line of work has provided additional insights into the social organization of *R. cyathiformis*. During a long-term study, a colony divided so that about half its members left to form a new colony just a few feet away from the parent colony. A behavioral analysis before and after colony fission revealed that inclusive fitnesses of the "rebels" who left the colony, as well as of the "loyalists" who stayed in the old colony, increased as a result of the fission (Gadagkar and Joshi 1985), probably because of the prevailing low efficiency of brood rearing before colony fission associated with high levels of aggression. The second most dominant individual before colony fission became the queen on the new colony. Here, therefore, as an instance where a wasp managed to establish herself as a queen, yet avoided the cost of challenging the original queen as well as the risk of failure that lone foundresses face. Interestingly, loyalists and rebels both behaved as internally coordinated groups well before the actual fission, synchronizing among themselves their times of being on or away from the nest and avoiding members of the other group (Gadagkar and Joshi 1985).

perhaps, the next potential queen. In short, every colony has a potential queen already differentiated even when the original queen is present. An interesting fact is that these potential queens are often young and aggressive individuals who seldom forage for food or do other work. This is in complete contrast to the North American paper wasp *Polistes exclamans*, where older foragers tend to become replacement queens (Strassmann and Meyer 1983).

Fig. 5.13. Behavioral profiles of individually identified females in a *Ropalidia cyathiformis* colony analyzed by principal components analysis. Each point represents one wasp (either before or after the queen was removed) plotted in the coordinate space of the first two principal components. When the queen labeled *Former Queen* was removed the wasp labeled *Potential Queen* became the replacement queen (her modified behavioral profile is labeled *New Queen*). When this new queen was in turn removed, the wasp labeled *Potential Queen 2* became the next replacement queen. (From Gadagkar 1987, courtesy of Verlag J. Peperny.)

Fig. 5:13



Primitively eusocial wasps of the kind we are dealing with provide rich model systems for studying the role of interactions between the adults in determining social organization. But are interactions between adults sufficient to tell us all there is to know about social organization? In other words, are all females at emergence potentially capable of assuming any role in the colony? This has by and large been assumed to be so (e.g., Queller and Strassmann 1989; Reeve, this volume), but since there has never been a direct test there is no firm evidence one way or the other. An important exception is the study by Richards and Richards (1951), who clearly recognized the significance of this question and demonstrated a slight preimaginal caste

Caste

In summary, when viewed with an emphasis on queen-worker dichotomy, social organization appears to achieve efficient division of labour and colony harmony, but when viewed with an emphasis on interindividual variability a rich mosaic of complex behavioral strategies becomes evident, suggesting ways by which individual selection might mold worker behavior (see Origin of Social Life : a perspective from studying independent-founding polistines). It must be stressed, however, that the foregoing discussion of social organization is based on only a few studies of a few species. Future studies of these and the large number of as-yet unstudied species are bound to revise our current ideas (see for e.g., Itô and Higashi 1987; Itô et al. 1988).

differentiation in a number of polistine species. However, most of their material, barring a few species of *Polistes* and *Mischocyttarus*, consisted of highly eusocial swarm-founding Neotropical polistines.

As far as the independent-founding genera are concerned, there appears to have been only one direct test of the null hypothesis that "all eclosing females are potentially capable of laying eggs" (Gadagkar et al. 1988:176, 1990b). This hypothesis was tested by collecting a number of nests of *R. marginata* and isolating emerging females in individual cages. Of 299 virgin female wasps from 39 nests so tested, only 150 laid eggs. The remaining 149 died without doing so in spite of living, on average, longer than the time taken by the egg layers to lay their first eggs. This result clearly suggests that there is some preimaginal biasing of caste. But what factors determine which females will become egg layers and which will become non egg layers? Among a large number of variables studied only two were significantly correlated with the probability of a wasp becoming an egg layer. One of these variables, the number of empty cells in the parent nest, is likely to be a strong indicator of the queen's declining influence on a colony. The second correlate, the rate of food consumption by wasps in the experiment, varied considerably in spite of all animals being housed in individual cages and being provided unlimited access to food.

On the basis of these results Gadagkar et al. (1988) have

If the queen or other adults bias the future caste of their colony's brood, one might expect the brood in a preemergence colony to be channeled toward worker development and the brood in a mature postemergence colony to be channeled toward queenlike development. If such channeling is based on differential larval

Indirect evidence suggests the existence of a similar caste-biasing system in *B. grisea*. Pardi and Marino Piccioli (1970, 1981) found two distinct classes of females: (1) larger "queenlike" females who were fertilized more often, were more oophagous, and were mainly nest foundresses and (2) "workerlike" females who had the opposite traits and foraged more often. These authors postulated that preimaginal trophic factors determine the two "morphophysiological" conditions.

proposed a model for preimaginal biasing of caste in primitively eusocial insects. With a decline in the queen's influence resulting from poor health, old age, or a temporary programmed shift in her physiology and/or behavior, a set of processes would be initiated that has two consequences: (1) the accumulation of empty cells and (2) the production of a class of female offspring programmed to feed more and so to have a high probability of becoming reproductively competent. In contrast, when the queen's influence is high there are two rather different consequences: (1) the absence of empty cells and (2) the production of a class of female offspring programmed to feed relatively less. Preimaginal caste bias is undoubtedly partial, leaving considerable potential in the adult stage for environmental and other social factors to influence caste.

The above discussion suggests that some form of preimaginal caste biasing already occurs at the level of social evolution represented by the independent-founding polistines. While postemergence events undoubtedly influence caste, preemergence factors should be studied with greater vigor and better techniques. Whether all adult females are equally capable of assuming any role in the colony in the absence of postemergence social interactions is still an open question for most species.

Hunt (1988) has pointed out a factor that can potentially exaggerate such differences in larval nourishment. This is the tendency of *Mischocyttarus* larvae in preemergence colonies to give up saliva to the soliciting adults more readily than larvae in postemergence colonies (see Food and Feeding Habits). The result is an exaggeration of the difference in nourishment between larvae from pre- and postemergence phases of the colony cycle, thus providing a more powerful mechanism of channeling emerging females into their respective worker and reproductive roles.

Although no quantitative data on larval nourishment in pre- and postemergence colonies are available, there is some evidence that *R. marginata* nests with relatively better nourished larvae produce potential egg layers and those with relatively poorly nourished larvae produce potential non egg layers (Gadagkar et al. 1990c).

nourishment, one would expect larvae in preemergence colonies to be poorly nourished compared with larvae in mature postemergence colonies. Although no quantitative data on larval nourishment in pre- and postemergence colonies are available, there is some evidence that *R. marginata* nests with relatively better nourished larvae produce potential egg layers and those with relatively poorly nourished larvae produce potential non egg layers (Gadagkar et al. 1990c).

In most highly eusocial insects such as honey bees, termites, and most ants, the simple answer to those questions could be that individuals who accept sterile worker roles simply have no other choice; over evolutionary time they have lost the ability to reproduce on their own. Workers in many species are capable of laying eggs, but with rare exception (e.g., Anderson 1963) they lay only haploid, male-producing eggs, and even then such worker reproduction is usually suppressed in the presence of the queen (Hamilton 1964b, 1972; Wilson 1971; Trivers and Hare 1976; Fletcher and Ross 1985; Ross 1985; Page and Erickson 1988;

answer such questions.

Why do social wasps live in groups? Why do some individuals accept the role of sterile worker? During colony founding, why do some individuals nest with others even if it means few or no opportunities to lay their own eggs? During the early post emergence phase of the colony cycle, why do many females stay to help their mothers produce more offspring? Primitively eusocial polistine wasps are attractive model systems in insect sociobiology because they seem to provide the opportunity to

founding polistines

Origin of Social Life : a perspective from studying independent-

If the answer is negative, as it seems to be for some species, then we should be viewing these primitively eusocial insect societies from a rather different perspective and suitably modify our theories to explain altruism.

solitary life if
of inclusive fitness and say that group life is favored over
volume). We may broadly generalize Hamilton's (1964 a,b) concept
inclusive fitness and kin selection see Ross and Carpenter, this
relatives if the group consists of close kin (for discussion of
opportunities to gain inclusive fitness by caring for their
if the former means sterility? Workers in social groups may have
How can social life be more advantageous than solitary life

than simply its maintenance.
exist, allows us to focus on the origin of social life rather
this chapter, where real choices concerning reproductive roles
study of primitively eusocial species such as those considered in
sometimes be more advantageous than solitary life. Besides the
social life, even if it means partial or full sterility, must
not because they are incapable of doing anything else but because
strongly suggest that individuals are accepting subordinate roles
new colonies (Gadagkar and Joshi 1984, 1985). All these facts
Gadagkar, unpubl.); and females leave their natal nests to found
away, and replace mother queens (Jeanne 1972; Yamane 1986;
1979, 1981; Gadagkar 1987) daughters sometimes challenge, drive
the most dominant female is lost or removed (Jeanne 1972; Little
(Table 5.1); subordinate females adopt the role of the queen if
colonies coexist in *Mischocyttarus*, *Ropalidia*, and *Parapolybia*
option of direct reproduction. Single- and multiple-foundress
in which the animals almost certainly have a choice regarding the
In primitively eusocial species there are several situations
Ratneiks 1988; Ratneiks and Visscher 1989; Visscher 1989).

Many features of the biology of the genera considered in this chapter suggest that conditions necessary for high r between workers and the brood they rear may not be met (see also Ross and Carpenter, this volume). In *B. petiolata* only 15% of foundress associations consist exclusively of former nestmates (Keeping and Crewe 1987), so that co-foundresses are not often

only 0.25 while they are related to their sons by 0.5). Instead of brothers (workers are related to their brothers by favor of their sisters or must be able to rear their own sons Furthermore, workers either must successfully skew investment in any half-sisters or more distantly related individuals. egg layer mated to a single male so that workers need not rear Realization of high r requires that colonies consist of a single between a female and her full (= super) sisters is 0.75. such as is found in Hymenoptera, where genetic relatedness be achieved with a male haploid (haplodiploid) genetic system, relatives reared to be greater than that to offspring. This may greater than 0.5, that is, for the genetic relatedness to inequality may be obtained is for r in the group mode to be offspring reared in the solitary mode. One way in which this genetic relatedness to these individuals and m is the number of relatives) reared in the group mode, r_1 is the coefficients of where n is the number of individuals (offspring or other

$$\sum_{i=1}^n r_i > 1/2 m \quad \dots \dots \dots (5.1)$$

Polygyny or multiple mating by the queen should pose no great difficulty for attaining high levels of inclusive fitness if workers discriminate between full sisters and less-related individuals, giving preferential aid to the former (Gadagkar 1985b). However, studies of nestmate discrimination in *R. marginata* suggest that the labels and templates used in discrimination are not produced individually, but rather are acquired from a common external source, namely the natal nest or nestmates, making it unlikely that different levels of genetic relatedness can be effectively recognized among members of the same colony (Venkataraman et al. 1988). Thus, it seems likely that workers in these primitively eusocial wasps often rear complex mixtures of full sisters, half-sisters, nieces,

et al. 1986)

and *R. marginata* ($r = 0.53$, calculated from data in Muralidharan and *Mischocyttarus immarginatus* ($r = 0.765$) (Strassmann et al. 1989) measured in *Mischocyttarus basimacula* ($r = 0.435$), Muralidharan et al. 1986). Relatedness between sisters has been the egg layer is known in at least one species (*R. marginata*: conspecifics in *M. flavitarsis* (Litte 1979). Multiple mating by (Ito 1986b) and nests frequently are usurped by foreign 1984), *R. variegata jacobsoni* (Yamane 1986) and *R. fasciata* nests (Litte 1977), *R. cyathiformis* (Gadagkar and Joshi 1982b, more than one egg layer, has been reported in *M. mexicanus* fall Gadagkar et al. 1990a). Polygyny, the simultaneous presence of *marginata*, queen supersedeure is common (Jeanne 1972; Yamane 1986; closely related. In *M. dreswenti*, *R. variegata jacobsoni*, *R.*

daughters, brothers, nephews, sons, and cousins without the ability to discriminate on the basis of genetic relatedness. - Because r in such societies rarely exceeds 0.5, it has been increasingly suspected that haplodiploidy is not as important a factor in the origin of insect sociality as was earlier thought (Evans 1977a; West-Eberhard 1978a; Andersson 1984; Stubblefield and Charnov 1986; Venkataraman et al. 1988; Gadagkar 1990a,b).

Even if r is rarely much greater than 0.5 (or even less than 0.5), the inequality in Eq. 5.1 can nonetheless be achieved if n is sufficiently greater than m . The ecology of independent-founding polistine wasps suggests that there must be substantial benefits to group living, that is, n must often be greater than m (see also Reeve, this volume). For instance, the probability of survival of single-foundress colonies is certainly small, as the nest and its brood are extremely vulnerable to predators and parasites (e.g., Suzuki and Murai 1980). Yet the effectiveness of protection from such enemies by even one or two supernumerary adults is obvious to anyone who has watched these wasp colonies. Furthermore, multiple foundresses are more likely to be able to rebuild a damaged nest than are solitary foundresses. This latter factor favours multiple-foundress nests even when the destructive agent is itself indifferent to group size, as in the case of predation by birds or destruction by typhoons (Litte 1977, 1979, 1981; Itô 1983, 1984b, 1985 a,b,c, 1986a, 1987b; Kojima 1989). In addition, a combination of high adult mortality and long development times in *R. marginata* have been shown to enhance inclusive fitness of workers relative to that of solitary

The presence of permanently sterile worker castes is the most prominent and seemingly paradoxical feature of the highly eusocial insects. Attempts to explain the evolution of eusociality have sometimes obscured the rather obvious fact that

contribute to achieving the inequality in Eq. 5.1. The evidence for preimaginal caste bias in *R. marginata* (see 1979, 1983), and empirical (Michener and Brothers 1974) studies. theoretical (Stubblefield and Charnov 1986), modeling (Craig manipulation and subfertility hypotheses have found support in rather easily. The general ideas embodied in the parental be so small that the inequality in Eq. 5.1 may be satisfied (subfertility hypothesis: West-Eberhard 1975). That is, m will it "easier" to give up reproduction and accept a worker role variation in the quantity of food obtained as larvae) will find subfertility females produced by whatever cause (even by accidental related idea, which circumvents this problem is the idea that successful in making them overcome parental manipulation. A with this is whether counterselection on the offspring would be manipulation hypothesis: Alexander 1974). A significant problem more grandchildren than her wildtype counterpart (parental helping to rear her remaining (fertile) offspring leaves behind manipulates a fraction of her offspring into being sterile and attained is if ecological conditions exist such that a parent who Another way by which the inequality in Eq. 5.1 may be

many social insect species do not possess permanently sterile workers but could nevertheless be forerunners of the highly eusocial state. The evolutionary forces that promoted the origin of such primitive levels of sociality may thus have been quite different from those discussed above. The statements that "social behavior in insects is in part mutualistic" and that "social colonies without altruism are therefore considered a probability" (Lin and Michener 1972:131) may well prove to have been prophetic. The theory of reciprocal altruism proposed by Trivers (1971), an individual-level selection theory which suggests that aid may be given today with the hope of getting a return tomorrow, has for some mysterious reason seldom been applied to social insect colonies. A related individual selection model has been discussed by West-Eberhard (1978a) and Gadagkar (1985a). Consider two wasps that come together and nest jointly. If their joint productivity (say, 21 offspring) is even slightly greater than the sum of their individual productivities (say, $10 + 10 = 20$ offspring) in the solitary mode, and if the roles of queen and sterile worker are assigned randomly, then wasps who take the risk of joint nesting will, on average, produce 10.5 offspring and thereby do better than those who shy away from the risks of joint nesting. In this "gambling" model, benefits of group living can be infinitesimally small, and abilities of parents to manipulate their offspring or of colony members to discriminate on the basis of differential genetic relatedness are not necessary.


operative at this final stage. words, the benefits of haplodiploidy for social evolution become begin to have selective value (*the recognition stage*). In other recognize and give preferential aid to closer relatives will roles. It is precisely at this stage that the ability to begin to lose reproductive options and to get trapped into worker living become increasingly unavailable to some individuals, who manipulation becomes increasingly effective, benefits of group ability to manipulate offspring can be selected for. As food supply leading to subfertility can be exploited, and the manipulation (*the manipulation stage*). Accidental variations in Once group living is established, the stage is set for

Dawkins 1979; Brockmann et al. 1979). (Tinbergen 1932, 1935; Tinbergen and Kruyt 1938; Brockmann and seem to possess an appropriately diverse array of behaviors repertoire to permit the necessary interactions; solitary wasps societies by mutualism is a sufficiently complex behavioral 1988). The only prerequisite for the evolution of incipient among kin will evolve more easily (West-Eberhard 1978a; Schwarz for recognition of genetic relatedness, although group living possible without any preadaptation for parental manipulation or benefits of gambling (*the gambling stage*: Fig. 5.14). This is living comes from mutualism, reciprocal altruism, and the credibility to the idea that the initial incentive for group Many features of the genera discussed in this chapter lend

A possible route to eusociality

Fig. 5.14. The route to eusociality, a hypothesis concerning the evolution of the highly eusocial state from the solitary state through the gambling, manipulation, and recognition stages. The examples given for each stage are tentative as our knowledge of the causes and consequences of group living in most social insect groups is rather sketchy.

Fig. 5.14



Any attempt to collate the rapidly accumulating information on polistine biology may appear aimless unless there is a well-defined structure around which this information is organized. Therefore, recognition of the existence of two behaviourally distinct groups by Jeanne (1980a), namely the "independent-founding polistines and the "swarm-founding polistines," was perhaps the most significant recent conceptual advance in our understanding of polistine wasps. Much work since then has aimed to further consolidate such a classification. We may now have reached the stage, however, of beginning to question this classification and to start looking at exceptions to the

Concluding Remarks

This route to eusociality (Gadagkar 1990d) explains one otherwise curious fact. The ability to discriminate among the members of a colony on the basis of relatedness seems to be absent in all primitively eusocial species studied but present in the highly eusocial ants and honey bees (reviewed in Venkataraman et al. 1988). If haplodiploidy were important for the origins of insect eusociality one would expect workers in primitively eusocial species to exploit the genetic asymmetries thus created by discriminating between close and distant relatives. But if eusociality originated because of mutualistic benefits, as assumed here, and its subsequent maintenance in highly eusocial forms is due largely to haplodiploidy, the observed distribution of kinship discrimination abilities is no longer a paradox.

characteristics of each group. To what extent do species of *Ropalidia*, and *Parapolybia* depend on chemical defense against ants? To what extent is nest founding by several females of different ages, arriving simultaneously and sometimes having well-defined social roles in their previous colonies, really different from swarm founding? What means of communication, if any, do the independent-founding species use in choosing and reaching a new nesting site? This is not to imply that we have reached the end of the utility of Jeanne's (1980a) classification. Indeed, it is a tribute to its continuing utility that we expect significant new advances in polistine biology to be triggered by standing this classification on its head and looking for exceptions!

It is a pleasure to thank Kenneth G. Ross, Robert W. Mathews, Robert L. Jeanne, Madhav Gadgil, and Seetha Bhagavan for many helpful comments on an earlier draft of this chapter and G. Jayashree, S. Veda and U. N. Shyamala for much assistance in its preparation. I thank Robert L. Jeanne for suggesting the terms "determinate nesting cycle species", "indeterminate nesting cycle species" and "preimaginal caste bias". My own work was supported by grants from the Department of Science and Technology, Government of India, and the Indian National Science Academy.

Acknowledgements

1. Akre, R. D. 1982. Social wasps. pp. 1-105. In H. R. Hermann (ed.), Social Insects, Vol. 4. Academic Press, New York.
2. Alexander, R. D. 1974. The evolution of social behavior. Annu. Rev. Ecol. Syst. 5:325-383.
3. Anderson, R. H. 1963. The laying worker in the Cape honey bee, *Apis mellifera capensis*. J. Apic. Res. 2: 85-92.
4. Andersson, M. 1984. The evolution of eusociality. Annu. Rev. Ecol. Syst. 15: 165-189.
5. Belavadi, V. V., and R. Govindan. 1981. Nesting habits and behaviour of *Ropalidia (Icaridia) marginata* (Hymenoptera: Vespidae) in south India. Colemania 1: 95-101.
6. Brockmann, H. J., and R. Dawkins. 1979. Joint nesting in a digger wasp as an evolutionarily stable preadaption to social life. Behaviour 71: 203-245.
7. Brockmann, H. J., A. Grafen, and R. Dawkins. 1979. Evolutionarily stable nesting strategy in a digger wasp. J. Theor. Biol. 77: 473-496.
8. Brooks, R. W., and D. B. Wahl. 1987. Biology and mature larva of *Hemipimpla puchripennis* (Saussure), a parasite of *Ropalidia* (Hymenoptera: Ichneumonidae, Vespidae). J. New York Entomol. Soc. 95: 547-552.
9. Carpenter, J. M. 1982. The phylogenetic relationships and natural classification of the Vespidae (Hymenoptera). Syst. Entomol. 7: 11-38.
10. Carpenter, J. M. 1987a. Phylogenetic relationships and classification of the Vespinae (Hymenoptera: Vespidae). Syst. Entomol. 12: 413-431.
11. Carpenter, J. M. 1987b. On "The evolutionary genetics of social wasps" and the phylogeny of the Vespinae (Hymenoptera: Vespidae). Insectes Soc. 34: 58-64.
12. Carpenter, J. M. 1988a. The phylogenetic system of the Stenogastrinae (Hymenoptera: Vespidae). J. New York Entomol. Soc. 96: 140-175.
13. Carpenter, J. M., and J. W. Wenzel. 1988. A new species and nest type of *Mischocyttarus* from Costa Rica (Hymenoptera: Vespidae; Polistinae), with descriptions of nests of three related species. Psyche 95: 89-99.

14. Chandrashekara, K., and R. Gadagkar, 1990. Evolution of eusociality: lessons from social organization in *Ropalidia marginata* (Lep.) (Hymenoptera: Vespidae). In Proc. 11th International Congress of IUSSI, Bangalore, India (in press).
15. Chandrashekara, K., S. Bhagavan, S. Chandran, P. Nair, and R. Gadagkar, 1990. Perennial indeterminate colony cycle in a primitively eusocial wasp. In Proc. 11th International Congress of IUSSI, Bangalore, India (in press).
16. Craig, R. 1979. Parental manipulation, kin selection, and the evolution of altruism. Evolution 33: 319-334.
17. Craig, R. 1983. Subfertility and the evolution of eusociality by kin selection. J. Theor. Biol. 100: 379-397.
18. Dantas de Araújo, C. Z. 1982. Bionomia comparada de *Mischocyttarus [sic] dreswenti* das regiões subtropical (Curitiba, PR) e tropical (Belém, PA) do Brasil (Hymenoptera, Vespidae). Dusenía 13: 165-172.
19. Darchen, R. 1976a. *Ropalidia cincta*, guêpe sociale de la savane de Lamto (Cote-d'Ivoire) (Hym. Vespidae). Ann. Soc. Entomol. France (N.S.) 12: 579-601.
20. Das, B. P., and V. K. Gupta. 1989. The social wasps of India and the adjacent countries (Hymenoptera: Vespidae). Oriental Ins. Monogr. no. 11: 1-292.
21. Davis, T. A. 1966a. Nest structure of a social wasp (*Ropalidia variegata*) varying with siting of leaves. Nature 210: 966-967.
22. Davis, T. A. 1966b. Observations on *Ropalidia variegata* (Smith) (Hymenoptera: Vespidae). Entomol. News 77: 271-277.
23. Evans, H. E. 1977a. Extrinsic versus intrinsic factors in the evolution of insect sociality. Bioscience 27: 613-617.
24. Fitzgerald, D. V. 1950. Notes on the genus *Ropalidia* (Hymenoptera: Vespidae) from Madagascar. Proc. R. Entomol. Soc. London (A) 25: 81-86.
25. Fletcher, D. J. C., and K. G. Ross. 1985. Regulation of reproduction in eusocial Hymenoptera. Annu. Rev. Entomol. 30: 319-343.
26. Gadagkar, R. 1980. Dominance hierarchy and division of labour in the social wasp *Ropalidia marginata* (Lep.) (Hymenoptera: Vespidae). Curr. Sci. 49: 772-775.
27. Gadagkar, R. 1985a. Evolution of insect sociality - a review of some attempts to test modern theories. Proc. Indian Acad. Sci. (Anim. Sci.) 94: 309-324.

28. Gadagkar, R. 1985b. Kin recognition in social insects and other animals - a review of recent findings and a consideration of their relevance for the theory of kin selection. Proc. Indian Acad. Sci. (Anim. Sci.) 94: 587-621.
29. Gadagkar, R. 1987. Social structure and the determinants of queen status in the primitively eusocial wasp *Ropalidia cyathiformis*. pp. 377-378 In J. Eder and H. Rembold (eds.), Chemistry and Biology of Social Insects. J. Peperny, Munich.
30. Gadagkar, R. 1990a. The haplodiploidy threshold and social evolution. Curr. Sci. (in press).
31. Gadagkar, R. 1990b. Social biology of *Ropalidia*: investigations into the origins of eusociality. In Proc. 11th International Congress of IUSSI, Bangalore, India (in press).
32. Gadagkar, R. 1990c. Evolution of eusociality: the advantage of assured fitness returns. Phil. Trans. R. Soc. Lond. B. (in press).
33. Gadagkar, R. 1990d. Origin and evolution of eusociality: a perspective from studying primitively eusocial wasps. J. Genet. (in press).
34. Gadagkar, R., and N. V. Joshi. 1982a. A comparative study of social structure in colonies of *Ropalidia*. pp. 187-191 In M. D. Breed, C. D. Michener, and H. E. Evans (eds.), The Biology of Social Insects. Westview, Boulder, CO.
35. Gadagkar, R., and N. V. Joshi. 1982b. Behaviour of the Indian social wasp *Ropalidia cyathiformis* on a nest of separate combs (Hymenoptera: Vespidae). J. Zool., London 198: 27-37.
36. Gadagkar, R., and N. V. Joshi. 1983. Quantitative ethology of social wasps: time-activity budgets and caste differentiation in *Ropalidia marginata* (Lep.) (Hymenoptera: Vespidae). Anim. Behav. 31: 26-31.
37. Gadagkar, R., and N. V. Joshi. 1984. Social organisation in the Indian wasp *Ropalidia cyathiformis* (Fab.) (Hymenoptera: Vespidae). Z. Tierpsychol. 64: 15-32.
38. Gadagkar, R., and N. V. Joshi. 1985. Colony fission in a social wasp. Curr. Sci. 54: 57-62.
39. Gadagkar, R., M. Gadgil, N. V. Joshi, and A. S. Mahabal. 1982a. Observations on the natural history and population ecology of the social wasp *Ropalidia marginata* (Lep.) from peninsular India (Hymenoptera: Vespidae). Proc. Indian Acad. Sci. (Anim. Sci.) 91: 539-552.

40. Gadagkar, R., M. Gadgil, and A. S. Mahabal. 1982b. Observations on population ecology and sociobiology of the paper wasp *Ropalidia marginata* (Lep.) (Family Vespidae). Proc. Symp. Ecol. Anim. Popul. Zool. Surv. India 4: 49-61.
41. Gadagkar, R., C. Vinutha, A. Shanubhogue, and A. P. Gore. 1988. Pre-imaginal biasing of caste in a primitively eusocial insect. Proc. R. Soc. London (B) 233: 175-189.
42. Gadagkar, R., K. Chandrashekara, S. Chandran, and S. Bhagavan. 1990a. Serial polygyny in *Ropalidia marginata*: implications for the evolution of eusociality. In Proc. 11th International Congress of IUSSI, Bangalore, India (in press).
43. Gadagkar, R., S. Bhagavan, R. Malpe, and C. Vinutha. 1990b. On reconfirming the evidence for pre-imaginal caste bias in a primitively eusocial wasp. Proc. Indian Acad. Sci. (Anim. Sci.) 99: 141-150.
44. Gadagkar, R., S. Bhagavan, K. Chandrashekara, and C. Vinutha. 1990c. The role of larval nutrition in pre-imaginal biasing of caste in the primitively eusocial wasp *Ropalidia marginata* (Lep.). Submitted.
45. Gadgil, M., and A. S. Mahabal. 1974. Caste differentiation in the paper wasp *Ropalidia marginata* (Lep.). Curr. Sci. 43: 482.
46. Gorton, R. E. 1978. Observations on the nesting behavior of *Mischocyttarus imarginatus* (Rich.) (Vespidae: Hymenoptera) in a dry forest in Costa Rica. Insectes Soc. 25: 197-204.
47. Hamilton, W. D. 1964a. The genetical evolution of social behaviour. I. J. Theor. Biol. 7: 1-16.
48. Hamilton, W. D. 1964b. The genetical evolution of social behaviour. II. J. Theor. Biol. 7: 17-52.
49. Hamilton, W. D. 1972. Altruism and related phenomena, mainly in social insects. Annu. Rev. Ecol. Syst. 3: 193-232.
50. Hensen, R. V., and L. H. M. Blommers. 1987. Review of the Malagasy species of *Belonogaster* Saussure (Hymenoptera, Vespidae). Tijdschr. Entomol. 130: 11-31.
51. Hermann, H. R., and J. T. Chao. 1984a. Nesting biology and defensive behavior of *Mischocyttarus* (*Monocyttarus*) *mexicanus cubicola* (Vespidae: Polistinae). Psyche 91: 51-65.
52. Hermann, H. R., and J. T. Chao. 1984b. Distribution of *Mischocyttarus* (*Monocyttarus*) *mexicanus cubicola* in the United States. Florida Entomol. 67: 516-520.

53. Hermann, H. R., and J. T. Chao. 1984c. Morphology of the venom apparatus of *Mischocyttarus mexicanus cubicola* (Hymenoptera: Vespidae: Polistinae). J. Georgia Entomol. Soc. 19: 339-344.
54. Herre, E. A., D. M. Windsor, and R. B. Foster. 1986. Nesting associations of wasps and ants on lowland Peruvian ant-plants. Psyche 93: 321-330.
55. Hook, A. W., and H. E. Evans. 1982. Observations on the nesting behaviour of three species of *Ropalidia* Guérin-Ménéville (Hymenoptera: Vespidae). J. Aust. Entomol. Soc. 21: 271-275.
56. Hunt, J. H. 1988. Lobe erection behavior and its possible social role in larvae of *Mischocyttarus* paper wasps. J. Insect Behav. 1: 379-386.
57. Hunt, J. H., I. Baker, and H. G. Baker. 1982. Similarity of amino acids in nectar and larval saliva: the nutritional basis for trophallaxis in social wasps. Evolution 36: 1318-1322.
58. Itô, Y. 1983. Social behaviour of a subtropical paper wasp, *Ropalidia fasciata* (F.): field observations during founding stage. J. Ethol. 1: 1-14.
59. Itô, Y. 1984b. Social behaviour and social structure of neotropical paper wasps, *Mischocyttarus angulatus* Richards and *M. basimacula* (Cameron). J. Ethol. 2: 17-29.
60. Itô, Y. 1985a. A comparison of frequency of intra-colony aggressive behaviours among five species of polistine wasps (Hymenoptera: Vespidae). Z. Tierpsychol. 68: 152-167.
61. Itô, Y. 1985b. Colony development and social structure in a subtropical paper wasp, *Ropalidia fasciata* (F.) (Hymenoptera: Vespidae). Res. Popul. Ecol. 27: 333-349.
62. Itô, Y. 1985c. Social behaviour of an Australian paper wasp, *Ropalidia plebeiana*, with special reference to the process of acceptance of an alien female. J. Ethol. 3: 21-25.
63. Itô, Y. 1986a. On the pleometrotic route of social evolution in the Vespidae. Monit. Zool. Ital. (N.S.) 20: 241-262.
64. Itô, Y. 1986b. Social behaviour of *Ropalidia fasciata* (Hymenoptera: Vespidae) females on satellite nests and on a nest with multiple combs. J. Ethol. 4: 73-80.
65. Itô, Y. 1987a. Social behaviour of the Australian paper wasp, *Ropalidia revolutionalis* (de Saussure) (Hymenoptera: Vespidae). J. Ethol. 5: 115-124.

66. Itô, Y. 1987b. Role of pleometrosis in the evolution of eusociality in wasps. pp. 17-34 In Y. Itô, J. L. Brown, and J. Kikkawa (eds.), Animal Societies: Theories and Facts. Japan Scientific Societies Press, Tokyo.
67. Itô, Y., and S. Higashi. 1987. Spring behaviour of *Ropalidia plebeiana* (Hymenoptera: Vespidae) within a huge aggregation of nests. Appl. Entomol. Zool. 22: 519-527.
68. Itô, Y., and Sk. Yamane. 1985. Early male production in a subtropical paper wasp *Ropalidia fasciata* (Hymenoptera: Vespidae). Insectes Soc. 32: 403-410.
69. Itô, Y., O. Iwahashi, Sô. Yamane, and Sk. Yamane. 1985. Overwintering and nest reutilization in *Ropalidia fasciata* (Hymenoptera, Vespidae). Kontyû 53: 486-490.
70. Itô, Y., Sô. Yamane, and J. P. Spradbery. 1988. Population consequences of huge nesting aggregations of *Ropalidia plebeiana* (Hymenoptera: Vespidae). Res. Popul. Ecol. 30: 279-295.
71. Iwata, K. 1976. Evolution of Instinct: Comparative Ethology of Hymenoptera. Amerind, New Delhi.
72. Jarvis, J. U. M. 1981. Eusociality in a mammal: cooperative breeding in naked mole-rat colonies. Science 212: 571-573.
73. Jeanne, R. L. 1970a. Chemical defense of brood by a social wasp. Science 168: 1465-1466.
74. Jeanne, R. L. 1970b. Note on a bat (*Phylloderma stenops*) preying upon the brood of a social wasp. J. Mammal. 51: 624-625.
75. Jeanne, R. L. 1972. Social biology of the neotropical wasp *Mischocyttarus drewseni*. Bull. Mus. Comp. Zool. Harvard Univ. 144: 63-150.
76. Jeanne, R. L. 1975a. The adaptiveness of social wasp nest architecture. Quart. Rev. Biol. 50: 267-287.
77. Jeanne, R. L. 1979a. Construction and utilization of multiple combs in *Polistes canadensis* in relation to the biology of a predaceous moth. Behav. Ecol. Sociobiol. 4: 293-310.
78. Jeanne, R. L. 1979b. A latitudinal gradient in rates of ant predation. Ecology 60: 1211-1224.
79. Jeanne, R. L. 1980a. Evolution of social behavior in the Vespidae. Annu. Rev. Entomol. 25: 371-396.
80. Jeanne, R. L. 1981a. Chemical communication during swarm emigration in the social wasp *Polystia sericea* (Olivier). Anim. Behav. 29: 102-113.

81. Jeanne, R. L., and E. G. Castellan Bermudez. 1980. Reproductive behavior of a male neotropical social wasp, *Mischocyttarus drewseni* (Hymenoptera: Vespidae). J. Kansas Entomol. Soc. 53: 271-276.
82. Jeanne, R. L., H. A. Downing, and D. C. Post. 1983. Morphology and function of sternal glands in *Polistine* wasps (Hymenoptera: Vespidae). Zoomorphology 103: 149-164.
83. Keeping, M. G. 1984. A beetle predacious on the brood of a social wasp. J. Entomol. Soc. Sth. Africa 47: 355-358.
84. Keeping, M. G. 1990. Rubbing behaviour and morphology of van der Vecht's gland in *Belonogaster petiolata* (Hymenoptera: Vespidae). J. Insect Behav. 3: 85-104.
85. Keeping, M. G., and R. M. Crewe. 1983. Parasitoids, commensals and colony size in nests of *Belonogaster* (Hymenoptera: Vespidae). J. Entomol. Soc. Sth. Africa 46: 309-323.
86. Keeping, M. G., and R. M. Crewe. 1987. The ontogeny and evolution of foundation associations in *Belonogaster petiolata* (Hymenoptera: Vespidae). pp. 383-384 In J. Eder and H. Rembold (eds.), Chemistry and Biology of Social Insects. J. Peperny, Munich.
87. Keeping, M. G., D. Lipschitz, and R. M. Crewe. 1986. Chemical mate recognition and release of male sexual behavior in polybine wasp, *Belonogaster petiolata* (Hymenoptera: Vespidae). J. Chem. Ecol. 12: 773-779.
88. Kojima, J. 1982a. Nest architecture of three *Ropalidia* species (Hymenoptera: Vespidae) on Leyte Island, the Philippines. Biotropica 14: 272-280.
89. Kojima, J. 1982b. Notes on rubbing behavior in *Ropalidia gregaria* (Hymenoptera, Vespidae). New Entomol. 31: 17-19.
90. Kojima, J. 1983a. Defense of the pre-emergence colony against ants by means of a chemical barrier in *Ropalidia fasciata* (Hymenoptera, Vespidae). Jap. J. Ecol. 33: 213-223.
91. Kojima, J. 1983b. Occurrence of the rubbing behavior in a paper wasp, *Parapolybia indica* (Hymenoptera, Vespidae). Kontyû 51: 158-159.
92. Kojima, J. 1983c. Peritrophic sac extraction in *Ropalidia fasciata* (Hymenoptera, Vespidae). Kontyû 51: 502-508.
93. Kojima, J. 1984a. *Ropalidia* wasps in the Philippines (Hymenoptera, Vespidae). (I) Subgenus *Icarola*. Kontyû 52: 522-532.

94. Kojima, J. 1984b. Construction of multiple independent combs in *Ropalidia fasciata* (Hymenoptera: Vespidae). Jap. J. Ecol. 34: 233-234.
95. Kojima, J. 1984c. Division of labor and dominance interaction among co-foundresses on pre-emergence colonies of *Ropalidia fasciata* (Hymenoptera, Vespidae). Biol. Mag. Okinawa 22: 27-35.
96. Kojima, J. 1989. Growth and survivorship of preemergence colonies of *Ropalidia fasciata* in relation to foundress group size in the subtropics (Hymenoptera: Vespidae). Insectes Soc. 36: 197-218.
97. Kojima, J., and M. G. Keeping. 1985. Larvae of *Belonogaster juncea colonialis* Kohl and *B. petiolata* (Deger) (Hymenoptera: Vespidae). J. Entomol. Soc. Sth. Africa 48: 233-239.
98. Kukuk, P. F., G. C. Eickwort, M. Raveret-Richter, B. Alexander, R. Gibson, R. A. Morse, and F. Ratnieks. 1989. Importance of the sting in the evolution of sociality in the Hymenoptera. Ann. Entomol. Soc. Amer. 82: 1-5.
99. Lin, N., and C. D. Michener. 1972. Evolution of sociality in insects. Quart. Rev. Biol. 47: 131-159.
100. Lütte, M. 1977. Behavioral ecology of the social wasp, *Mischocyttarus mexicanus*. Behav. Ecol. Sociobiol. 2: 229-246.
101. Lütte, M. 1979. *Mischocyttarus flavitarsis* in Arizona: social and nesting biology of a polistine wasp. Z. Tierpsychol. 50: 282-312.
102. Lütte, M. 1981. Social biology of the polistine wasp *Mischocyttarus labiatus*: survival in a Colombian rain forest. Smithsonian Contr. Zool. 327: 1-27.
103. Marino Piccioli, M. T. 1968. The extraction of the larval peritrophic sac by the adults in *Belonogaster*. Monit. Zool. Ital. (N.S.) 2 (Suppl.): 203-206.
104. Marino Piccioli, M. T., and L. Pardi. 1970. Studi sulla biologia di *Belonogaster* (Hymenoptera, Vespidae). I. Sull'etogramma di *Belonogaster griseus* (Fab.). Monit. Zool. Ital. (N.S.) 3 (Suppl.): 197-225.
105. Marino Piccioli, M. T., and L. Pardi. 1978. Studies on the biology of *Belonogaster* (Hymenoptera Vespidae). 3. The nest of *Belonogaster griseus* (Fab.). Monit. Zool. Ital. (N.S.) 10 (Suppl.): 179-228.
106. Matthews, R. W. 1968a. *Microstigmus comes*: sociality in a sphecid wasp. Science 160: 787-788.

107. Michener, C.D. 1969. Comparative social behavior of bees. Annu. Rev. Entomol. 14: 299-342.
108. Michener, C. D., and D. J. Brothers. 1974. Were workers of eusocial Hymenoptera initially altruistic or oppressed? Proc. Natl. Acad. Sci. USA 71: 671-674.
109. Muralidharan, K., M. S. Shaila, and R. Gadagkar. 1986. Evidence for multiple mating in the primitively eusocial wasp *Ropalidia marginata* (Lep.) (Hymenoptera: Vespidae). J. Genet. 65: 153-158.
110. Oster, G. F., and E. O. Wilson. 1978. Caste and Ecology in the Social Insects. Princeton University Press, Princeton, NJ.
111. Page, R. E., and E. H. Erickson. 1988. Reproduction by worker honey bees. Behav. Ecol. Sociobiol. 23: 117-126.
112. Pardi, L. 1948. Dominance order in *Polistes* wasps. Physiol. Zool. 21: 1-13.
113. Pardi, L. 1977. Su alcuni aspetti della biologia di *Belonogaster* (Hymenoptera, Vespidae). Boll. Ist. Entomol. Univ. Bologna 33: 281-299.
114. Pardi, L., and M. T. Marino Piccioli. 1970. Studi sulla biologia di *Belonogaster* (Hymenoptera, Vespidae). 2. Differenziamento castale incipiente in *B. griseus* (Fab.). Monit. Zool. Ital. (N.S.) 3 (Suppl.): 235-265.
115. Pardi, L., and M. T. Marino Piccioli. 1981. Studies on the biology of *Belonogaster* (Hymenoptera Vespidae). 4. On caste differences in *Belonogaster griseus* (Fab.) and the position of this genus among social wasps. Monit. Zool. Ital. (N.S.) 14 (Suppl.): 131-146.
116. Post, D. C., and R. L. Jeanne. 1982b. Sternal glands in three species of male social wasps of the genus *Mischocyttarus* (Hymenoptera: Vespidae). J. New York Entomol. Soc. 90: 8-15.
117. Queller, D. C., and J. E. Strassmann. 1989. Measuring inclusive fitness in social wasps. pp. 103-122 In M. D. Breed and R. E. Page (eds.), The Genetics of Social Evolution. Westview, Boulder, CO.
118. Rasnitsyn, A. P. 1988. An outline of evolution of the hymenopterous insects (Order Vespida). Oriental Ins. 22: 115-145.
119. Ratnieks, F. L. W. 1988. Reproductive harmony via mutual policing by workers in eusocial Hymenoptera. Amer. Nat. 132: 217-236.

120. Ratnieks, F. L. W., and P. K. Visscher. 1989. Worker policing in the honeybee. *Nature* 342: 796-797.
121. Richards, O. W. 1962. A Revisional Study of the Masarid Wasps (Hymenoptera, Vespidae). British Museum (Natural History), London.
122. Richards, O. W. 1969. The biology of some W. African social wasps (Hymenoptera: Vespidae, Polistinae). *Mem. Soc. Entomol. Ital.* 48: 79-93.
123. Richards, O. W. 1971. The biology of the social wasps (Hymenoptera, Vespidae). *Biol. Rev. (Cambridge)* 46: 483-528.
124. Richards, O. W. 1978a. The Social Wasps of the Americas, Excluding the Vespinae. British Museum (Natural History), London.
125. Richards, O. W. 1978b. The Australian social wasps (Hymenoptera: Vespidae). *Aust. J. Zool.* 61 (Suppl.): 1-132.
126. Richards, O. W. 1982. A revision of the genus *Belonogaster* de Saussure (Hymenoptera: Vespidae). *Bull. Brit. Mus. (Nat. Hist.) Entomol. Ser.* 44: 31-114.
127. Richards, O. W., and M. J. Richards. 1951. Observations on the social wasps of South America (Hymenoptera Vespidae). *Trans. R. Entomol. Soc. London* 102: 1-170.
128. Ross, K. G. 1985. Aspects of worker reproduction in four social wasp species (Insecta: Hymenoptera: Vespidae). *J. Zool., London (A)* 205: 411-424.
129. Roubaud, E. 1916. Recherches biologiques sur les gpes solitaires et sociales d'Afrique. La genese de la vie sociale et l'volution de l'instinct maternel chez les vespides. *Ann. Sci. Nat. Zool.* (sr. 10) 1: 1-160.
130. Schwarz, M. P. 1988. Intra-specific mutualism and kin-association of cofoundresses in allodapine bees (Hymenoptera; Anthophoridae). *Monit. Zool. Ital. (N. S.)* 22: 245-254.
131. Seeley, T. D. 1985. Honeybee Ecology: A Study of Adaptation in Social Life. Princeton University Press, Princeton, NJ.
132. Sekijima, M., M. Sugijura, and M. Matsura. 1980. Nesting habits and brood development of *Parapolybia indica* Saussure (Hymenoptera: Vespidae). *Bull. Fac. Agric. Mie Univ.* 61: 11-23.
133. Spradbery, J. P. 1973a. Wasps: An Account of the Biology and Natural History of Social and Solitary Wasps. University of Washington Press, Seattle.

134. Spradbery, J. P. 1975. The biology of *Stenogaster concinna* van der Vecht, with comments on the phylogeny of *Stenogastrinae* (Hymenoptera: Vespidae). J. Aust. Entomol. Soc. 14: 309-318.
135. Starr, C. K. 1985b. Enabling mechanisms in the origin of sociality in the Hymenoptera - the sting's the thing. Ann. Entomol. Soc. Amer. 78: 836-840.
136. Starr, C. K. 1989b. In reply, is the sting the thing? Ann. Entomol. Soc. Amer. 82: 6-8.
137. Strassmann, J. E., and D. C. Meyer. 1983. Gerontocracy in the social wasp, *Polistes exclamans*. Anim. Behav. 31: 431-438.
138. Strassmann, J. E., C. R. Hughes, D. C. Queller, S. Turillazzi, R.ervo, S. K. Davis, and K. F. Goodnight. 1989. Genetic relatedness in primitively eusocial wasps. Nature 342: 268-269.
139. Stubblefield, J. W., and E. L. Charnov. 1986. Some conceptual issues in the origin of eusociality. Heredity 57: 181-187.
140. Sugjura, M., M. Sekijima, and M. Matsuura. 1983a. Intracolony polyethism in *Parapolybia indica* (Hymenoptera, Vespidae) I. Behaviour and its changes in the foundress and worker wasps in relation to colony development. Bull. Fac. Agric. Mie Univ. 66: 27-43 (in Japanese).
141. Sugjura, M., M. Sekijima, and M. Matsuura. 1983b. Life cycle of *Parapolybia indica* (Hymenoptera, Vespidae), with special reference to the colony development. Bull. Fac. Agric. Mie Univ. 66: 11-25 (in Japanese).
142. Suzuki, H., and M. Murai. 1980. Ecological studies of *Ropalidia [sic] fasciata* in Okinawa Island. I. Distribution of single- and multiple-foundress colonies. Res. Popul. Ecol. 22: 184-195.
143. Tinbergen, N. 1932. Über die Orientierung des Bienenwölfes (*Philanthus triangulum* Fabr.). Z. Vergl. Physiol. 16: 305-334. [Translated into English and reprinted in: 1972. The Animal in its World; Explorations of an Ethologist, Vol. I. Field Studies. Harvard University Press, Cambridge, MA. pp. 103-127.]
144. Tinbergen, N. 1935. Über die Orientierung des Bienenwölfes (*Philanthus triangulum* Fabr.). II. Die Bienenjagd. Z. Vergl. Physiol. 21: 699-716. [Translated into English and reprinted in: Tinbergen, N. 1972. The Animal in its World; Explorations of an Ethologist, Vol. I. Field Studies. Harvard University Press, Cambridge, MA. pp. 128-145.]

145. Tinbergen, N., and W. Kruyt. 1938. Über die Orientierung des Bienenwölfes (*Philaethus triangulum* Fabr.). III. Die Bevorzugung bestimmter Wegmarken. Z. Vergl. Physiol. 25: 292-334. [Translated into English and reprinted in: Tinbergen, N. 1972. The Animal in its World; Explorations of an Ethologist, Vol. I. Field Studies. Harvard University Press, Cambridge, MA. pp. 146-196.]
146. Trivers, R. L. 1971. The evolution of reciprocal altruism. Quart. Rev. Biol. 46: 35-57.
147. Trivers, R. L., and H. Hare. 1976. Haplodiploidy and the evolution of social insects. Science 191: 249-263.
148. Vecht, J. van der. 1941. The Indo-Australian species of the genus *Ropalidia* (=Icaria) (Hym., Vespidae). Treubia 18: 103-190.
149. Vecht, J. van der. 1962. The Indo-Australian species of the genus *Ropalidia* (Icaria) (Hymenoptera, Vespidae) (2nd part). Zool. Verhand. (Leiden) 57: 1-72.
150. Vecht, J. van der. 1965. The geographical distribution of the social wasps (Hymenoptera, Vespidae). Proc. 12th Int. Congr. Entomol., pp 440-441.
151. Vecht, J. van der. 1966. The East Asiatic and Indo-Australian species of *Polybioides* *Buysson* and *Parapolybia* *Saussure* (Hym., Vespidae). Zool. Verhand. (Leiden) 82: 1-42.
152. Vecht, J. van der. 1967. Bouwproblemen van sociale wespen. Verhand. K. Nederl. Akad. Wetens. Afd. Natuurk. 76: 59-68.
153. Vecht, J. van der. 1977a. Studies of Oriental *Stenogastriinae* (Hymenoptera: Vespidae). Tijdschr. Entomol. 120: 55-75.
154. Vecht, J. van der. 1977b. Important steps in the evolution of nest construction in social wasps. Proc. 8th Int. Congr. Int. Union Study Soc. Insects, p. 319.
155. Venkataraman, A. B., V. B. Swarnalatha, P. Nair, and R. Gadagkar. 1988. The mechanism of nestmate discrimination in the tropical social wasp *Ropalidia marginata* and its implications for the evolution of sociality. Behav. Ecol. Sociobiol. 23: 271-279.
156. Visscher, P. K. 1989. A quantitative study of worker reproduction in honey bee colonies. Behav. Ecol. Sociobiol. 25: 247-254.
157. West-Eberhard, M. J. 1975. The evolution of social behavior by kin selection. Quart. Rev. Biol. 50: 1-33.

158. West-Eberhard, M. J. 1978a. Polygyny and the evolution of social behavior in wasps. *J. Kansas Entomol. Soc.* 51: 832-856.
159. Wilson, E. O. 1971. *The Insect Societies*. Harvard University Press, Cambridge, MA.
160. Windsor, D. M. 1972. Nesting association between two neotropical *polybinae* wasps (Hymenoptera, Vespidae). *Biotropica* 4: 1-3.
161. Winston, M. L. 1987. *The Biology of Honey Bees*. Harvard University Press, Cambridge, MA.
162. Yamane, S., and J. Kojima. 1982. *Pseudonomadina* new genus, with a description of a new species from the Philippines (Hymenoptera, Trigonalidae). *Kontyû* 50: 183-188.
163. Yamane, S., and M. Terayama. 1983. Description of a new subspecies of *Bakeromyrmex typicus* Rohwer parasitic on the social wasp *Parapolybia varia Fabricius* in Taiwan (Hymenoptera: Trigonalidae). *Mem. Kagoshima Univ. Res. Ctr. South Pacific* 3: 169-173.
164. Yamane, S. 1980. Social Biology of the *Parapolybia Wasps* in Taiwan. Ph.D. dissertation, Hokkaido University, Sapporo.
165. Yamane, S. 1984. Nest architecture of two Oriental paper wasps, *Parapolybia varia* and *P. nodosa*, with notes on its adaptive significance (Vespidae, Polistinae). *Zool. Jahrb. Abt. Syst. Ökol. Geogr. Tiere* 111: 119-141.
166. Yamane, S. 1985. Social relations among females in pre- and postemergence colonies of a subtropical paper wasp, *Parapolybia varia* (Hymenoptera: Vespidae). *J. Ethol.* 3: 27-38.
167. Yamane, S. 1986. The colony cycle of the Sumatran paper wasp *Ropalidia (Icaridia) variegata jacobsoni* (Buysson), with reference to the possible occurrence of serial polygyny (Hymenoptera: Vespidae). *Monit. Zool. Ital.* (N.S.) 20: 135-161.
168. Yamane, S., and S. Yamane. 1979. *Polistine* wasps from Nepal (Hymenoptera, Vespidae). *Insecta Matsumurana* (N.S.) 15: 1-37.

pendix 5.1. Common Parasites of *Belonogaster*, *Mischocyttarus*, *Parapolybia*, and dependent-founding *Kopalidia*

Host species	Parasite	Locality	Stage affected	Remarks	Reference
<i>Belonogaster</i>	<i>Pediobius kopalidia</i> (Eulophidae: Hymenoptera)	Ghana, Africa	Pupa	High proportion of the pupae destroyed	Richards 1969
<i>B. juncea petiolata</i>	<i>Anacamptomyia</i> sp. (Tachinidae: Diptera) and <i>Campotyypus apicalis</i> (Ichneumonidae: Hymenoptera)	Transvaal, S. Africa	Prepupa and pupa		Keeping & Crewe 1983
<i>Mischocyttarus ewseni</i>	Strepsiptera	Brazil, S. America	Adult	Only 2 out of 760 adults were affected	Jeanne 1972
<i>flavitaris</i>	<i>Chalcoela iphitalis</i> (Pyralidae: Lepidoptera)	Arizona, USA	Larva and pupa	Common cause of brood mortality	Little 1979
<i>flavitaris</i>	<i>Monodontomerus</i> sp. (Torymidae: Hymenoptera)	Arizona, USA	Larva and pupa	Rare	Little 1979
<i>flavitaris</i>	Strepsiptera	Arizona, USA	Pupa and adult		Little 1979

Appendix 5.1 Contd

Host Species	Parasite	Locality	Stage affected	Remarks	Reference
<i>M. labiatus</i>	<i>Megaselia</i> sp. (Phoridae: Diptera)	Colombia, S. America	Brood	Responsible for a large proportion of nest failures than any other single factor	Litte 1981
<i>Parapolybia varia</i>	<i>Bakeronymus typicus</i> (Trigonaliidae: Hymenoptera)	Taiwan	Larva		Yamane and Terayama 1983
<i>Ropalidia cyathiformis</i>	Unidentified species of Ichneumonidae (Hymenoptera)	Bangalore, India	Larva		Gadagkar, unpubl.
<i>R. fasciata</i>	<i>Arthula formosana</i> (Ichneumonidae: Hymenoptera)	Okinawa, Japan	Larva	Very high percentage of cells parasitized	Ito 1983
<i>R. flavobrunea lapinga</i>	<i>Pseudonomadina biceps</i> (Trigonaliidae: Hymenoptera)	Philippine Islands	Brood	In addition to larvae and pupae, live adults of the parasite were found in active wasp nests	Yamane & Kojima 1982

Appendix 5.1 Contd

Species	Parasite	Locality	Stage affected	Remarks	Reference
<i>formosa</i>	<i>Hemipimpla pulchripennis</i> (Ichneumonidae: Hymenoptera)	Madagascar	Larva		Brooks & Wahl 1987
<i>marginata</i>	<i>Koralliomysia portentosa</i> (Tachinidae: Diptera)	Bangalore, India	Larva		Belavadi & Govindan 1981
<i>marginata</i>	Strepsiptera	Bangalore, India	Adult		Belavadi & Govindan, 1981, Gadagkar, unpubl.
<i>marginata</i>	Unidentified species of Tachinidae (Diptera), Toromyidae (Hymenoptera), Ichneumonidae (Hymenoptera)	Bangalore, India	Larva		K. Chandra shekara & Gadagkar unpubl.

Host species	Predator	Locality	Stage affected	Remarks	Reference
<i>Belonogaster petiolata</i>	<i>Hoplostomus fuliginus</i> (Scarabaeidae: Coleoptera)	S. Africa Transvaal,	Pupa		Keeping 1984
<i>Mischocyttarus drewsi</i>	<i>Monomorium pharaonis</i> , <i>Camponotus abdominalis</i> (Formicidae: Hymenoptera)	Brazil S. America	Brood		Jeanne 1972
<i>M. drewsi</i>	Spiders	Brazil, S. America	Adults away from the nest		Jeanne 1972
<i>M. flavitarsis</i>	Birds	Arizona, USA	Whole nest or pieces of the nest	Inferred	Little 1977
<i>M. flavitarsis</i>	Spiders, preying mantids	Arizona, USA	Adults away from the nest		Little 1977
<i>M. labiatus</i>	Army ants	Colombia, S. America	Brood	Seen only once	Little 1977
<i>M. mexicanus</i>	Ants, birds: blue jays and probably Carolina wrens, scrub jays, yellowthroats, cardinals, thrashers, mockingbirds, robins	Mexico	Whole nest or pieces of the nest	Predation by blue jays was observed and the involvement of the other birds was inferred	Little 1977
<i>Mischocyttarus</i> spp.	Bats	Brazil, S. America	Whole nest		Jeanne 1970b

Appendix 5.2 contd

Reference	Species	Predator	Locality	Stage affected	Remarks
Sekijima et al. 1980	<i>rapolybia dica</i>	<i>Vespa tropica</i> (Vespidae: Hymenoptera)	Japan	Pupa and larva	Most important enemy
Yamane 1980	<i>varia</i>	Ants	Taiwan	Brood	
Yamane 1980	<i>varia</i>	<i>Vespa tropica</i> (Vespidae: Hymenoptera)	Taiwan	Pupa and larva	Most important factor regulating population levels
Gadagkar unpubl.	<i>pallida</i> , <i>rginata</i> & <i>cyathiformis</i>	<i>Vespa tropica</i> (Vespidae: Hymenoptera)	Southern India	Pupa and larva	Most important factor regulating population levels