

New research finds that more than 90% of global aquaculture faces substantial risk from environmental change

Date: June 26, 2023

Source: University of California - Santa Barbara

Summary: Many of the world's largest aquatic food producers are highly vulnerable to human-induced environmental change, with some of the highest-risk countries in Asia, Latin America and Africa demonstrating the lowest capacity for adaptation, a landmark study has shown.

FULL STORY

Many of the world's largest aquatic food producers are highly vulnerable to human-induced environmental change, with some of the highest-risk countries in Asia, Latin America and Africa demonstrating the lowest capacity for adaptation, a landmark study has shown.

The study shows that more than 90% of global "blue" food production, in both capture fisheries and aquaculture, faces substantial risks from environmental change, with several leading countries in Asia and the United States set to face the greatest threats to production.

The authors behind the new paper produced the first-ever global analysis of environmental stressors impacting the production quantity and safety of blue foods around the world, ranking countries for the first time according to their exposure from key stressors.

A total of 17 stressors were surveyed, including algal blooms, sea level rise, changing temperatures and pesticide exposure.

"Environmental stressors do not care about national borders," noted Ben Halpern, co-lead author and professor at UC Santa Barbara and director of the National Center for Ecological Analysis and Synthesis.

"Stressors get moved by air, water, species, and humans, connecting land to sea and ecosystem to ecosystem."

The research published by *Nature Sustainability*, titled "Vulnerability of Blue Foods to Human-induced Environmental Change" is one of seven scientific papers published by the Blue Food Assessment (BFA) as part of a global effort to inform future aquatic food sustainability.

Alongside climate change, the report highlights that highly vulnerable blue food production systems are found across all continents, including some of the world's largest blue food producers such as Norway, China, and the United States, yet also argues there is too often a lack

of understanding around the complexity of stressors causing environmental change.

"We have only scratched the surface in our understanding of how environmental stressors are connected, and how they can both negatively impact the production and safety of the resulting blue foods," said Ling Cao, co-lead author and professor at the State Key Laboratory of Marine Environmental Science at Xiamen University.

"Understanding the complexity of these stressors, and their cascading impacts, will be essential in developing successful adaptation and mitigation strategies," she said.

Species invasion, inland eutrophication or algal blooms, ocean warming and sea level rise were cited by the paper as the main threats to blue food production in the U.S., with freshwater and marine fisheries facing disproportionately large risks.

As the largest blue food producer, China's freshwater aquaculture is also highly exposed to inland eutrophication and severe weather events, the research shows.

The authors also argue that special attention should be paid to countries facing high exposure to environmental change yet not possessing adequate capacity for adaptation, including Bangladesh, Eswatini, Guatemala, Honduras and Uganda.

In terms of production systems, the paper finds that marine fisheries were generally more vulnerable to climate-related stressors, particularly rising temperatures and acidification, whilst aquaculture was more susceptible to the effects of diseases and hypoxia, or low oxygen levels.

"Although we have made some progress with climate change, our adaptation strategies for blue food systems facing environmental change are still underdeveloped and need urgent attention," said Rebecca Short, co-lead author and researcher at the Stockholm Resilience Centre.

Among the report's key recommendations is a call for more transboundary collaboration and adaptation strategies which recognize that the ecosystems that blue food production rely upon are highly interconnected, with environmental change in one area having potential knock-on effects elsewhere.

The authors also call for a diversification of blue food production in high-risk countries to cope with the impact of environmental change unless sufficient mitigation and adaptation strategies are adopted.

Likewise, the paper highlights the urgent need for greater stakeholder engagement in understanding, monitoring and mitigating pressures on blue food production systems.

Indigenous knowledge will be critical for strategic planning and policies to mitigate and adapt to environmental change, particularly for artisanal fisheries and heavy marine fisheries-dependent countries, such as Small Island Developing States (SIDS).

The research also includes an extended dataset which ranks countries around the world based on the exposure of their blue food production systems to the various environmental stressors.

Story Source:

[Materials](#) provided by **University of California - Santa Barbara**. Original written by Sonia Fernandez. *Note: Content may be edited for style and length.*

Journal Reference:

1. Ling Cao, Benjamin S. Halpern, Max Troell, Rebecca Short, Cong Zeng, Ziyu Jiang, Yue Liu, Chengxuan Zou, Chunyu Liu, Shurong Liu, Xiangwei Liu, William W. L. Cheung, Richard S. Cottrell, Fabrice DeClerck, Stefan Gelcich, Jessica A. Gephart, Dakoury Godo-Solo, Jessie Ihilani Kaull, Fiorenza Micheli, Rosamond L. Naylor, Hanna J. Payne, Elizabeth R. Selig, U. Rashid Sumaila, Michelle Tigchelaar. **Vulnerability of blue foods to human-induced environmental change.** *Nature Sustainability*, 2023; DOI: [10.1038/s41893-023-01156-y](https://doi.org/10.1038/s41893-023-01156-y)

Cite This Page:

MLA

APA

Chicago

University of California - Santa Barbara. "New research finds that more than 90% of global aquaculture faces substantial risk from environmental change." ScienceDaily. ScienceDaily, 26 June 2023. <www.sciencedaily.com/releases/2023/06/230626164317.htm>.

Explore More

from ScienceDaily

RELATED STORIES

Study Into Global Daily Air Pollution Shows Almost Nowhere on Earth Is Safe

Mar. 7, 2023 — In a new study of daily ambient fine particulate matter (PM2.5) across the globe, a new study has found that only 0.18% of the global land area and 0.001% of the global population are exposed to ...

Biodiversity in Africa and Latin America at Risk from Oil Palm Expansion, New Report Warns

Nov. 28, 2022 — Zero deforestation commitments may inadvertently leave vital habitats in Latin America and Africa vulnerable to agricultural expansion, a new study has ...

The Use of Wild Mammals in Traditional Medicine

Dec. 9, 2020 — Investigators identified 565 mammalian species that have been used to source products used in traditional medicine around the world, especially in Asia, Africa, and Latin ...

The Dangers of Collecting Drinking Water

Nov. 4, 2020 — Fetching drinking water in low and middle income countries can cause serious injury, particularly for women. A new study reveals dangers including falls, traffic accidents, animal attacks, and ...