


INDIAN INSTITUTE OF SCIENCE Centre for Ecological Sciences BANGALORE-560 012, INDIA



GEOGRAPHICAL DISTRIBUTION OF PLANT SPECIES IN KARNATAKA

MADHAV GADGIL, C. J. SALDANHA, SURESH H. S. MALATI HUGDE AND VENKAT RAO

January 1987

Technical Report No. 12

INTRODUCTION

This is a report of our attempt to build up a computerised data base on geographical distribution of flowering plants in Karnataka. It constitutes part of a larger programme which in addition to Karnataka would eventually include Nilgiri biosphere reserve and Peninsular India. The information for this data base has been taken from "Flora of Karnataka" Vol. 1 by Prof. C. J. Saldanha (appendix-1)

Information has been coded to cover 6 fields which are as follows:

- 1) Zone
- 2) Altitude
- 3) Habitat
- 4) Life form
- 5) Abundance
- 6) Phytogeographic affinity

To each of these variables as far as they apply to Karnataka have been identified and coded as far as phytogeography of the plants concerned. Total number of tax a coded in 64 families is 1100.

Methodology

To prepare a complete account of geographical distribution of plants in Karnataka work was taken up in 2 steps.

- 1. Proparation of thick list of plants
- 2. Collecting various information available for each plant species, organizing them in systematic way

- I. Preparation of check list: Firstly check list of the Indian families was prepared and arranged in alphabetical order. Each family was assigned a code number, for easy retrieval. Out of this first 63 families of cronquest system of classification were taken for the analysis, the list is given in appendix. Check list of all the recorded species of plants of peninsular India was prepared then assigned a code number for every genus and species. Model sheet is inclued in appendix-1.
- 2. <u>Information</u> :- Various information collected for 63 families are explained below:-

This is the crucial part of work. Information about the plant was collected under 6 different heads which are important and obvious.

Zone: Zones were made for the Geographical region of Karnataka.

Karnataka was divided into 6 broad ecological zones and assigned code. They are

- 01 Coastal littoral
- 02 Coastal low lands
- 03 Up Ghats
- 04 Western plateau
- 05 Northern plateau
- 06 Southern plateau
- 07 All regions

Coastal littoral: (01) This region represents coastal belt of 320 kms with a average width of 20 kms. The coast in often soudy and occasionally with rocky shore average rainfall 250 cms supports than dume and mangrage veget them.

Coastal low lands: (02) This zone is not extensive in Karnataka. It extends from 5 km north (Uttara Kannada) to 65 km in south (south kanara) receives rainfall on an average of 250 cm. The soil is fertile alluvial with latentic on exposed surfaces. This zone has agriculture belt as well as patches of evergreen forest/evergreen scrub.

Up Ghats: (03) This is a chain of mountains called Sahyadris with an average elivation of 900 m and deep valleys of 200 M with peak raised to 1827M (Kudremukh). This range has two faeles wind ward and leeward.

Wind ward had which appears plunging into sea near Karwar receives good rain fall from south west monsoon and supports luxuriant vegetation.

Leeward side which is undulating hills receives lesser rainfall supports moist deciduous forest. We have included Bababudan range in this category. They represent separate chain of mountains with an average altitude of 1400M, highest 1923 M SL(Mullainagiri) very close to Western Ghats Vegetation pattern similar.

Soil is lateritic. Along with the forest belt this region harbours rich Areca plantations and paddy fields.

Western plateau: (04) This is a strip at eastern foot of the ghats and a part of Deccan plateau includes parts of Belgaum, Dharwar, Shimoga, Chikmagalur, Hassan and Mysore. We have included BillyIrlrangana hills in this region, which is at the southern tip of this region, join western ghats in Nilgiris and supports luxurient vegetation from dry scub to shola-grass land.

Morthern idateau (05) It is undulating plateau of elevation 309 450M, a vant stretch of Monotonous Lands with inclated hills

leeping with low elevation. Soil mainly deep black and modium black. It is a prominant irrigation belt of the state natural vegetation very less except for dry scrub forest.

Southern plateau: (06) Undulating much higher area of mean elevation 800M with isolated low elevation ranges, major one Shivana Samudram to Bellary. Soil type is red soil with isolated patches of deep red and deep red and black soil. Matural vegetation scanty except for dry deciduous and scrub. For each region seperate code number is given last code number 07. All region represents entire state.

Altitude Karnataka has charactistic altitudinal range. It ranges from 0 to M to 1923M. this is divided into 4 class.


Туро	Code No.
0.300	01
250800	02
800-1400	0.3
1200 and above	. 04

This system is a standarised system used by french Institute, Pondicherry also.

0-300,250-850,850-1200- range represent tropical vegetation belt and 1200 and above range represents tropical mountaine vegetation type(shola-grass land vegetation type).

Habitat: This is dwelling place of plant. This can be natural or

artificial(Nan-made). We have under this 22 categories. For convenience they can be arranged



Forests: Broad classification of Indian forests is based on climatic consideration by MaG. Champion 1936.

Another approach to classification by Legris & Pascal(1964) and more detailed mapping of the region in 1982.

Both the approaches agree that two basic types exists.

- 1. Moist forest type
- 2. Dry forest type

Moist forest type includes evergreen, semievergreen and moist deciduous.

Evergreen and semievergreen

Lower slopes and valleys of western ghats are good examples.

They are close canopy forests and nurtured by south west monsoon.

Main associations are Dipterocarpus - Kingledendron - Vatoria.

Dipterocarpus - Mesua - Palaquium, Etc.

Ecological importance in considerable and disturbance would result in schievergreen forests.

Moist deciduous type

Decreasing rain fall on the leeward side results in moist deciduous forests. Canopy will be dense while in leaf. Lary's fall during dry season. This belt runs from Belgaum to Kodaqu. Main association - Tectona-Dillenia-Lagerstroemia-Terminalia.

Dry evergreen type not represented in Karnataka.

Dry Type

Dry Deciduous: Bioclimate of Ostern part of Maidan permits climax veg tation Dry Deciduous. Canopy is open and leafless during dry months. Major associations include

Anogeissus- Tenminalia-Tectona ,

Ancquissus-Hardwickia,

Anogeissus-Chloroxylon Albizia and

Albizia - Acacia

Scrub: They are dry forests in several parts of Karnataka. Trees with tiny leaflets and spines. Diminent tree is Acacia.

Grass Land: In Kurnataka grass lands are in high altitudes of Ghats, Bababudan hills and Biligirirangana hills.

Lithophytes: Lithophytes are those which grow on rocks and rocky formations.

Non-cultivated Lands: They constitute waste places which is usually occupied by exotics and weeds.

Sandy Beach: This is made up of characteristic vegetation - sand dune vegetation which is psamaphytic has direct sun, salt laiden winds and shifting sand. There are about 99 species of plants we

recorded (A.G. Untawale and Sayeda Wafar, 1986). Main species includes Ipom capuscapre, Spinifix littorees.

Presh water habitats include rivers: streams, rivulats, etc in the flowing fresh water category. Pends, Tanks and ditches are included in standing fresh water category. Marshy area surrounding this, is Fresh water marsh.

Marine habitats include mangrove and related vegetation.


 $\underline{\text{Man made:}}$ This includes the manipulated habitats which is created by man. They include:


Agriculture includes irrighted, nonirrighted and fall ws. Plantations includes forestry and demostic.

Parks and gardens, road sides/hedges, railway tracks.

So all the habitat categories can be listed with their excess(appendix-2).

Life form: Life form of a plant is defined by stature, structur, amount of biomass they put in and their life cycles This class has 13 categories. They can be arranged for convenience.





They can be listed with their respective codes. (appendix-2).

Abundance: Abundance is number of representatives of that species. It is the qualitative quantification. discussed under 7 different headings as given below.

Code	Heading
01	Absent
02	Ruro
03	Frequent
04	Common
05	Locally abundant
06	Distribution in patches
07	Distribution scattered

01 Absent This does not fit into present analysis but when analysis includes peninsular India then it will be useful.

<u>Mare</u> In which species restricts itself to an area or distributes scantly around an area.

Common and locally abundant-distinction is made by the fact that common stands for commoness of species all along the state and locally abundant - abundant in that particular area where appears are collected.

Phyto Geographic Affinity It is the distribution of species along the global scale or otherwise the limit of the species. Claute last

evolve! through equs. Our continent "Gondwane" has also undergone several quotagical and quantific to add changes. Of his have also correspondingly responded to the changes.

Affinities are discussed under 7 herds.

Civile	Head.	1 '
01	Endemic	Ve
02	Indo arabic	•
03	Indo malayan	
04	Himalayan	
05	Introduced	
06	Nootropical	•
07	Pantropical	

Endemic: (01) Having one restricted area (f distribution.

Indoarabic: (02) Which is related to the parts of Africa and dry areas of Pakistan atc. include Arabia also.

Indo molayan: (03) It is a continuous bolt of relating areas such as Srilanka, Peninsular India, Burma, South East Asia, Helaya, etc.
Introduced: (05) Introduction by man, sime do grow as uscape but not yet naturalised.

Neotropical: (06) From central American countries which have become naturalised.

Pantropical: (07) Which has distribution of entire tropics.

This would give a brief account of parameters chosen for the organisation of the information.

Analysis: Central computational facility Doc-10 system is usel for

analysis. Typical calabiling inflamation for a species really Achyminthus is a late.

0080101, 07, 019203. 2022, 02, 04, 07, ACHYRANTHUS ASPERA

We will just try to mid so with up old s

008 0101 - stands for the hame of the plant species

Achyranthus aspera of Amaranthaceae

07 - stands for zones - All regions

010203 - Altitude - 0-1400M

2022- Habitat - Rhadsides/hedges and non culltivated lands

02 - life form - Perronial herb

04 - Abundance - Common

07 - Phytogoographic affinity - Pantropical

Information Retrieval

From the information gathered on various parameters following kind of information can be retrieved.

Following are the few that can be answered.

1. How many families are represented in each gargraphical zero?

Zines	01	02	03	04	05	06
No. of Comilies	36	38	55	57	42	53
2 tofa rulics rulics rules unted	57	50 3	87.3	90.5	66.7	R4.1

2 - What is the species insit - Cooch polymorphic ነበን

Z⊝ne	01	02	03	04	05	06
No of species	143	163	517	197	281	387
% of spe cies	13.9	15.9	53.2	48.3	27.3	37.6
Density logS/ logA	.5979	. 6757	.7058	.6386	.4927	. 4547

S = N. If spucios

 Λ = Geographical area in sq Km

- 3. Which is the zone having highest species density level? "Upghats"
- 4. What is the percentage of species distribution for a given family in each geographical zones? For example if one wants to know a of species occurance for the families Amaranthaceae and Lauraceae.

Families	01	02	03	04	05	06
Amaranth aceae	50%	43.8%	65.6%	50%	71.9%	75%
Lauraceae	4.8%	9.5%	92.98	11.98	2.48	9.5%

5. What is the composition of a particular zones in terms of life forms. For example, if we want to know composition of constal littoral and upper ghats.

	Castal Li No.	8	Upper Ne	Ghats %			
Annual herb	35	25.5	75	14.1			
Perrenial herb	33	24.1	73	13.7			
Climbing herb	9	6.6	40	7 : 5			
Climbing shrub	5	; 3,6	50	9,4			
Understarcy shrub	0	0	16	3.0			
Exprise + shrub	22	15.1	68	12.8			
Troc	31	22.6	203	, 38 1			
Epiphytos	0	0	6	1.1			
H.1.S.P	0	0	0	0			
H.T.R.P.)	7	1	•2			
H.P.S.P.	0	0	0	0			
H.P.R.P.	1	٠7	1	0.2			
Shprophyte	0	Ö	0	0			
6. What is the percent	age of:	species in	each	zone nt m			
particular_altitude_class?							
Altitude	01	02		03 91			
Z enes							
01	43.2	8.6		9.0 3 1			
02	46.8	10.0		10.0 5 2			

	Altitude		01	02	03	31
2 ones						
01			43.2	8.6	9.0	3 1
02			46.8	10.0	10.0	5/2
03		ŧ	49.2	36.1	35.9	72.1
01			37.9	34.2	35.2	22.4
05	×		31.6	19.5	19.9	6.9
06			35.1	25.9	29.48	6.9

- 7. Which are the families having representations in all zones of Karnataka?
- 1. Aiz aceae
- 2. Amaranthaceae
- 3. Annonaceae
- 4. Aristolochiaceae
- 5. Capparacene
- Coratophyllaceae
- 7. Chenery xliaceae
- 8. Dill miaccae
- 9. Dri democre
- 14 Ebonaculad
- Habitac carracond
- 12 all above no
- 13.Lauraceae
- 14. Locythichcome
- 15.Mhgnoliacuae
- 16.Malvacene
- 17.Monispermaceae
- 18.Moracoac
- 19.Papavoraceae
- 20. Pittosporaceae
- 21.Plumbaginaceae
- 22.Polygonaceae

- 23.Partulachecau
- 24.Ranunculaceae
- 25.Sapotacuae
- 26.Sterculiaceae
- 27.Tiliaceae
- 28.Urticaceae
- 29.Violaceae

APPENDIX-1:LIST OF FAMILIES

ee5 ATZOACEAE AMARANTHACEAE $\{\hat{\mathcal{G}}(t)\}$ ANCISTROCLADACEAE $\Theta \downarrow \Theta$ ANNONACEAE 011 ARISTOLOCHIACEAE 919 026 BEGONIACEAE 030 BIXACEAE 031 DOMBACACEAE 033 BRASSICACEAE 038 CACTACEAE CAPPARACEAE 042 CARYOPHYLLACEAE 046 050 CERATOPHYLLACEAE 051 CHENOPODIACEAE 058 CLUSIACEAE 067 CRASSULACEAE 070 CUCURBITACEAE 074 DATISCACEAE 077 DILLENIACEAE DIPTEROCARPACEAE 681 092 DROSERACEAE 083 EBENACEAE 085 ELACOCARPACEAE 096 FABACEAE/CAESALPINIACEAE 097 FABACEAE/MIMOSOIDEAE Ø 98 **FABACEAEZPAPLIONACEAE** 100 FLACOURTIACEAE 110 REPNANDIACÉAE 126 LAURACEAE LECYTHIDACEAF 127 137 MAGNOL TACEAL 139 MALVACEAE 144 MENTSPERMACEAE 1.13 HURACEAS USG MYTTLATICACEAE MYRS THACFAR 154 157 HET RHIBOUNACE VE $J_{\rm c} (t, t, t)$ MYCTAGINACEAF 16 1 MAMORINES - EVE 165OCHNACEAE PAPAVERACEAE PASSIFLORACEAE 175 PHYTOLACCACEAE 177 178 PIPERACEAE 177 PITTOSPORACEAE

182

187

189

196

200 204

205

PLUMBAGINACEAE

PULYGENACIÉAE

ROSACEAE,

SABIACEAE

SALICACEAE

PORTULACAČEAE RANUNCULACEAE

SAPOTACEAE SAXIFRAGACEAE STERCUL TACEAE SYMPLOCACEAE TAMARICACEAE 230 THEACEAE 233 TIL IACEAE 237 TURNERACEAE 239 LILMACEAE 246 URTICACEAE VIOLACEAE

APPENDIX-2 : HABITAI CATAGORIES

- 01 EVERGREEN/SCHIEVERGREEN
- 02 HOIST DECIDUOUS
- 93 DEGRADED EVERGREEN
- 94 DEGRADED DECIDIOUS
- 05 SCRUB
- 96 FLOWING FRESH WATER
- 97 STANDING FRESH WATER
- 08 FRESH WATER MARSH
- 09 BRACKISH WATER
- 10 DRACKISH MARSH
- 11' SANDY BEACHES
- 12 LITHOPHYTES
- 13 NATURAL GRASS LANDS
- 14 IRRIGATED
- 15 NON IRRIGATED
- 16 FAULOW
- 17 PARKS AND GARDENS
- 18 FORESTRY PLANTALLON
- 19 NOW FORESTRY PLANTATION
- 139 ROAD SIDES/HEDGES
- 21 RAILWAY TRACKS
- 22 NON CULTIVATED LAND

APPENDIX-3 : LIFE FORM CATAGORIES

- 01 ANNUAL HERB
- 02 PERENNIAL HERB
- 03 CLIMBING HERB
- ed CLIMBING SHRUE
- 95 UNDER STOPEY SHRUD
- 06 EXPOSED SHRUB
- 97 TREE

₽,

- 08 EPIPHYTE
- 09 HETEROTROPHIC TOTAL STEM PARASITE
- 10 HETEROTROPHIC TOTAL ROOT PARASITE
- 11 HETEROTROPHIC PARTIAL STEM PARASITE
- 12 HETEROTROPHIC PARTIAL ROOT PARASITE
- 13 SAPROPHYTE