

# **GLOBAL AND NATIONAL SOILS AND TERRAIN DIGITAL DATABASES (SOTER)**

**Procedures Manual**



FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED  
NATIONS



# GLOBAL AND NATIONAL SOILS AND TERRAIN DIGITAL DATABASES (SOTER)

## Procedures Manual



United Nations Environment Programme



International Society of Soil Science



International Soil Reference and Information Centre



Food and Agriculture Organization of the  
United Nations

[Copyrights and disclaimer]

## Preface

Based on a discussion paper "Towards a Global Soil Resources Inventory at Scale 1:1M" prepared by Sombroek (1984), the International Society of Soil Science (ISSS) convened a workshop of international experts on soils and related disciplines in January 1986 in Wageningen, the Netherlands, to discuss the "Structure of a Digital International Soil Resources Map annex Data Base" (ISSS, 1986a). Based on the findings and recommendations of this workshop a project proposal was written for SOTER, a World SOils and **TER**rain Digital Data Base at a scale of 1:1 million (ISSS, 1986b).

A small international committee was appointed to propose criteria for a "universal" map legend suitable for compilation of small scale soil-terrain maps, and to include attributes required for a wide range of interpretations such as crop suitability, soil degradation, forest productivity, global soil change, irrigation suitability, agro-ecological zonation, and risk of droughtiness. The committee compiled an initial list of attributes. The SOTER approach received further endorsement at the 1986 ISSS Congress in Hamburg, Germany.

A second meeting, sponsored by the United Nations Environment Programme (UNEP), was held in Nairobi, Kenya, in May 1987 to discuss the application of SOTER for preparing soil degradation assessment maps. Two working groups (legend development and soil degradation assessment) met concurrently during this meeting. The legend working group was charged with the task of developing Guidelines for a World Soils and Terrain Digital Database at a 1:1 M scale, to propose general legend concepts, to prepare an attribute file structure, and to draft an outline for a Procedures Manual (ISSS, 1987).

Following the Nairobi meeting, UNEP formulated a project document: "Global Assessment of Soil Degradation" and asked ISRIC to compile, in close collaboration with ISSS, FAO, the Winand Staring Centre and the International Institute for Aerospace Survey and Earth Sciences (ITC), a global map on the status of human-induced soil degradation at a scale of 1:10 million, and to have this accompanied by a first pilot area at 1:1 million scale in South America where both status and risk of soil degradation would be assessed on the basis of a digital soil and terrain database as envisaged by the SOTER proposal. In this context ISRIC subcontracted the preparation for a first draft of a Procedures Manual for the 1:1 M pilot study area to the Land Resource Research Centre of Agriculture Canada<sup>1</sup>.

The first draft of the Procedures Manual (Shields and Coote, 1988) was presented at the First Regional Workshop on a Global Soils and Terrain Digital Database and Global Assessment

---

<sup>1</sup> Presently the Centre for Land and Biological Resources Research

of Soil Degradation held in March 1988 in Montevideo, Uruguay (ISSS, 1988). The proposed methodology was then tested in a pilot area, covering parts of Argentina, Brazil and Uruguay (LASOTER). Soil survey teams of the participating countries collected soils and terrain data to assess the workability of the procedures as proposed in the draft Manual. During two correlation meetings and field trips minor changes were suggested, while further modifications were recommended at a workshop that concluded the data collection stage. The comments from both workshops were incorporated in the January 1989 version of the Procedures Manual (Shields and Coote, 1989).

Application of the SOTER methodology in an area along the border between the USA and Canada (NASOTER), revealed additional shortcomings in the second version of the Manual. Also, the first tentative interpretation of the LASOTER data as well as the integration of the attribute data into a Geographic Information System demonstrated the need for further modifications.

A third revised version of the Manual was compiled by the SOTER staff (ISRIC, 1990a) and circulated for comments amongst a broad spectrum of soil scientists and potential users of the database. A workshop on Procedures Manual Revisions was convened at ISRIC, Wageningen, to discuss the revised legend concepts and definitions (ISRIC, 1990b).

Based on the recommendations of this workshop, the proposed modifications were further elaborated, resulting in a fourth draft version of the Procedures Manual (ISRIC, 1991). This Manual consisted of three parts, the first of which dealt with terrain and soil characteristics. The second part treated land use in a summary way in the expectation that a more comprehensive structure for a land use database would become available from other organizations. In the third part information on related files and climatic data needed for SOTER applications were described. In each section definitions and descriptions of the attributes to be coded were given, while in the first section an explanation of the mapping approach was provided.

Unlike the 1st and 2nd versions of the Manual, the later versions did not elaborate upon the soil degradation assessment as this is considered to be an interpretation of the database. Guidelines for this and other interpretations will be subject of separate publications. Technical specifications (e.g. table definitions, primary keys, table constraints etc.) and a user manual for the SOTER database will also be published separately.

A second SOTER workshop organized by UNEP was convened in February 1992 in Nairobi. At this meeting FAO expressed its full support for the SOTER programme and indicated that it was prepared to use the SOTER methodology for storing and updating its own data on world soil and terrain resources. To facilitate the use of SOTER data by FAO it was decided to use the FAO-Unesco Soil Map of the World Revised Legend (FAO, 1988) as a basis for characterising the soils component of the SOTER database.

To take account of these decisions a fifth version of the Manual was prepared in 1992 with active participation by FAO. The main arrangement of this latest version of the Manual is similar to the fourth version, with the difference that the Manual now consists of two parts only, the first one dealing with soils and terrain, and the second one dealing with the accessory databases in which land use, vegetation and climatic data can be stored.

No further revisions of the Manual are planned until more experience has been gained in the application of the methodology according to the current guidelines. Nevertheless, all

comments are welcome, and should be sent to the Manager of the SOTER project<sup>1</sup>.

Vincent van Engelen  
Wen Ting-tiang  
editors

**Note with the 1995 revised edition**

This version incorporates some additional attributes in the horizon part of the database related to soluble salts. Also FAO soil units of 1988 have been added as an annex. No other changes have been made with respect to the 1993 version.

The editors.

---

<sup>1</sup> c/o Director, International Soil Reference and Information Centre, P.O.Box 353, 6700 AJ Wageningen, The Netherlands.

## Acknowledgements

The SOTER project, an initiative of the ISSS, was very effectively supported by Working Group DM of the ISSS under chairmanship of M.F. Baumgardner. The project has benefited enormously from the experience of a wide range of soil and other natural resource scientists from all over the world. Our special thanks go to the following persons who were very active in the compilation of the manual:

D.R. Coote  
J.H.M. Pulles  
J. Shields

Participation of the following persons in the various workshops and pilot area studies is also highly appreciated:

C. Alvarez, D. Arnold, A. Ayoub, N.H. Batjes, M.F. Baumgardner, P. Brabant, R. Brinkman, P.A. Burrough, T. Calhoun, A. Califra, C. Clerici, T.T. Cochrane, R.M. Di Giacomo, E. Di Landro, P.J. Fasolo, N. Fernandez, I.P. Garbouchev, R. Godagnone, M. Ilaiwi, E. Klamt, J. Lamp, K.B. MacDonald, J.H. Molfino, F.N. Muchena, F. Nachtergaele, L.R. Oldeman, J. Olmos, W.L. Peters, C. Petraglia, R. Pötter, M.F. Purnell, W. Reybold, J.C. Salazar, C. Scoppa, J.L. Seghal, D. Sims, W.G. Sombroek, R.F. van de Weg, G. Varallyay, D. Yost, J.A. Zinck.



# Contents

	Page
PREFACE	iii
ACKNOWLEDGEMENTS	iv
<b>PART I: SOILS AND TERRAIN</b>	<b>1</b>
1 General introduction	3
Aim	3
Central database	3
Characteristics	3
Procedures	4
2 Mapping approach and database construction	5
SOTER mapping approach	5
SOTER source material	7
Associated and miscellaneous data	8
3 SOTER differentiating criteria	9
Terrain	9
Terrain components	10
Soil components	11
SOTER unit mappability	13
SOTER approach at other scales	13
4 SOTER database structure	15
Geometric database	15
Attribute database	17
5 Additional SOTER conventions	21
SOTER unit codes	21
Minimum size of the SOTER unit	22
Number of soil and terrain components	22
Representative soil profiles	22
Updating procedures	23
6 Attribute coding	25
Terrain	25
Terrain component	31

---

	Page
Terrain component data	32
Soil component	35
Profile	39
Horizon data	41
<b>PART II: LAND USE AND VEGETATION</b>	<b>59</b>
7 Land cover	61
Land use	61
Vegetation	62
<b>PART III: MISCELLANEOUS FILES</b>	<b>67</b>
8 Reference files	69
Source map	69
Laboratory information	70
Soil profile database	71
9 Climate	73
Climate station	73
Climate data	74
Various climate characteristics	75
Additional conventions	77
Data sources	77
ANNEX 1 Hierarchy of landforms	79
ANNEX 2 FAO soil unit codes	85
ANNEX 3 Hierarchy of land use	89
ANNEX 4 Hierarchy of vegetation	93
ANNEX 5 ISO country codes	99
ANNEX 6 SOTER data entry forms	101
REFERENCES	119
GLOSSARY	123
RELATED PUBLICATIONS	124

## List of figures

	Page
1 Relations between a SOTER Unit and their composing parts and major separating criteria	6
2 Terrain subdivided according to major landforms	10
3 Terrain further subdivided according to lithology	10
4 Terrain components differentiated according to surface forms	10
5 Terrain components differentiated according to slope gradients	10
6 SOTER units after differentiating soils	11
7 SOTER units, their terrain components, attributes, and location	16
8 SOTER attribute database structure with area and point data	16
9 Examples of slightly dissected and dissected landscapes as indicated by the density of the drainage on 1:50 000 maps	30
10 Texture groups of parent material	34
11 Texture classes of fine earth	54

## List of tables

1 Non-spatial attributes of a SOTER unit	18
2 Hierarchy of major landforms	26
3 Hierarchy of lithology	30
4 Size classes for structure elements of various types	52
5 Attributes of land use and vegetation files	61
6 Hierarchy of land use; land use orders, groups, and systems	62
7 Hierarchical vegetation classes	63
8 Related tables	70
9 Attribute list for climate station and climate data files	74
10 Example of various kinds of climatic data recorded for a climate station	77

# **PART I**

## **SOILS AND TERRAIN**

## Chapter 1

# General introduction

### AIM

The aim of the SOTER project is to utilize current and emerging information technology to establish a World Soils and Terrain Database, containing digitized map units and their attribute data (ISSS, 1986b). The main function of this database is to provide the necessary data for improved mapping and monitoring of changes of world soil and terrain resources.

It is composed of sets of files for use in a Relational DataBase Management System (RDBMS) and Geographic Information System (GIS). It is capable of delivering accurate, useful and timely information to a wide range of scientists, planners, decision-makers and policy-makers.

### CENTRAL DATABASE

In the initial phases of the SOTER project no concrete plans have been formulated for the physical establishment of a centralized database. Rather, a separate database will be set up for each area for which a land resource inventory is being undertaken according to the SOTER methodology. The common approach does, however, guarantee the possibility of merging the individual databases into a global database if and when this becomes feasible. Through its basic activities SOTER also intends to contribute to the establishment of national and regional soil and terrain databases, founded upon the same commonly acceptable principles and procedures, so as to further facilitate the exchange of land resource information and ultimate incorporation into a global database.

### CHARACTERISTICS

The database has the following characteristics:

- it is structured to provide a comprehensive framework for the storage and retrieval of uniform soil and terrain data that can be used for a wide range of applications at different scales,
- it will contain sufficient data to allow information extraction at a resolution of 1:1 million, both in the form of maps and tables,
- it will be compatible with global databases of other environmental resources,

- it will be amenable to periodic updating and purging of obsolete and/or irrelevant data, and
- be accessible to a broad array of international, regional and national environmental specialists through the provision of standardized resource maps, interpretative maps and tabular information essential for the development, management and conservation of environmental resources.

## **PROCEDURES**

The database is supported by a Procedures Manual which translates SOTER's overall objectives into a workable set of arrangements for the selection, standardization, coding and storing of soil and terrain data.

SOTER requires soils from all corners of the world to be characterised under a single set of rules. As the FAO-Unesco (1974-1981) Soil Map of the World was designed for this purpose, SOTER has adopted the recently Revised Legend (FAO, 1988) as the main tool for differentiating and characterizing its soil components. As there is no universally accepted system for world-wide classification of terrain, SOTER has designed its own system, presented in Chapter 6 of this Manual, which is partly based on earlier FAO work.

The input of soil and terrain data into the SOTER database is contingent upon the availability of sufficiently detailed information. Although some additional information gathering may be required when preparing existing data for acceptance by the database, the SOTER approach is not intended to replace traditional soil surveys. Hence this manual cannot be used as guidelines for soil survey procedures or any other methodology for the collection of field data. Nor does it present a methodology for the interpretation of remotely sensed data. Several handbooks on these techniques are available and details of land resource survey methodology should be contained within them.

## Chapter 2

# Mapping approach and database construction

Within the context of the general objectives of SOTER, as defined in chapter 1, the following subjects will be treated in more detail:


- the procedure for delineating areas with a homogeneous set of soil and terrain characteristics,
- the construction of an attribute database related to the mapping units and based on well-defined differentiating criteria,
- the development of a methodology that should be transferable to and useable by developing countries for national database development at the same or at a larger scale (technology transfer).

### **SOTER MAPPING APPROACH**

The methodology of mapping of land characteristics outlined in this manual originated from the idea that land (in which terrain and soil occur) incorporates processes and systems of interrelationships between physical, biological and social phenomena evolving through time. This idea was developed initially in Russia and Germany (landscape science) and became gradually accepted throughout the world. A similar integrated concept of land was used in the land systems approach developed in Australia by Christian and Stewart (1953) and evolved further by Cochrane *et al.* (1981, 1985), McDonald *et al.* (1990) and Gunn *et al.* (1990). SOTER has continued this development by viewing land as being made up of natural entities consisting of combinations of terrain and soil individuals.

Underlying the SOTER methodology is the identification of areas of land with a distinctive, often repetitive, pattern of landform, lithology, surface form, slope, parent material, and soil. Tracts of land distinguished in this manner are named SOTER units. Each SOTER unit thus represents one unique combination of terrain and soil characteristics. Figure 1 shows the representation of a SOTER unit in the database and gives an example of a SOTER map, with polygons that have been mapped at various levels of differentiation.

**FIGURE 1**  
Relations between a SOTER Unit and their composing parts and major separating criteria



**Example (see figure 1)**

The map shown in figure 1 could have the following legend:

*SOTER description unit*

- 317 one terrain type with one terrain component and one soil component
- 318 one terrain type consisting of an association of two terrain components each having a particular soil component
- 319 one terrain type, consisting of an association of two terrain components, the first having one soil component and the second having an association of two soil components
- 320 one terrain type, consisting of an association of three terrain components, the first having one soil component, the second having an association of three soil components and the third having one soil component
- 321 one terrain type with one terrain component having an association of two soil components (occurs as two polygons)
- 322 one terrain type, consisting of an association of two terrain components each with a soil component

The SOTER mapping approach in many respects resembles physiographic soil mapping. Its main difference lies in the stronger emphasis SOTER puts on the terrain-soil relationship as compared to what is commonly done in traditional soil mapping. This will be true particularly at smaller mapping scales. At the same time SOTER adheres to rigorous data entry formats necessary for the construction of an universal terrain and soil database. As a result of this approach the data accepted by the database will be standardized and will have the highest achievable degree of reliability.

The methodology presented in this manual has been developed for applications at scale of 1:1 million and has been tested successfully in pilot areas in North and South America.



Nevertheless, the methodology also is intended for use at larger scales connected with the development of national soil and terrain databases. A first testing of such a detailed database was carried out in São Paulo State of Brazil at a scale of 1:100 000 (Oliviera and van den Berg, 1992). The SOTER methodology also lends itself well to the production of maps and associated tables at scales smaller than 1:1 million.

Attributes of terrain, soil and other units as used by SOTER are hierarchically structured to facilitate the use of the procedures at scales other than the reference scale of 1:1 million.

### **SOTER SOURCE MATERIAL**

Basic data sources for the construction of SOTER units are topographic, geomorphological, geological and soil maps at a scale of 1:1 million or larger (mostly exploratory and reconnaissance maps). In principle all soil maps that are accompanied by sufficient analytical data for soil characterization according to the revised FAO-Unesco Soil Map of the World Legend (FAO, 1988) can be used for mapping according to the SOTER approach. Seldom, however, will an existing map and accompanying report contain all the required soil and terrain data. Larger scale (semi-detailed and detailed) soil and terrain maps are only suitable if they cover sufficiently large areas. In practice such information will be mostly used to support source material at smaller scales.

As SOTER map sheets will cover large areas, often they will include more than one country, and correlation of soil and terrain units may be required. Where no maps of sufficient detail exist for a certain study area, or where there are gaps in the available data, it may still be possible to extract information from smaller scale maps (e.g. the FAO-Unesco Soil Map of the World at 1:5 million scale or similar national maps), provided that some additional fieldwork is carried out, where necessary in conjunction with the use of satellite imagery. Hence there will often be a need for additional field checks, sometimes supported by satellite imagery interpretation and extra analytical work to complement the existing soil and terrain information. This should be carried out, however, within the context of complementing, updating or correlating existing surveys. It must be stressed that SOTER specifically excludes the undertaking of new land resource surveys within its programme.

Where it is necessary to include an area in the SOTER database for which there is insufficient readily available information, then it is recommended that a survey be carried out according to national soil survey standards, while at the same time ensuring that all parameters required by the SOTER database but not already part of the data being collected. This will ease the subsequent conversion from the national data format into the SOTER data format.

SOTER uses the 1:1 million Operational Navigation Charts and its digital version, the Digital Chart of the World (DMA, 1992), for its base maps. Although it aims at eventual world-wide coverage, the SOTER approach does not envisage a systematic mapping programme, and hence does not prescribe a standard block size for incorporation in the database. Nevertheless, SOTER does recommend that at its reference scale of 1:1 million a block should cover a substantial area (e.g. 100 000 km<sup>2</sup>).

**ASSOCIATED AND MISCELLANEOUS DATA**

SOTER is a land resource database. For many of its applications SOTER data can only be used in conjunction with data on other land-related characteristics but SOTER does not aspire to be able to provide all these data. Nevertheless to obtain a broad characterisation of tracts of land in terms of these complementary characteristics, the SOTER database does include files on climate, vegetation and land use. The former file is in the form of point data, that can be linked to SOTER units through GIS software. Vegetation and land use information is, on the other hand, provided at the level of SOTER units. However, it should be stressed that for specific applications, information on these characteristics should be obtained from specialized databases such as a climatic database. This also applies to natural resource data (e.g. groundwater hydrology) and socio-economic data (e.g. farming systems) which do not form part of the SOTER database.

Miscellaneous data refers to background information that is not directly associated with land resources. SOTER stores information on map source material, laboratory methods, and soil databases from which profile information has been extracted.

## Chapter 3

# SOTER differentiating criteria

The major differentiating criteria are applied in a step-by-step manner, each step leading to a closer identification of the land area under consideration. In this way a SOTER unit can be defined progressively into terrain, terrain component and soil component. Successively an area can thus be characterized by its terrain, its consisting terrain components and their soil components.

The level of disaggregation at each step in the analysis of the land depends on the level of detail or resolution required and the information available. The reference scale of SOTER being 1:1 million, this Manual provides the necessary detail to allow mapping at that scale.

### **TERRAIN**

#### **Physiography**





Physiography is the first differentiating criterion to be used in the characterisation of SOTER units. The term physiography is used in this context as the description of the landforms of the earth's surface. It can best be described as identifying and quantifying as far as possible the major landforms, based on the dominant gradient of their slopes and their relief intensity (see Chapter 6). In combination with a hypsometric (absolute elevation above sea-level) grouping, and a factor characterizing the degree of dissection, a broad subdivision of an area can be made and delineated on the map (see Figure 2), referred to as first and second level major landform in Table 2 of chapter 6. In this way three major landforms can be distinguished in Figure 2.

#### **Parent material**

Areas corresponding to major or regional landforms can be subdivided according to lithology or parent material (see Chapter 6). This will lead to a further definition of the physiographic units by the second differentiating criterion: lithology. The result is shown in Figure 3.

Terrain, in the SOTER context, is thus defined as a particular combination of landform and lithology which characterizes an area. It also possesses one or more typical combinations of surface form, mesorelief, parent material aspect and soil. These form the rationale for a further subdivision of the terrain into terrain components and soil components.

There is no limit to the number of subdivisions that can be applied to the terrain (and terrain components). It is, however, expected that in most cases a maximum of 3 or 4 terrain components and 3 soil components will be sufficient to adequately describe the terrain.



**TERRAIN COMPONENTS**

**Surface form, slope, etc.**

The second step in the subdivision is the identification of areas, within each terrain, with a particular (pattern of) surface form, slope, mesorelief and, in areas covered by unconsolidated material, texture of parent material. This will result in a further partitioning of the terrain into terrain components as is shown in Figures 4 and 5.

It should be noted that at this level of separation it is not always possible at a scale of 1:1 million to map terrain components individually, because of to the complexity of their occurrence. In such cases the information related to non-mappable terrain components is stored in the attribute database only, and no entry is made into the geometric database.


## SOIL COMPONENTS

The final step in the differentiation of the terrain is the identification of soil components within the terrain components. As with terrain components, soil components can be mappable or non-mappable at the considered scale. In the case of mappable soil components, each soil component represents a single soil within a SOTER unit (see Figure 6). However, at a scale of 1:1 million it often will be difficult to separate soils spatially, and a terrain component is likely to comprise a number of non-mappable soil components. In traditional soil mapping procedures such a cluster is known as a soil association or soil complex (two or more soils which, at the scale of mapping, cannot be separated). Non-mappable terrain components (of which there must be at least two in a SOTER unit) are by definition associated with non-mappable soil components. Nevertheless, in the attribute database each non-mappable terrain component can be linked to one or more specific (but non-mappable) soil components. Non-mappable soil components, as in the case of the non-mappable terrain components, do not figure in the geometric database.

### Differences in classification

As the SOTER soil components are characterized according to the FAO-Unesco Soil Map of the World Legend, so the criteria used for separating soil components within each terrain component are based on FAO diagnostic horizons and properties. At the SOTER reference scale of 1:1 million, soils must, in general, be characterized up to the 3rd (i.e. subunit) level following the guidelines provided for this in the annex to the Revised Legend (FAO, 1988).

For soils classified according to Soil Taxonomy (Soil Survey Staff, 1975, 1990 and 1992), the FAO sub-unit level corresponds roughly to the subgroup level. As many of the diagnostic horizons and properties as used by Soil Taxonomy are similar to those employed by FAO, generally there will not be many problems at this level of classification in translating Soil Taxonomy units into FAO units. A major difference between the two systems is the use in Soil Taxonomy of soil temperature and soil moisture regimes, particularly at suborder level. Since these characteristics do not feature in the FAO classification, and SOTER being basically a land resource database, intends to keep climatic data (including those related to soil climate) separated from land and soil data, a more drastic conversion will be required of Soil Taxonomy units which are defined in terms of soil temperature and soil moisture



characteristics. Nevertheless, experience has shown that even in these cases conversion from Soil Taxonomy great groups to FAO sub-units usually will not necessitate major adjustments of to the boundaries of soil mapping units.

### **Differences in use**

In addition to diagnostic horizons and properties, soil components can also be separated according to other factors, closely linked to soils, that have a potentially restricting influence on land use or may affect land degradation. These criteria, several of which are listed by FAO as phases, can include both soil (sub-surface) and terrain (surface, e.g. micro-relief) factors.

### **Soil profiles**

For every soil component at least one, but preferably more, fully described and analyzed reference profiles should be available from existing soil information sources. Following judicious selection, one of these reference profiles will be designated as the representative profile for the soil component. The data from this representative profile must be entered into the SOTER database in accordance with the format as indicated in sections *Profile* and *Horizon data* in Chapter 6 of this Manual. This format is largely based upon the FAO Guidelines for Soil Description (FAO, 1990), which means that profiles described according to FAO or to the Soil Survey Manual (Soil Survey Staff, 1951), from which FAO has derived many of its criteria, can be entered with little or no reformatting being necessary. Compatibility between the FAO-ISRIC Soil Database (FAO, 1989) and the relevant parts of the SOTER database also will facilitate transfer of data already stored in databases set up according to FAO-ISRIC standards.

### **Horizons**

It is recommended that for SOTER the number of horizons per profile is restricted to a maximum of five subjacent horizons, reaching a depth of at least 150 cm where possible. Except for general information on the profile, including landscape position and drainage, each horizon has to be fully characterised in the database by two sets of attributes based on chemical and physical properties. The first set consists of single value data that belong to the representative profile. The second set holds the maximum and minimum values of each numeric attribute, derived from all available reference profiles. In case there is only one reference profile for a soil component then it will obviously not be possible to complete these additional tables.

### **Optional and mandatory data**

Both sets of horizon data consist of mandatory and optional data. Where mandatory data are missing, the SOTER database will accept expert estimates for such values. They will be flagged as such in the database. Optional data should only be entered where the information on them is reliable. For the representative profile these must be measured data.

As with terrain components, the percentage cover of the soil component within the terrain component is indicated. The relative position and relationship of soil components vis-à-vis each other within a terrain component is recorded in the database as well.

## **SOTER UNIT MAPPABILITY**

### **SOTER units in the database and on the map**

At the reference scale of 1:1 000 000 a SOTER unit is composed of a unique combination and pattern of terrain, terrain component and soil component. A SOTER unit is labelled by a SOTER unit identification code that allows retrieval from the database of all terrain, terrain component and soil component data, either in combination or separately. The inclusion of the three levels of differentiation in the attribute database does not imply that all components of a SOTER unit can be represented on a map, as the size of individual components, or the intricacy of their occurrence, may preclude cartographic presentation. The areas shown on a SOTER map can thus correspond to any of the three levels of differentiation of a SOTER unit: terrain, terrain components or soil components. The components not mapped are known to exist, and their attributes are included in the database, although their exact location and extent cannot be displayed on a 1:1 million map.

### **Differences**

In an ideal situation, at least from the point of view of geo-referencing the data, a SOTER unit on the map would be similar to a soil component in the database, i.e. the soil component of the SOTER unit could be delineated on a map. However, at the SOTER reference scale of 1:1 million it is unlikely that many SOTER units can be distinguished on the map at soil component level. This would only be possible if the landscape is relatively uncomplicated. A more common situation at this scale would be for a SOTER unit to consist of terrain with non-mappable terrain components linked to an assemblage of non-mappable soil components (a terrain component association) or, alternatively, a SOTER unit with mappable terrain components that contain several non-mappable soil components (a similar situation as with a soil association on a traditional soil map).

Thus, while in the attribute database a SOTER unit will hold information on all levels of differentiation, a SOTER map will display units whose content varies according to the mappability of the SOTER unit components. The disadvantage of not being able to accurately locate terrain components and/or soil components is therefore only relevant when data of complex terrains are being presented in map format. It does not affect the capability of the SOTER database to generate full tabular information on terrain, terrain component and soil component attributes while at the same time indicating the spatial relationship between and within these levels of differentiation.

## **SOTER APPROACH AT OTHER SCALES**

### **Smaller scales**

The methodology presented in this manual has been developed for applications at a scale of 1:1 million, which is the smallest scale still suitable for land resource assessment and monitoring at national level. However, as potentially the most complete universal terrain and soil database, SOTER is also suited to provide the necessary information for the compilation of smaller scale continental and global land resource maps and associated data tables. The methodology was tested by FAO for the compilation of the physiographic base for a future update of the Soil Map of the World (Eschweiler, 1993 and Wen, 1993).

Flexibility to cater for a wide range of scales is achieved through adopting a hierarchical structure for various major attributes, in particular those that are being used as differentiating criteria (landform, lithology, surface form, etc.). Examples of such hierarchies are given in this Manual for land use and vegetation (see Chapter 7). Different levels of these hierarchies can be related to particular scales. A hierarchy for the soil component can be derived from the FAO-Unesco Soil Map of the World Legend, with the level of soil groupings being related to extremely small scale maps, as exemplified by the map of world soil resources at 1:25 million (FAO, 1991). Soil units (2nd level) can be used for 1:5 million world soil inventory maps, while the soil subunits are most suitable for 1:1 million mapping. The density per unit area of point observations will vary according to the scale employed, with larger scales requiring a more compact ground network of representative profiles, as soils are being characterized in more detail.

A simplification of the database can be applied at scales substantially smaller than the reference scale of 1:1 million, but only the most elementary soil physical and chemical data are relevant if the scale is smaller than 1:10 million. It is thus necessary to realize that the SOTER database discussed in this Manual is meant for a scale of 1:1 million only, and that expansion or contraction of the data set will be necessary when changing the resolution of the SOTER database.

### **Larger scales**

As a systematic and highly organized way of mapping and recording terrain and soil data, the SOTER methodology can easily be extended to include reconnaissance level inventories, i.e. at a scale between 1:1 million and 1:100 000 (e.g. Oliveira and van den Berg, 1992).

Adjustments to the content of the attribute data set are necessary if SOTER maps at scales other than 1:1 million are being compiled. With an increase in resolution, the highest level constituents of a SOTER unit, i.e. the terrain, will gradually lose importance, and may disappear altogether at a scale of 1:100 000. This is because in absolute terms the area being mapped is becoming smaller, and terrain alone may not continue to offer sufficient differentiating power. Conversely, the lower part of the SOTER unit will gain in importance with more detailed mapping. At larger scales SOTER units will thus become delineations of soil entities, with the information on terrain becoming incorporated in the soil attributes. Hence scale increases require more detailed information on soils for most practical applications. Additional attributes which might be included could be soil micronutrient content, composition of organic fraction, detailed slope information, etc.



## Chapter 4

# SOTER database structure

In every discipline engaged in mapping of spatial phenomena, two types of data can be distinguished:

- geometric data, i.e. the location and extent of an object represented by a point, line or surface, and topology (shapes, neighbours and hierarchy of delineations),
- attribute data, i.e. characteristics of the object.

These two types of data are present in the SOTER database. Soils and terrain information consist of a geometric component, which indicates the location and topology of SOTER units, and of an attribute part that describes the non-spatial SOTER unit characteristics. The geometry is stored in that part of the database that is handled by Geographic Information System (GIS) software, while the attribute data is stored in a separate set of attribute files, manipulated by a Relational Database Management System (RDBMS). A unique label attached to both the geometric and attribute database connects these two types of information for each SOTER unit (see Figure 7, in which part of a map has been visualized in a block diagram).


The overall system (GIS plus RDBMS) stores and handles both the geometric and attribute database. This manual limits itself to the attribute part of the database only, in particular through elaborating on its structure and by providing the definitions of the attributes (Chapter 6). A full database structure definition is given by Tempel (1994b).

A relational database is one of the most effective and flexible tools for storing and managing non-spatial attributes in the SOTER database (Pulles, 1988). Under such a system the data is stored in tables, whose records are related to each other through the specific identification fields (primary keys), such as the SOTER unit identification code. These codes are essential as they form the links between the various subsections of the database, e.g. the terrain table, the terrain component and the soil component tables. Another characteristic of the relational database is that when two or more components are similar, their attribute data need only to be entered once. Figure 8 gives a schematic representation of the structure of the attribute database. The blocks represent tables in the SOTER database and the solid lines between the blocks indicate the links between the tables.


### GEOMETRIC DATABASE

The geometric database contains information on the delineations of the SOTER unit. It also holds the base map data (cultural features such as roads and towns, the hydrological network and administrative boundaries). In order to enhance the usefulness of the

**FIGURE 7**  
SOTER units, their terrain components (tc), attributes, and location



**FIGURE 8**  
SOTER attribute database structure with area and point data (1:M = one to many, M:1 = many to one relations)



database, it will be possible to include additional overlays for boundaries outside the SOTER unit mosaic. Examples of such overlays could be socio-economic areas (population densities), hydrological units (watersheds) or other natural resource patterns (vegetation, agro-ecological zones).

### ATTRIBUTE DATABASE

The attribute database consists of sets of files for use in a Relational DataBase Management System (RDBMS). The attributes of the terrain and terrain component are either directly available or can be derived from other parameters during the compilation of the database. Only for horizon data, two types of attributes can be distinguished, depending on their importance and availability: mandatory attributes and optional attributes

Many of the horizon parameters of the soil component consist of measured characteristics of which the availability varies considerably. However, there is a minimum set of soil attributes that are generally needed if any realistic interpretation of the soil component of a SOTER unit is to be expected. Therefore their presence is considered mandatory. Other soil horizon attributes are of lesser importance and their presence in the database is considered optional. Whether a horizon attribute is mandatory or optional is indicated in the chapter describing the attributes. It is imperative that, in order to preserve the integrity of the SOTER database, a complete list of mandatory attributes is entered for each soil component. Optional attributes are accepted by the database as and when available.

Each of the attributes can be divided into descriptive (e.g. landform) and numerical (e.g. pH, slope gradient) data.

Under the SOTER system of labelling (see *SOTER unit codes* in Chapter 5 for a detailed description of the labelling conventions) all SOTER units are given a unique identification code, consisting of 4 digits. In the terrain component and soil component tables this identification code is completed with subcodes for terrain component and soil component number.

Where identical terrain components and soil components occur in several SOTER units in different proportions, a separation between the tables holding the data on proportion/position of the terrain component and soil component (terrain component block and soil component block) and the tables holding the data of the terrain component and soil component (terrain component data block and profile and horizon blocks) is made (see Figure 8).

Thus, the terrain component information is split into two tables:

- the *terrain component table* which indicates the SOTER unit to which the terrain component belongs and the proportion that it occupies within that unit
- the *terrain component data table* which holds all specific attribute data for the terrain component

In the first table there is space for an entry for each individual terrain component within a SOTER unit, while in the second table only entries are made for data of these terrain components if they possess a not previously occurring set of attribute values.

TABLE 1  
Non-spatial attributes of a SOTER unit

<b>TERRAIN</b>		
1 SOTER unit_ID	6 slope gradient	11 dissection
2 year of data collection	7 relief intensity	12 general lithology
3 map_ID	8 major landform	13 permanent water surface
4 minimum elevation	9 regional slope	
5 maximum elevation	10 hypsometry	
<b>TERRAIN COMPONENT</b>		<b>TERRAIN COMPONENT DATA</b>
14 SOTER unit_ID	18 terrain component data_ID	26 texture group non-consolidated parent material
15 terrain component number	19 dominant slope	27 depth to bedrock
16 proportion of SOTER unit	20 length of slope	28 surface drainage
17 terrain component data_ID	21 form of slope	29 depth to groundwater
	22 local surface form	30 frequency of flooding
	23 average height	31 duration of flooding
	24 coverage	32 start of flooding
	25 surface lithology	
<b>SOIL COMPONENT</b>		<b>HORIZON (* = mandatory)</b>
33 SOTER unit_ID	63 profile_ID*	96 soluble K <sup>+</sup>
34 terrain component number	64 horizon number*	97 soluble Cl <sup>-</sup>
35 soil component number	65 diagnostic horizon*	98 soluble SO <sub>4</sub> <sup>-</sup>
36 proportion of SOTER unit	66 diagnostic property*	99 soluble HCO <sub>3</sub> <sup>-</sup>
37 profile_ID	67 horizon designation	100 soluble CO <sub>3</sub> <sup>-</sup>
38 number of reference profiles	68 lower depth*	101 exchangeable Ca <sup>++</sup>
39 position in terrain component	69 distinctness of transition	102 exchangeable Mg <sup>++</sup>
40 surface rockiness	70 moist colour*	103 exchangeable Na <sup>+</sup>
41 surface stoniness	71 dry colour	104 exchangeable K <sup>+</sup>
42 types of erosion/deposition	72 grade of structure	105 exchangeable Al <sup>+++</sup>
43 area affected	73 size of structure elements	106 exchangeable acidity
44 degree of erosion	74 type of structure*	107 CEC soil*
45 sensitivity to capping	75 abundance of coarse fragments*	108 total carbonate equivalent
46 rootable depth	76 size of coarse fragments	109 gypsum
47 relation with other soil components	77 very coarse sand	110 total carbon*
	78 coarse sand	111 total nitrogen
	79 medium sand	112 P <sub>2</sub> O <sub>5</sub>
	80 fine sand	113 phosphate retention
	81 very fine sand	114 Fe dithionite
	82 total sand*	115 Al dithionite
	83 silt*	116 Fe pyrophosphate
	84 clay*	117 Al pyrophosphate
	85 particle size class	118 clay mineralogy
	86 bulk density*	
	87 moisture content at various tensions	
	88 hydraulic conductivity	
	89 infiltration rate	
	90 pH H <sub>2</sub> O*	
	91 pH KCl	
	92 electrical conductivity	
	93 soluble Na <sup>+</sup>	
	94 soluble Ca <sup>++</sup>	
	95 soluble Mg <sup>++</sup>	
<b>PROFILE</b>		
48 profile_ID		
49 profile database_ID		
50 latitude		
51 longitude		
52 elevation		
53 sampling date		
54 lab_ID		
55 drainage		
56 infiltration rate		
57 surface organic matter		
58 classification FAO		
59 classification version		
60 national classification		
61 Soil Taxonomy		
62 phase		

In the same way the soil component information is stored in three tables:

- the *soil component table* holds the proportion of each soil component within a SOTER unit/terrain component combination and its position within the terrain component;
- the *profile table* holds all attribute data for the soil profile as a whole;
- the *horizon table* holds the data for each individual soil horizon. To be able to give some degree of variability it consists of four sets of attribute values:
  - a. single values taken from the representative profile, either: (1) measured, or (2) estimated (only for mandatory attributes);
  - b. maximum (measured) values taken from all available profiles within the soil component;
  - c. minimum (measured) values taken from all available profiles within the soil component.

For the profile and horizon tables the same conditions for the terrain component data table are valid. Only soil profiles not previously described may be entered. For profile/horizon data describing soils occurring in various soil components only one entry is necessary.

The horizon tables must contain all mandatory measured data: (a1) data set. In case data is not available for some of the quantifiable attributes, SOTER will allow expert estimates to be used for attributes of the representative profile: (a2) data set. Measured and estimated values of the representative profile will thus be stored separately.

To be able to indicate the variability within a soil component various statistical parameters can be determined. Data from the representative profile are considered as modal values. However, considering the small number of profiles generally available for the compilation of the soil component, it is not realistic to aim at standard deviations and means. Therefore only maximum and minimum values of the profiles of the same soil component give an indication of the range of variation that exist within the component. They will be stored respectively in the (b) and (c) data sets.

It is strongly recommended that in conjunction with the SOTER database a national soil profile database be established along the lines of the FAO-ISRIC Soil Database (FAO, 1989), in which, amongst others, all representative profiles would be accommodated.

All mandatory and optional attributes for the soil component, as well as all other non-spatial attributes of the SOTER units, are listed in Table 1. The listing for the soil component attributes is compatible, but contains some additional items, with the data set that is stored in the FAO-ISRIC Soil Database.

The database can be asked to calculate automatically a number of derived parameters from the values entered for the mandatory and optional attributes. These include, amongst others, CEC per 100 g clay, base saturation and textural class.



## Chapter 5

# Additional SOTER conventions

The various conventions described in this chapter form an addition to those characterized in Chapter 2. They mainly concern rules governing the minimum size of a SOTER unit, both in absolute and relative terms, as well as criteria determining the selection of representative profiles, relations with associated databases, type of data and missing data.

SOTER database management procedures, such as date stamps and backup procedures, are not treated in this manual, but are to be described in a separate manual (Tempel, in prep.).

### SOTER UNIT CODES

Each SOTER unit is assigned an identifying code that is unique for the database in question. Tentatively, the SOTER coding will consist of a simple numbering system. This code will normally range from 1 to 999, or 9999 for large maps. The terrain components within each SOTER unit are given single digit extension numbers separated by a slash (/) and ranked according to the size of the component. A similar single digit extension number is used to code the soil components. This means that a maximum of 10 terrain components (first digit with values from 0-9) each with 10 soil components (second digit) can be stored in the database. The component extension numbers are separated from the SOTER unit code by a slash. The identification code of a soil component in the database thus can range from 1/11 to 9999/99. Numbering is not strictly sequential, as the total number of terrain components per SOTER unit and soil components per terrain component is limited (see chapter 5.4), and identification codes like 1/17 (7 soil components within terrain component 1) or 25/53 (3 soil components in terrain component 5) are unlikely to occur.

When individual databases are merged into regional and global databases, then the SOTER identification codes can be preceded by the ISO code for the country. When databases of neighbouring countries are entered into one database, then cross-boundary SOTER units will have different codes in each country. If a GIS is used the SOTER units of one country can automatically be given the code of their counterpart on the other side of the border (assuming that proper correlation has been carried out), otherwise this has to be done manually.

At national level this coding convention is only applicable to 1:1 million maps. For larger scale maps and databases there is no need to follow a unified system.

### **MINIMUM SIZE OF THE SOTER UNIT**

As a general rule of thumb the minimum size of a single SOTER unit is  $0.25 \text{ cm}^2$  on the map which, at a scale of 1:1 million, equals  $25 \text{ km}^2$  in the field. This is the smallest area that can still be cartographically represented. Mostly such tiny units will correspond to narrow elongated features (floodplains, ridges, valleys) or strongly contrasting terrain and soil features. In general, SOTER units will be much larger.

If there are gradual changes in landscape features, new SOTER units can be delineated when any one terrain component or soil component of a unit changes in area by more than 50%.

### **NUMBER OF SOIL AND TERRAIN COMPONENTS**

Within a SOTER unit terrain components and soil components can occupy any percentage of the terrain and terrain component respectively, provided the total area of each component is not less than what is indicated in the previous section. In theory this would allow for an unlimited number of terrain components within each SOTER unit, or soil components within each terrain component. In practice this is unlikely to occur, as many terrain components and soil components cover sizeable areas. SOTER recommends that a minimum area of 15% of the SOTER unit is taken into account when defining terrain and soil components, unless the SOTER unit in question is very large, or it involves strongly contrasting terrain or soil components, when the percentage coverage can be less.

Most commonly it is expected that a SOTER unit would be subdivided into up to 3 or 4 terrain components, each with not more than 3 soil components, resulting in a maximum of 12 subdivisions. Obviously, the proportional areal sum of soil components within each terrain component, and terrain components within each SOTER unit, will always be 100%.

It is advisable that map compilers exercise restraint in subdividing terrain into terrain and soil components. Only those criteria that can be considered important for analyzing a landscape in subsequent interpretations should be selected. Significant changes in attributes such as parent material, surface form and slope gradient, which at the same time should cover substantial areas, qualify as criteria for defining new SOTER units. Terrain components should be split into soil components only if there are clear changes in diagnostic criteria which will reflect in land use or land degradation aspects. Minor changes in any of these criteria should be considered as part of the natural variability that at a scale of 1:1 million can be expected to occur within each SOTER unit. Discretion in defining terrain and soil components is absolutely necessary in order not to generate an excessive number of components and so lengthening the time required for coding, entering and processing of data.

### **REPRESENTATIVE SOIL PROFILES**

The representative profile used to typify a specific soil component is chosen from amongst a number of reference profiles with similar characteristics. Where possible SOTER will rely on a selection of reference profiles made by the original surveyors. It is envisaged that all reference profiles taken into consideration be stored in a national soil profile database, preferably based on the FAO-ISRIC Soil Database format. The SOTER database includes a key to national databases.



The SOTER database also includes a code that shows how many reference profiles were considered for the selection of the representative profile, and were used to determine the maximum and minimum values of attributes as well.

### UPDATING PROCEDURES

SOTER units and their attributes are unique in both space and time, and although soil and in particular terrain characteristics are thought to have a high degree of temporal stability, it might become necessary to update certain attributes from time to time. At present, there is no procedure for updates of the geographic data, such as the boundaries of the SOTER units. However, replacing (parts of) map sheets by more recent maps will involve changes in attribute data as well, for which the guidelines below can be used.

Updating the attribute database could become necessary because of *missing data*, *incorrect data* or *obsolete data* in the database. If there are some data gaps, the voids can be filled when additional data becomes available. Incorrect data, which include data that is being replaced by (a set of) more reliable data (e.g. a representative profile is being substituted by another, more representative profile) can be replaced by new data, although a note has to be made of this in the database. In contrast, obsolete data is not simply replaced by more up-to-date information. Instead, old data is downloaded into a special database containing obsolete data, after which the latest data is entered into the regular database. In this way the database with obsolete data can be used for the monitoring of changes over time. When certain parameters are measured at regular intervals, then periodic updating will become necessary.

The SOTER unit Identification code does indicate to which level of differentiation the SOTER unit can be mapped. The database is capable of generating a number of relational data that are pertinent to each SOTER unit, and between the SOTER units (e.g. percentage of each soil component within terrain component or SOTER unit, total area of all terrain components with identical terrain component data code, etc.).



## Chapter 6

# Attribute coding

Note that the numbers preceding the attributes in Table 1 are identical to the numbers of the attributes in this chapter, written in the left margin. They also figure on the SOTER data entry forms (see Annex 5 for a pro forma).

The SOTER unit identification code, referring to the map unit, is completed in the database by two additional digits, separated from the SOTER unit code by a slash. The first digit represents the terrain component number. The second digit constitutes the soil component number. Eventually, the SOTER unit identification code will be the unique identifier for SOTER units on a worldwide scale (see also *SOTER unit codes* in Chapter 5).

However, for compilers of SOTER data on a national or regional scale it is sufficient to attach locally unique identification codes to each SOTER unit, taking into account the coding conventions explained in the section *SOTER unit codes*. These identification codes will be converted into globally unique identifiers before entry into a continental or worldwide SOTER database.

Class limits as used in this manual are defined as follows. The upper class limit is included in the next class. For example, slope class 2-5% (item 9) includes all slopes from 2.0 to 4.9%. Hence, a slope of 5% would fall in slope class 5-8%.

### TERRAIN

#### 1 *SOTER unit\_ID*

The SOTER unit\_ID is the identification code of a SOTER unit on the map and in the database. It links the mapped area to the attributes in the database and in particular, it identifies which terrain belongs to a SOTER unit. SOTER units which have identical attributes carry the same SOTER unit\_ID. In other words the SOTER unit\_ID is similar to a code for a mapping unit on a conventional soil map.

For each SOTER map, a unique code (up to 4 digits) is assigned to every SOTER unit that has been distinguished. On most SOTER maps 2 or 3 digits will suffice.

#### 2 *year of data collection*

The year in which the original terrain data were collected will serve as the time stamp for each SOTER unit. Where the SOTER unit has been composed on the basis of several sources of information, it is advisable to use the major source for dating it. In this manner a link between the SOTER unit and the major source of information, which

TABLE 2  
Hierarchy of major landforms

1st level	2nd level	gradient (%)	relief intensity
<b>L</b> level land	<b>LP</b> plain	<8	<100m/km
	<b>LL</b> plateau	<8	<100m/km
	<b>LD</b> depression	<8	<100m/km
	<b>LF</b> low-gradient footslope	<8	<100m/km
	<b>LV</b> valley floor	<8	<100m/km
<b>S</b> sloping land	<b>SM</b> medium-gradient mountain	15-30	>600m/2km
	<b>SH</b> medium-gradient hill	8-30	>50m/slope unit
	<b>SE</b> medium-gradient escarpment zone	15-30	<600m/2km
	<b>SR</b> ridges	8-30	>50m/slope unit
	<b>SU</b> mountainous highland	8-30	>600m/2km
	<b>SP</b> dissected plain	8-30	<50m/slope unit
<b>T</b> steep land	<b>TM</b> high-gradient mountain	>30	>600m/2km
	<b>TH</b> high-gradient hill	>30	<600m/2km
	<b>TE</b> high-gradient escarpment zone	>30	>600m/2km
	<b>TV</b> high gradient valleys	>30	var.
<b>C</b> land with composite landforms	<b>CV</b> valley	>8	var.
	<b>CL</b> narrow plateau	>8	var.
	<b>CD</b> major depression	>8	var.

Notes: var. = variable.

should be listed under *map\_ID*, can easily be made. The year of compiling the data according to the SOTER procedures is thus not recorded, unless the compilation itself has resulted in some major reinterpretation based on additional sources of information, like fresh satellite imagery. In general the year of compilation can be deduced from the year in which the data was entered into the database, as both years are likely to be the same or very close to each other. It is assumed that the year in which the terrain data were collected also applies to the terrain component data, and no separate date entry is required for this.

### 3 *map\_ID*

The source map identification code from which the data were derived for the compilation of the SOTER units. There is room for 12 characters.

### 4 *minimum elevation*

Absolute minimum elevation of the SOTER unit, in metres above sea level. Both the minimum and maximum elevation can be read from a contoured topographic map.

### 5 *maximum elevation*

Absolute maximum elevation of the SOTER unit, in metres above sea level.

### 6 *slope gradient*

The dominant slope angle, expressed as a percentage, prevailing in the terrain.

## 7 relief intensity

The relief intensity is the median difference between the highest and lowest point within the terrain per specified distance. This specified distance can be variable, but is expressed in m/km in the database.

## 8 major landform

Landforms are described foremost by their morphology and not by their genetic origin, or processes responsible for their shape. The dominant slope is the most important differentiating criterion, followed by the relief intensity. The relief intensity is normally given in metres per kilometre, but for distinction between hills and mountains it is practical to use two kilometre intervals (see Table 2).

At the highest level of landform separation, suitable for scales equal to or smaller than 1:10 million, four groups are being distinguished (adapted from Remmelzwaal, 1991). They can be subdivided when the position of the landform vis-a-vis the surrounding land is taken into consideration.

Where not clear from the gradient or relief intensity, the distinction between the various second level landforms follows from the description in Annex 1.

### REGIONAL LANDFORMS

Major landforms can be further characterized according to three criteria. These are:

1. regional slope
2. hypsometry
3. dissection

The differentiating power of these criteria is highest with respect to level lands, although they can be used for sloping lands with a relief intensity of less than 600 m/2 km as well. For steep lands with a high relief intensity they have little utility, with the exception of the hypsometric level.

## 9 regional slope

A refining of slope classes compared to those used for major landforms is possible. The dominant slopes can be broken down into the following classes:

### a) Simple landforms

<b>W</b>	0- 2 %	flat, wet*
<b>F</b>	0- 2 %	flat
<b>G</b>	2- 5 %	gently undulating
<b>U</b>	5- 8 %	undulating
<b>R</b>	8-15 %	rolling
<b>S</b>	15-30 %	moderately steep

<b>T</b>	30-60 %	steep
<b>V</b>	≥ 60 %	very steep

\* wet is defined as < 90% permanent water surface > 50% (see also item 12)

b) Complex landforms\*\*

<b>CU</b>	Cuesta-shaped
<b>DO</b>	Dome-shaped
<b>RI</b>	Ridged
<b>TE</b>	Terraced
<b>IN</b>	Inselberg covered (occupying at least 1% of level land)
<b>DU</b>	Dune-shaped
<b>IM</b>	With intermontane plains (occupying at least 15%)
<b>WE</b>	With wetlands (occupying at least 15%)
<b>KA</b>	Strong karst

\*\* in the case of complex landforms, the protruding landform should be at least 25 m high (if not it is to be considered mesorelief) except for terraced land, where the main terraces should have elevation differences of at least 10 m.

These subdivisions are mainly applicable to level landforms, and to some extent to sloping landforms. They are not to be used for steep lands, except in the case of mountains with intermontane plains, but may be used for lands with complex landforms, where the subdivision can be related to the constituent landform with the lesser slope.

## 10 *hypsometry*

The hypsometric level is, for level and slightly sloping land (relief intensity of less than 50 m) an indication of the height above sea level of the local base level. For lands with a relief intensity of more than 50 m the hypsometric is used to indicate the height above the local base (i.e. local relief).

a) Level lands and sloping lands (relief intensity < 50 m/slope unit)

<b>1</b>	< 300 m	very low level (plain etc.)
<b>2</b>	300- 600 m	low level
<b>3</b>	600-1500 m	medium level
<b>4</b>	1500-3000 m	high level
<b>5</b>	≥ 3000 m	very high level

b) Sloping lands (relief intensity > 50 m/slope unit)

<b>6</b>	< 200 m	low (hills etc.)
<b>7</b>	200-400 m	medium
<b>8</b>	≥ 400 m	high

c) Steep and sloping lands (relief intensity > 600 m/2 km)

<b>9</b>	600-1500 m	low (mountains etc.)
<b>10</b>	1500-3000 m	medium

- |           |             |           |
|-----------|-------------|-----------|
| <b>11</b> | 3000-5000 m | high      |
| <b>12</b> | 5000 m      | very high |

### **11** *dissection*

The degree of dissection is difficult to quantify in a practical manner. Factors like coverage, slope and depth of dissected features all contribute to the intensity of landscape dissection. SOTER uses the drainage density as a qualitative measure of the degree of dissection. The higher the drainage density, the more dissected a tract of land is, and in general also the steeper the slopes of the dissected parts will be. The depth of dissection can be assumed to increase with an increased density of the drainage network and steeper landscape slopes. Conversely, a high drainage density on very flat land (dominant slopes < 2%) is not necessarily related to the dissection of the terrain, but could be an indication of the wetness of the land.

The most accurate way to measure the drainage density (defined as the average length of drainage channels per unit area of land, expressed as  $\text{km km}^{-2}$ ) is to actually measure the length of all well-defined, permanent and seasonal, streams and rivers within a representative block. This should be done on good quality 1:50,000 or larger maps. Techniques exist to speed up this measurement through intersection point counting (Verhasselt, 1961).

In practice the necessary material to carry out this measurement is often not available, and only quantitative estimates can be made. This should be done with aid of the most detailed material available (maps, aerial photos or satellite images). Only three classes are being distinguished:

- |          |                               |                    |
|----------|-------------------------------|--------------------|
| <b>1</b> | < 10 $\text{km km}^{-2}$      | slightly dissected |
| <b>2</b> | 10-25 $\text{km km}^{-2}$     | dissected          |
| <b>3</b> | $\geq 25$ $\text{km km}^{-2}$ | strongly dissected |


Figure 9 provides an illustration, at a scale of 1:50,000, of two of these classes. The degree of dissection is not applicable to land with a relief intensity of more than 600 m.

### **12** *general lithology*

For each SOTER unit a generalized description of the consolidated or unconsolidated surficial material, underlying the larger part of the terrain, is given. Major differentiating criteria are petrology and mineralogical composition (Holmes, 1968, Strahler, 1969). At the 1:1 million scale the lithology should at least be specified down to group level. Codes are shown in Table 3.

### **13** *permanent water surface*

Indicate the percentage of the SOTER unit that is largely (i.e. > 90%, thus excluding small islands etc.) permanently (i.e. more than 10 month/year) covered by water. Bodies of water large enough to be delineated on the map are not considered part of a SOTER unit.



**TABLE 3**  
Hierarchy of lithology

major class	group	type
I igneous rock	IA acid igneous	IA1 granite
		IA2 grano-diorite
		IA3 quartz-diorite
		IA4 rhyolite
..	II intermediate igneous	II1 andesite, trachyte, phonolite
..		II2 diorite-syenite
..	IB basic igneous	IB1 gabbro
..		IB2 basalt
..		IB3 dolerite
..	IU ultrabasic igneous	IU1 peridotite
..		IU2 pyroxenite
..		IU3 ilmenite, magnetite, ironstone, serpentine
M metamorphic rock	MA acid metamorphic	MA1 quartzite
		MA2 gneiss, migmatite
		MA3 slate, phyllite (pelitic rocks)
		MA4 schist
..	MB basic metamorphic	MB1 slate, phyllite (pelitic rocks)
..		MB2 schist
..		MB3 gneiss rich in ferro-magnesian minerals
..		MB4 metamorphic limestone (marble)
S sedimentary rock	SC clastic sediments	SC1 conglomerate, breccia
		SC2 sandstone, greywacke, arkose
		SC3 siltstone, mudstone, claystone
		SC4 shale
		SC5 ironstone
..	SO organic	SO1 limestone, other carbonate rocks
..		SO2 marl and other mixtures
..		SO3 coals, bitumen & related rocks
..	SE evaporites	SE1 anhydrite, gypsum
..		SE2 halite
U unconsolidated	UF fluvial	
	UL lacustrine	
	UM marine	
	UC colluvial	
	UE eolian	
	UG glacial	
	UP pyroclastic	
	UO organic	



**TERRAIN COMPONENT**

This section includes attributes to identify any terrain component, its percentage within the SOTER unit (15-100%) and a link to the complete set of attribute data of a terrain component (see *Terrain component data*).

**14 SOTER unit\_ID**

See SOTER unit\_ID under paragraph *Terrain*.

**15 terrain component number**

The sequence number of the terrain component in the terrain. The largest terrain component in the SOTER unit comes first, followed by the second in size, and so on. The combination SOTER unit\_ID and terrain component number (e.g. 2034/1) gives the complete identification code for each terrain component within the database.

**16 proportion of SOTER unit**

The proportion that the terrain component occupies within the SOTER unit. As stated in section Number of soil and terrain components in Chapter 5, a terrain component normally covers not less than 15% of a the terrain. The sum of all terrain components should be 100%.

<b>Examples</b>	SOTER unit_id = 2034,	SOTER unit_id = 2034
	terrain component number = 1	terrain component number = 2
30%	proportion within SU = 70%	proportion within SU =

**17 terrain component data\_ID**

If two (or more) terrain components are completely similar, then their data will only be entered once in the database. The data code has the format SOTER unit\_ID/terrain component number. When referring to an already described terrain component data\_ID the first terrain component with a particular attribute content will also be used for subsequent identical terrain components. In case a terrain component has not been described before in the database, then its code will also be used as its data code (four plus one digits).

<b>Examples</b>	
case A (two terrain components, both not yet described in the attribute database)	
SOTER unit_ID = 2034,	SOTER unit_ID = 2034
terrain component number = 1	terrain component number = 2
proportion within SU = 70%	proportion within SU = 30%
terrain component data_ID= 2034/1	terrain component data_ID= 2034/2
case B (two terrain components, one already described (marked with *), one not yet)	
SOTER unit_ID = 2035	SOTER unit_ID = 2035
terrain component number = 1	terrain component number = 2
proportion within SU = 60%	proportion within SU = 40%
terrain component data_ID= 2034/2*	terrain component data_ID= 2035/2

**TERRAIN COMPONENT DATA****18** *terrain component data\_ID*

See terrain component data\_ID under section *Terrain*.

*SLOPE CHARACTERISTICS*

Items 19-21 characterize the slope of the terrain component.

**19** *dominant slope*

Dominant slope gradient of the terrain component, %.

**20** *length of slope*

Estimated dominant length of slope, m.

**21** *form of slope*

The form of the dominant slope (only entered if the dominant slope gradient > 2%)

**U** Uniform slope.

**C** Concave, lower slope with decreasing gradient downslope.

**V** Convex, upper slope with decreasing gradient upslope.

**I** Irregular slope.

*MESO-RELIEF*

Items 22-24 characterize the meso-relief or local surface forms.

**22** *local surface form*

A number of characteristic meso-relief or local surface forms can be recognised at the 1:1 million scale (Day, 1983; FAO, 1977; Soil Survey Staff, 1951), in addition to the slope form as listed below (this list is not exhaustive).

**H** hummocky      very complex pattern of slopes extending from somewhat rounded depressions or kettleholes of various sizes to irregular conical knolls or knobs. There is a general lack of concordance between knolls or depressions. Slopes ranges are large and vary generally between 4 % and 70 %.

**M** mounded      coverage (at least 5 %) by isolated mounds more than 2.5 m high.

**K** towered      coverage (at least 5 %) by isolated steep sided karst towers more than 2.5 m high.

<b>R</b>	ridged	coverage (at least 5 %) by parallel, sub-parallel or intersecting usually sharp-crested ridges (elongated narrow elevations) more than 2.5 m high.
<b>T</b>	terraced	level areas (less than 2 % slope) bounded on one side by a steep slope more than 2.5 m high with another flat surface above it.
<b>G</b>	gullied	coverage (at least 5 %) by steep-sided gullies more than 2.5 m deep.
<b>S</b>	strongly dissected	areas with a drainage density of more than 25 km km <sup>-2</sup> , the depth dissected of the drainage lines being at least 2.5 m.
<b>D</b>	dissected	areas with a drainage density of more than 10 km km <sup>-2</sup> , the depth of the drainage lines being at least 2.5 m.
<b>L</b>	slightly dissected	areas with a drainage density of less than 10 km km <sup>-2</sup> , the depth of the drainage lines being at least 2.5 m.

### 23 *average height*

The average height of the meso-relief (or depth where applicable) in metres, depth being indicated by a minus sign.

### 24 *coverage*

The estimated percentage coverage of the meso-relief elements within the terrain component

### 25 *lithology surficial material*


Description of the consolidated or unconsolidated surficial materials which underlie most of the terrain component. These include the types of rockmass from which parent material is derived, and other unconsolidated mineral or organic deposits. The same list of parent materials is used as was given for the terrain unit lithology (see Table 3). If the type level of parent material, already indicated at terrain level, does not vary then no further entry has to be made here.

### 26 *texture of non-consolidated parent material*

The texture group of particles <2 mm of the non-consolidated parent material, or the parent material at 2 m if the soil is deeply developed, is given. Figure 10 shows the different groups in a texture triangle.

### 27 *depth to bedrock*

The average depth to consolidated bedrock in metres. For depths less than 2 m the unit is 0.1 m. For depths more than 10 m the depth can be given to the nearest 5 metres.



## 28 surface drainage

Surface drainage of the terrain component (after Cochrane *et al.*, 1985 and Van Waveren *et al.*, 1987).

- |                         |   |
|-------------------------|---|
| <b>E</b> extremely slow | water ponds at the surface, and large parts of the terrain are waterlogged for continuous periods of more than 30 days            |
| <b>S</b> slow           | water drains slowly, but most of the terrain does not remain waterlogged for more than 30 days continuously                       |
| <b>W</b> well           | water drains well but not excessively, nowhere does the terrain remain waterlogged for a continuous period of more than 48 hours  |
| <b>R</b> rapid          | excess water drains rapidly, even during periods of prolonged rainfall  |
| <b>V</b> very rapid     | excess water drains very rapidly, the terrain does not support growth of short rooted plants even if there is sufficient rainfall |

## 29 depth of groundwater

The depth in metres of the mean ground water level over a number of years as experienced in the terrain component.

## **FLOODING**

Flooding is characterized by items 30-32:

### **30** *frequency*

Frequency of the natural flooding of the terrain component in classes after FAO (1990).

<b>N</b>	none
<b>D</b>	daily
<b>W</b>	weekly
<b>M</b>	monthly
<b>A</b>	annually
<b>B</b>	biennially
<b>F</b>	once every 2-5 years
<b>T</b>	once every 5-10 years
<b>R</b>	rare (less than once in every 10 years)
<b>U</b>	unknown

### **31** *duration*

Duration of the flooding of the terrain component in classes after FAO (1990).

<b>1</b>	less than 1 day
<b>2</b>	1-15 days
<b>3</b>	15-30 days
<b>4</b>	30-90 days
<b>5</b>	90-180 days
<b>6</b>	180-360 days
<b>7</b>	continuously

### **32** *start*

Give the month (indicated by a figure) during which flooding of the terrain component normally starts. Three entries are possible.

## **SOIL COMPONENT**

This section includes, besides the SOTER identification codes, all the attributes of the soil component (items 33 to 47). General attributes linked to the representative soil profile and horizon attributes are dealt with in the next sections, *Profile* and *Horizon data*.

### **33** *SOTER unit\_ID*

See SOTER unit\_ID under section *Terrain*. The SOTER unit\_ID given in the terrain chapter should also be used here.

**34** *terrain component number*

See terrain component number under section *Terrain component*. The terrain component number given in the terrain component chapter should also be used here.

**35** *soil component number*

The sequence number of the soil within the terrain component according to the ranking of the soil component within the terrain component (the largest soil component is given number 1, the second largest number 2, etc.). Soil components are the lowest level of differentiation of the SOTER units.

**36** *proportion of SOTER unit*

The proportion that the soil component occupies within the SOTER unit. As stated in the section *Number of soil and terrain components*, Chapter 5, a soil component normally occupies not less than 15% of the terrain. The sum of all soil components should be 100% for each SOTER unit.

**37** *profile\_ID*

Code for the representative profile. Any national code is permitted provided it is unique at a national level. An ISO country code (see Annex 4) should precede the national code. There is room for 12 characters.

**38** *number of reference profiles*

The number of reference profiles that were considered for the selection of the representative profile is indicated. These profiles have also contributed to the determination of maximum and minimum values for a number of chemical and physical parameters of the soil.

**39** *position in terrain component*

The relative position of the soil component within the terrain component is characterized by one of the following descriptions:

<b>H</b>	high	interfluvial, crest or higher part of the terrain component
<b>M</b>	middle	upper and middle slope or any other medium position within the terrain component
<b>L</b>	low	lower slope or lower part of the terrain component
<b>D</b>	lowest	depression, valley bottom or any other lowest part of the terrain component
<b>A</b>	all	all positions within the terrain component

**40** *surface rockiness*

The percentage coverage of rock outcrops according to the following classes (FAO, 1990):

<b>N</b>	none	0 %
<b>V</b>	very few	0- 2 %
<b>F</b>	few	2- 5 %
<b>C</b>	common	5-15 %
<b>M</b>	many	15-40 %
<b>A</b>	abundant	40-80 %
<b>D</b>	dominant	≥ 80 %

**41** *surface stoniness*

The percentage cover of coarse fragments (> 0.2 cm), completely or partly at the surface, is described according to the following classes (FAO, 1990):

<b>N</b>	none	0 %
<b>V</b>	very few	0- 2 %
<b>F</b>	few	2- 5 %
<b>C</b>	common	5-15 %
<b>M</b>	many	15-40 %
<b>A</b>	abundant	40-80 %
<b>D</b>	dominant	≥ 80 %

*OBSERVABLE EROSION*

Any visible signs of (accelerated) erosion are to be indicated according to type, area affected and degree. If more than two types of erosion are active at the same time, then only the dominant type is indicated (items 42-44).

**42** *types of erosion/deposition*

Characterization of the erosion or deposition type according to FAO (1990):

<b>N</b>	no visible evidence of erosion
<b>S</b>	sheet erosion
<b>R</b>	rill erosion
<b>G</b>	gully erosion
<b>T</b>	tunnel erosion
<b>P</b>	deposition by water
<b>W</b>	water and wind erosion
<b>L</b>	wind deposition
<b>A</b>	wind erosion and deposition
<b>D</b>	shifting sand
<b>Z</b>	salt deposition
<b>U</b>	type of erosion unknown

**43** *area affected*

The area affected by the above mentioned erosion. Classes according to UNEP-ISRIC (1988).

- 1** 0- 5 %
- 2** 5-10 %
- 3** 10-25 %
- 4** 25-50 %
- 5**  $\geq$  50 %

**44** *degree of erosion*

After FAO (1990).

- |          |          |  |
|----------|----------|--|
| <b>S</b> | slight   | Some evidence of loss of surface horizons. Original biofunctions largely intact.   |
| <b>M</b> | moderate | Clear evidence of removal or coverage of surface horizons. Original biofunctions partly destroyed.   |
| <b>V</b> | severe   | Surface horizons completely removed (with subsurface horizons exposed) or covered up by sedimentation of material from upslope. Original biofunctions largely destroyed. |
| <b>E</b> | extreme  | Substantial removal of deeper subsurface horizons (badlands). Complete destruction of original biofunctions.   |

**45** *sensitivity to capping*

The degree in which the soil surface has a tendency to capping and sealing (FAO, 1990):

- |          |          |   |
|----------|----------|---|
| <b>N</b> | none     | no capping or sealing observed  |
| <b>W</b> | weak     | the soil surface has a slight sensitivity to capping. Soft or slightly hard crust less than 0.5 cm thick.                                 |
| <b>M</b> | moderate | the soil has a moderate sensitivity to capping. Soft or slightly hard crust more than 0.5 cm thick, or hard crust less than 0.5 cm thick. |
| <b>S</b> | strong   | the soil surface has a strong sensitivity to capping. Hard crust more than 0.5 cm thick.  |

**46** *rootable depth*

Estimated depth in cm to which root growth is unrestricted by any physical or chemical impediment, such as an impenetrable or toxic layer. Strongly fractured rocks, such as shales, may be considered as rootable. Classes after FAO (1990).

- |          |              |           |
|----------|--------------|-----------|
| <b>V</b> | very shallow | $<$ 30 cm |
|----------|--------------|-----------|



<b>S</b>	shallow	30- 50 cm
<b>M</b>	moderately deep	50-100 cm
<b>D</b>	deep	100-150 cm
<b>X</b>	very deep	≥ 150 cm

#### 47 *relation with other soil components*

A free-format space of 254 characters is available to indicate succinctly the relationship between this soil component and adjoining soil components. Up to 254 characters are permitted.

For example: "Soil component A has formed in colluviated material derived from soil component B".

### PROFILE

#### 48 *profile\_ID*

Same as profile\_ID in the section *Soil component*.

#### 49 *profile database\_ID*

The identification code for the owner, institute or organisation that holds (part of) the national soil profile database. The code consists of an ISO code for the country (see Annex 5) and a sequence number (see also the section *Soil profile database* in Chapter 8).

### LOCATION OF THE REPRESENTATIVE PROFILE

The latitude and longitude, as accurate as possible, and expressed in decimal degrees. A profile of which the approximate location (i.e. accurate to the nearest full minute) is not known cannot be accepted in the SOTER database.

#### 50 *latitude*

The latitude is stored in decimal degrees north. Latitudes in the southern hemisphere are negative.

#### 51 *longitude*

The longitude is stored in decimal degrees east. Longitudes in the western hemisphere are negative.

#### 52 *elevation*

The elevation of the representative profile in metres above sea level, and at least indicated to the nearest 50 m contour (if this is not possible, no entry should be made).

**53** *sampling date*

The date at which the profile was described and sampled. In case these two activities were carried out on different dates, the date of sampling should be taken. The format is MM/YYYY.

**54** *lab\_ID*

The code for the soil laboratory that analyzed the samples: ISO country code followed by a sequence number.

**55** *drainage*

The present drainage of the soil component is described according to one of the classes mentioned below (after FAO, 1990).

<b>E</b>	excessively drained	Water is removed from the soil very rapidly.
<b>S</b>	somewhat excessively drained	Water is removed from the soil rapidly.
<b>W</b>	well drained	Water is removed from the soil readily but not rapidly.
<b>M</b>	moderately well drained	Water is removed from the soil somewhat slowly during some periods of the year. The soils are wet for short periods within rooting depth.
<b>I</b>	imperfectly drained	Water is removed slowly so that the soils are wet at shallow depth for a considerable period.
<b>P</b>	poorly drained	Water is removed so slowly that the soils are commonly wet for considerable periods. The soils commonly have a shallow water table.
<b>V</b>	very poorly drained	Water is removed so slowly that the soils are wet at shallow depth for long periods. The soils have a very shallow water table.

**56** *infiltration rate*

The basic infiltration rate, in cm/h, is indicated according to the following 7 categories (BAI, 1991).

<b>V</b>	very slow	< 0.1 cm h <sup>-1</sup>
<b>S</b>	slow	0.1- 0.5 cm h <sup>-1</sup>
<b>D</b>	moderately slow	0.5- 2.0 cm h <sup>-1</sup>
<b>M</b>	moderate	2.0- 6.0 cm h <sup>-1</sup>
<b>R</b>	rapid	6.0-12.5 cm h <sup>-1</sup>
<b>Y</b>	very rapid	12.5-25.0 cm h <sup>-1</sup>
<b>E</b>	extremely rapid	≥ 25 cm h <sup>-1</sup>

**57** *surface organic matter*

Any litter or other organic matter on the surface will be described according to thickness (in cm) and degree of decomposition (Soil Survey Staff, 1975):

<b>F</b>	fibric	weakly decomposed organic soil material (fibre content >2/3 of volume)
<b>H</b>	hemic	degree of decomposition intermediate between fibric and sapric (fibre content between 1/6 and 2/3 of volume)
<b>S</b>	sapric	highly decomposed organic soil material (fibre content <1/6 of volume)

**58** *classification*

Characterisation of profile according to the revised FAO-Unesco Soil Map of the World Legend (FAO, 1988). The codes as given in this publication must be entered (see Annex 2 and for a full description also FAO, 1989). Where possible the characterization should be up to subunit level.

**59** *classification version*

The year of publication of the version of the FAO Legend used for the characterization.

**60** *national classification*

The original national classification of the representative profile if different from item 58. Up to 12 characters are permitted.

**61** *Soil Taxonomy*

Only the Soil Taxonomy classification (for codes see FAO, 1989) for representative profiles as is indicated in the national database or relevant report, is given. No entry will be made for soil profiles that were not originally classified according to Soil Taxonomy.

**62** *phase*

Any potentially limiting factor related to surface or subsurface features of the terrain, and not already specifically described in the soil profile, can be made a phase (see FAO, 1989). The coding for phases currently used by FAO is given in the FAO-ISRIC Soil Database (FAO, 1989). A note should be made on the code for new phases recognized.

**HORIZON DATA**

This section provides the attributes for the various horizons that have been distinguished in the representative soil profile. In general, no more than 5 horizons should be described. Mandatory attributes must always be completed. If these data are not available, expert estimates are required. Expert estimates are also permitted for optional attributes. Measured data are entered as an actual value for the representative profile, and as maximum and

minimum values derived from all the reference profiles of the soil component. Mandatory attributes are marked both in Table 1 and in the text.

**63** *profile\_ID* (mandatory)

Same as *profile\_ID* in sections *Soil component* and *Profile*.

**64** *horizon number* (mandatory)

A consecutive number, starting with the surface horizon, is allocated to each horizon.

**65** *diagnostic horizon* (mandatory)

Descriptions are taken from the Revised Legend of the FAO/Unesco Soil Map of the World (FAO, 1988). For more precise definitions refer to this publication.

- HI** histic      An horizon which is more than 20 cm but less than 40 cm thick. It can be more than 40 cm but less than 60 cm thick if it consists of 75 percent or more, by volume, of sphagnum fibres or has a bulk density when moist of less than 0.1 kg.dm<sup>-3</sup>. A surface layer less than 25 cm thick qualifies as a histic horizon if, after having been mixed to a depth of 25 cm, it has 16% or more organic carbon and the mineral fraction contains more than 60% clay, or 8% or more organic carbon for intermediate contents of clay.
- MO** mollic      A horizon with the following properties for the upper 18 cm:  
 1) the soil structure is sufficiently strong that the horizon is not both massive and hard or very hard when dry. Very coarse prisms larger than 30 cm in diameter are included in the meaning of massive if there is no secondary structure within the prisms.  
 2) the chroma is less than 3.5 when moist, the value darker than 3.5 when moist and 5.5 when dry; the colour value is at least one unit darker than that of the C (both moist and dry). If a C horizon is not present, comparison should be made with the horizon immediately underlying the A horizon. If there is more than 40% finely divided lime, the limits of the colour value dry are waived; the colour value moist should then be 5 or less.  
 3) the base saturation (by NH<sub>4</sub>OAc) is 50% or more  
 4) the organic carbon content is at least 0.6% throughout the thickness of mixed soil, as specified below. It is at least petrocalcic or a petrogypsic horizon or a petroferric phase.
- FI** fimic      A man made surface layer 50 cm or more thick which has been produced by long continued manuring with earthy mixtures. If a fimic horizon meets the requirements of the mollic or umbric horizon, it is distinguished from it by an acid-extractable P<sub>2</sub>O<sub>5</sub> content which is higher than 250 mg.kg<sup>-1</sup> soil by 1 percent citric acid. Examples are the plaggen epipedon and the anthropic epipedon of Soil Taxonomy.

- UM** umbric Comparable to mollic in colour, organic carbon and phosphorus content, consistency, structure and thickness. However, the base saturation is less than 50%.
- OC** ochric The horizon is too light in colour, has too high a chroma, too little organic carbon, or is too thin to be a mollic or umbric, or is both hard and massive when dry. Finely stratified materials do not qualify as an ochric horizon, e.g. surface layers of fresh alluvial deposits.
- AR** argic A subsurface horizon which has a distinctly higher clay content than the overlying horizon. This difference may be due to an illuvial accumulation of clay, or to a destruction of clay in the surface horizon, or to a selective surface erosion of clay, or to biological activity or to a combination of two or more of these different processes. Sedimentation of surface materials, which are coarser than the subsurface horizon, may enhance a pedogenic textural differentiation. However, a mere lithological discontinuity, such as may occur in alluvial deposits, does not qualify as an argic horizon. When an argic horizon is formed by clay illuviation, clay skins may occur on ped surfaces, in fissures, in pores, and in channels. The texture must be sandy loam or finer with at least 8% clay.
- NA** natric An argic horizon with
- 1) a columnar or prismatic structure in some part of the horizon, or a blocky structure with tongues of an eluvial horizon in which there are uncoated silt or sand grains extending more than 2.5 cm into the horizon, and
  - 2) an exchangeable sodium percentage of more than 15% within the upper 40 cm of the horizon; or more exchangeable magnesium plus sodium than calcium plus exchange acidity within the upper 40 cm of the horizon if the saturation with exchangeable sodium is more than 15% in some subhorizon within 200 cm of the surface.
- CB** cambic An altered horizon lacking properties that meet the requirements of an argic, natric or spodic horizon; lacking the dark colours, organic matter content and structure of the histic horizon, or the mollic and umbric horizons. The texture is sandy loam or finer, with at least 8% of clay; the thickness is at least 15 cm with the lower depth at least 25 cm below the surface; soil structure is at least moderately developed or rock structure is absent in at least half the volume of the horizon; the CEC is more than 160 mmol<sub>c</sub> kg<sup>-1</sup> clay, or the content of weatherable minerals in the 0.050 to 0.200 mm fraction is 10% or more; the horizon shows alteration in a) stronger chroma, redder hue, or higher clay content than the underlying horizon, or b) evidence of removal of carbonates, or c) if carbonates are absent in the parent material and in the dust that falls on the soil, the required evidence of alteration is satisfied by the presence of soil structure and the absence of rock structure in more than 50% of the horizon; shows no cementation, induration or brittle consistence when moist.

- SP** spodic A spodic horizon meets one of the following requirements below a depth of 12.5 cm:
- 1) a subhorizon more than 2.5 cm thick that is continuously cemented by a combination of organic matter with iron and/or aluminium
  - 2) a sandy or coarse-loamy texture with distinct dark pellets of coarse silt size or larger or with sand grains covered with cracked coatings which consist of organic matter and aluminium with or without iron.
  - 3) one or more subhorizons in which a) if there is 0.1% or more extractable iron, the ratio of iron plus Al extractable by pyrophosphate at pH 10 to clay% is 0.2 or more, or if there is less than 0.1% extractable iron, the ratio of Al plus organic carbon to clay is 0.2 or more; and b) the sum of pyrophosphate-extractable Fe+Al is half or more of the sum of dithionite-citrate extractable Fe+Al; and c) the thickness is such that the index of accumulation of amorphous material in the subhorizons that meet the preceding requirements is 65 or more. This index is calculated by subtracting half the clay% from CEC at pH 8.2  $\text{mmol}_c \text{kg}^{-1}$  clay and multiplying the remainder by the thickness of the subhorizon in cm. The results of all subhorizons are then added.
- FA** ferralic The ferralic horizon has a texture that is sandy loam or finer with at least 8% of clay; is at least 30 cm thick; has a CEC equal to or less than  $160 \text{ mmol}_c \text{kg}^{-1}$  clay or has an effective CEC equal to or less than  $120 \text{ mmol}_c \text{kg}^{-1}$  clay (sum of  $\text{NH}_4\text{OAc}$  exchangeable bases plus 1M KCl-exchangeable acidity); has less than 10% weatherable minerals in the 0.050 to 0.200 mm fraction; has less than 10% water-dispersible clay; has a silt-clay ratio which is 0.2 or less; does not have andic properties; has less than 5% by volume showing rock structure.
- CA** calcic A horizon of accumulation of calcium carbonate. The horizon is enriched with secondary calcium carbonate over a thickness of 15 cm or more, has a calcium carbonate content of 15% or more and at least 5% greater than that of a deeper horizon. The latter requirement is expressed by volume if the secondary carbonates in the calcic horizon occur as pendants on pebbles, or as concretions or soft powdery forms. If such a calcic horizon rests on very calcareous materials (40% or more calcium carbonate equivalent), the percentage of carbonates need not decrease with depth.
- PC** petrocalcic A continuous cemented or indurated calcic horizon, cemented by calcium carbonate and in places by calcium and some magnesium carbonate. Accessory silica may be present. The petrocalcic horizon is continuously cemented to the extent that dry fragments do not slake in water and roots cannot enter. It is massive or platy, extremely hard when dry so that it cannot be penetrated by spade or auger, and very firm to extremely firm when moist. Non-capillary pores are filled; hydraulic conductivity is moderately slow to very slow. It is usually thicker than 10 cm.

<b>GY</b>	gypsic	The gypsic horizon is enriched with secondary calcium sulphate ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ), is 10 cm or more thick, has at least 5% more gypsum than the underlying horizon, and the product of the thickness (cm) and the percent of gypsum is 150 or more.
<b>PG</b>	petrogypsic	A gypsic horizon that is so cemented with gypsum that dry fragments do not slake in water and roots cannot enter. The gypsum content usually exceeds 60%.
<b>SU</b>	sulphuric	The sulphuric horizon forms as a result of artificial drainage and oxidation of mineral or organic materials which are rich in sulphides. It is at least 15 cm thick and characterized by a pH-H <sub>2</sub> O less than 3.5 and generally has jarosite mottles with a hue of 2.5Y or more and a chroma of 6 or more.
<b>AL</b>	albic	Clay and free iron oxides have been removed, or the oxides have been segregated to the extent that the colour of the horizon is determined by the colour of the primary sand and silt particles rather than by coatings of these particles. An albic horizon has a colour value moist of 4 or more, or a value dry of 5 or more, or both. If the value dry is 7 or more, or the value moist is 6 or more, the chroma is 3 or less. If the value dry is 5 or 6, or the value moist 4 or 5, the chroma is closer to 2 than to 3. If the parent materials have a hue of 5YR or redder, a chroma moist of 3 is permitted in the albic horizon where the chroma is due to the colour of uncoated silt or sand grains.

## 66 diagnostic property (mandatory)

Diagnostic properties (FAO, 1988).

<b>TC</b>	abrupt textural change	A clay increase between two layers, which takes place over a distance of less than 5 cm, where the lower layer shows a clay content of twice the clay content of the overlying layer if the latter has less than 20% clay, or an increase of 20% or more if the latter has 20% clay or more.
<b>AD</b>	andic properties	Soil materials which meet one or more of the following requirements: <ol style="list-style-type: none"> <li>1) acid oxalate extractable Al plus 1/2 acid oxalate extractable Fe is 2.0% or more in the fine earth fraction; bulk density of the fine earth fraction, measured in the field moist state, is <math>0.9 \text{ kg dm}^{-3}</math> or less; phosphate retention is more than 85%.</li> <li>2) more than 60% by volume of the whole soil is volcani-clastic material coarser than 2 mm; acid oxalate extractable Al plus 1/2 acid oxalate extractable Fe is 0.40% or more in the fine earth fraction.</li> <li>3) the 0.02 to 2.0 mm fraction is at least 30% of the fine earth fraction and meets one of the following: a) if the fine earth fraction has acid oxalate extractable Al plus 1/2 acid oxalate extractable Fe</li> </ol>

of 0.40% or less, there is at least 30% volcanic glass in the 0.02 to 2.0 mm fraction; or b) if the fine earth fraction has acid oxalate extractable Al plus 1/2 acid oxalate extractable Fe of 2.0% or more, there is at least 5% volcanic glass in the 0.02 to 2.0 mm fraction; or c) if the fine earth fraction has acid oxalate extractable Al plus 1/2 acid oxalate extractable Fe of between 0.40 and 2.0%, there is a proportional content of volcanic glass in the 0.02 to 2.0 mm fraction between 30 and 5%.

<b>CO</b>	calcareous	Soil material which shows strong effervescence with 10% HCl or which contains more than 2% calcium carbonate equivalent.
<b>CA</b>	calcaric	Soils which are calcareous throughout the depth between 20 and 50 cm.
<b>RO</b>	continuous hard rock	The underlying material is sufficiently coherent and hard when moist to make hand digging with a spade impractical. The material is continuous except for a few cracks produced in place without significant displacement of the pieces and horizontally distant to an average of 10 cm or more. The material considered here does not include subsurface horizons such as a duripan, a petrocalcic or a petrogypsic horizon or a petroferric phase.
<b>FA</b>	ferralic  properties	The term 'ferralic properties' is used in connection with Cambisols and Arenosols which have a CEC of less than 240 mmol <sub>c</sub> kg <sup>-1</sup> clay or less than 40 mmol <sub>c</sub> kg <sup>-1</sup> soil in at least one subhorizon of the cambic horizon or the horizon immediately underlying the A horizon.
<b>FI</b>	ferric properties	Many coarse mottles with hues redder than 7.5YR or chromas more than 5 or both; discrete nodules, up to 2 cm in diameter, the exteriors of the nodules being enriched and weakly cemented or indurated with Fe and having redder hues or stronger chromas than the interiors (Luvisols, Alisols, Lixisols and Acrisols).
<b>FL</b>	fluvic properties	Fluviatile, marine and lacustrine sediments, which receive fresh materials at regular intervals, and which, unless empoldered, have one or both of the following properties: 1) an organic carbon content that decreases irregularly with depth or that remains above 0.20% to a depth of 125 cm. Thin strata of sand may have less organic carbon if the finer sediments below, exclusive of buried horizons, meet the requirement; 2) stratification in at least 25% of the soil within 125 cm of the surface.



<b>GE</b>	geric properties	Soil materials which have either: 1) 1.5 cmol <sub>c</sub> kg <sup>-1</sup> clay or less of exchangeable bases (Ca, Mg, K, Na) plus unbuffered 1M KCl exchangeable acidity; or 2) a delta pH (pH KCl minus pH H <sub>2</sub> O) of +0.1 or more.
<b>GL</b>	gleyic and stagnic properties	Soil materials which are saturated with water at some period of the year or throughout the year, in most years, and which show evidence of reduction processes or of reduction and segregation of iron.
<b>GY</b>	gypsiferous	Soil material which contains 5% or more gypsum.
<b>IN</b>	inter fingering	Penetrations of an albic horizon into an underlying argic or natric horizon along ped faces, primarily vertical faces. The penetrations are not wide enough to constitute tonguing, but form continuous skeletons (ped coatings of clean silt or sand, more than 1 mm thick on the vertical ped faces).
<b>NI</b>	nitic properties	Soil material that has 30% or more clay, has a moderately strong angular blocky structure which falls easily apart into flat edged ('polyhedral' or 'nutty') elements which show shiny ped faces that are either thin clay coatings or pressure faces. This soil structure is apparently associated with the presence of significant amounts of active iron oxides and is indicative of a high effective moisture storage and favourable phosphate sorption - desorption properties.
<b>OR</b>	organic soil materials	Organic soil materials are: 1) saturated with water for long periods or are artificially drained and, excluding live roots, a) have 18% or more organic carbon if the mineral fraction is 60% or more clay, b) have 12% or more organic carbon if the mineral fraction has no clay, or c) have a proportional content of organic carbon between 12 and 18% if the clay content of the mineral fraction is less than 60%; or 2) never saturated with water for more than a few days and have 20% or more organic carbon.
<b>PE</b>	permafrost	Permafrost is a layer in which the temperature is perennially at or below 0°C.
<b>PL</b>	plinthite	Plinthite is an iron-rich, humus-poor mixture of clay with quartz and other diluents. It commonly occurs as red mottles, usually in platy, polygonal or reticulate patterns, and changes irreversibly to a hardpan or to irregular aggregates on exposure to repeated wetting and drying. In a moist soil, plinthite is usually firm but it can be cut with a spade. When irreversibly hardened the material is no longer considered plinthite. Such hardened material is shown as a petroferric or a skeletal phase.
<b>SA</b>	salic properties	The electric conductivity of the saturation extract is more than 15 dS m <sup>-1</sup> within 30 cm of the surface, or more than 4 dS m <sup>-1</sup> within 30 cm of the surface if the pH-H <sub>2</sub> O exceeds 8.5.

<b>SI</b>	slickensides	Slickensides are polished and grooved surfaces that are produced by one mass sliding past another. Some of them occur at the base of a slip surface where a mass of soil moves downward on a relatively steep slope. Slickensides are very common in swelling clays in which there are marked seasonal changes in moisture content.
<b>SM</b>	smeary consistence	Thixotropic soil material; it changes under pressure or by rubbing from a plastic solid into a liquefied stage and back to the solid condition. In the liquefied stage the material skids or smears between the fingers (Andosols).
<b>SO</b>	sodic properties	The exchangeable sodium percentage is 15% or more, or exchangeable sodium plus magnesium is 50% or more.
<b>SL</b>	soft powdery lime	Translocated authigenic lime, soft enough to be cut readily with a finger nail, precipitated in place from the soil solution rather than inherited from a soil parent material. It should be present in a significant accumulation (coatings on pores or structural faces).
<b>HU</b>	strongly humic	Soil material with an organic carbon content of more than 14 g/kg fine earth as a weighted average over a depth of 100 cm from the surface. This calculation assumes a bulk density of 1.5 kg dm <sup>-3</sup> .
<b>SU</b>	sulphidic materials	Sulphidic materials are waterlogged mineral or organic soil containing 0.75% or more sulphur (dry weight), mostly in the form of sulphides, having less than three times as much calcium carbonate equivalent as sulphur, and having a pH above 3.5. Sulphidic materials accumulate in a soil that is permanently saturated and having a pH above 3.5, generally with brackish water. If the soil is drained the sulphides oxidize to form sulphuric acid. The pH, which is normally near neutrality before drainage, drops below 3.5. At this point these materials become a sulphuric horizon. Sulphidic material differs from the sulphuric horizon in its reduced condition, its pH and the absence of jarosite mottles with a hue of 2.5Y or more or a chroma of 6 or more.
<b>TO</b>	tonguing	An albic horizon penetrates an argic horizon along ped surfaces, if peds are present. Tongues must have greater depth than width, have horizontal dimensions of 5 mm or more in fine textured argic horizons (clay, silty clay and sandy clay), 10 mm or more in moderately fine textured argic horizons, and 15 mm or more in medium or coarser textured argic horizons (silt loams, loams and sandy loams), and must occupy more than 15% of the mass of the upper part of the argic horizon.
<b>VE</b>	vertic properties	In connection with clayey soils which at some period in most years show one or more of the following: cracks, slickensides, wedge-shaped or parallelepiped structural aggregates, that are not in a combination, or are not sufficiently expressed, for the soils to qualify as Vertisols.

**WM** weatherable minerals Minerals included are those that are unstable in a humid climate relative to other minerals, such as quartz and 1:1 lattice clays, and that, when weathering occurs, liberate plant nutrients and iron or aluminium. They include: 1) clay minerals: all 2:1 lattice clays except aluminium-interlayered chlorite. Sepiolite, talc and glauconite are also included in the meaning of this group of weatherable clay minerals, although they are not always of clay size. 2) silt- and sand-size minerals: feldspars, feldspathoids, ferromagnesian minerals, glasses, micas, and zeolites.

## 67 horizon designation

Master horizon with subordinate characteristics according to the rules given below (for more details see FAO, 1990).

Master horizons

**H** H horizon/layer. Layer dominated by organic material, formed from accumulations of (partially) undecomposed organic material at the soil surface, which may be underwater. All H horizons are saturated with water for prolonged periods, or were once saturated but are now artificially drained. An H horizon may be on top of mineral soils or at any depth beneath the surface if it is buried.

**O** O horizon/layer. Layer dominated by organic material, consisting of (partially) undecomposed litter, such as leaves, twigs, moss etc., which has accumulated on the surface. They may be on top of either mineral or organic soils. An O horizon are not saturated with water for prolonged periods. The mineral fraction of such material is only a small percentage of the volume of the material and generally is much less than half the weight. An O horizon may be at the surface of a mineral soil or at any depth beneath the surface if it is buried.

**A** A horizon. Mineral horizon which formed at the surface or below an O horizon, and in which all or much of the original rock structure has been obliterated. The A horizon is characterised by one or more of the following:

- an accumulation of humified organic matter intimately mixed with the mineral fractions and not displaying properties characteristic of an E horizon (see below);
- properties resulting from cultivation, pasturing, or similar kinds of disturbance; or
- a morphology which is different from the underlying B or C horizon, resulting from processes related to the surface (e.g. vertisols).

**E** E horizon. Mineral horizon, in which the main feature is a loss of silicate clay, iron, aluminum, or some combination of these, leaving a concentration of sand and silt particles, and in which all or much of the original rock structure has been obliterated.

An E horizon is most commonly differentiated from an underlying B horizon by colour of higher value or lower chroma, or both; by coarser texture; or by a

combination of these. Although an E horizon is usually near the surface, below an O or A horizon, and above a B horizon, the symbol E may be used without regard to position in the profile for any horizon that meets the requirements, and that has resulted from soil genesis.

- B** B horizon. A B horizon has formed below an A, E, O or H horizon, and has as dominant feature the obliteration of all or much of the original rock structure, together with one or a combination of the following:
- illuvial concentration, alone or in combination, of silicate clay, iron, aluminum, humus, carbonates, gypsum or silica;
  - evidence of removal of carbonates;
  - residual concentration of sesquioxides;
  - coating of sesquioxides that make the horizon conspicuously lower in value, higher in chroma, or redder in hue than overlying and underlying horizons without apparent illuviation of iron;
  - alteration that forms silicate clay or liberates oxides or both and that forms a granular, blocky or prismatic structure if volume changes accompany the changes in moisture content, or
  - brittleness.

Layers with gleying but no other pedogenetic change are not considered a B horizon.

- C** C horizon/layer. A horizon or layer, excluding hard bedrock, that is little affected by pedogenetic processes and lacks properties of H, O, A, E or B horizons. Most are mineral layers, but some siliceous or calcareous layers (e.g. shells, coral and diatomaceous earth) are included. Sediments, saprolite and unconsolidated bedrock and other geological materials that commonly slake within 24 hours are included as C layers. Some soils form in highly weathered material that is considered a C horizon if it does not meet the requirements of an A, E or B horizon.

- R** R layer. Hard rock underlying the soil. Air dry chunks of an R layer will not slake within 24 hours if placed into water.

#### Subordinate properties

Subordinate distinctions and features within master horizons are indicated with lower case letters used as suffixes. The following subordinate properties may be used (see FAO, 1990 for more details).

- b** buried genetic horizon
- c** concretions or nodules
- f** frozen soil
- g** strong gleying
- h** accumulation of organic matter
- j** jarosite mottling
- k** accumulation of carbonates
- m** cementation or induration
- n** accumulation of sodium
- o** residual accumulation of sesquioxides

- p** ploughing or other disturbance
- q** accumulation of silica
- r** strong reduction
- s** illuvial accumulation of sesquioxides
- t** accumulation of silicate clay
- v** occurrence of plinthite
- w** development of colour or structure
- x** fragipan character
- y** accumulation of gypsum
- z** accumulation of salts more soluble than gypsum

**68** *lower depth* (mandatory)

The average depth of the lower boundary in cm (the upper boundary in the case of an **O** horizon).

**69** *distinctness of transition*

Abruptness of horizon boundary to underlying horizon (FAO, 1990).

- |          |         |         |
|----------|---------|---------|
| <b>A</b> | abrupt  | 0- 2 cm |
| <b>C</b> | clear   | 2- 5 cm |
| <b>G</b> | gradual | 5-15 cm |
| <b>D</b> | diffuse | ≥ 15 cm |

**70** *moist colour* (mandatory)

The Munsell colours (moist soil) should be given. Only integer values and chromas are accepted.

**71** *dry colour*

The Munsell colours (dry soil) should be given. Only integer values and chromas are accepted.

**STRUCTURE**

The grade, size and type of structure, defined according to FAO (1990), are described in items 72-74.

**72** *grade of structure*

- |          |               |   |
|----------|---------------|---|
| <b>N</b> | structureless | no observable aggregation or no orderly arrangement of natural planes of weakness (massive or single grain)   |
| <b>W</b> | weak          | soil with poorly formed indistinct peds, that are barely observable in place even in dry soil, breaks up into very few intact peds, many broken peds and much apedal material |

- M** moderate soil with well-formed distinct peds, durable and evident in disturbed soil which produces many entire peds, some broken peds and little apedal material
- S** strong soil with durable peds that are clearly evident in undisturbed (dry) soil, which breaks up mainly into entire peds

### 73 size of structure elements

TABLE 4

**Size classes for structure elements of various types (mm)** (Soil Survey Staff, 1951; FAO, 1990)

Size classes		Ranges of size of structure elements (mm)				
		platy	prismatic/columnar	(sub)ang.blocky	granul.	crumb
<b>V</b>	very fine	< 1	< 10	< 5	< 1	<1
<b>F</b>	fine	1- 2	10 - 20	5 - 10	1- 2	1-2
<b>M</b>	medium	2- 5	20 - 50	10 - 20	2- 5	2-5
<b>C</b>	coarse	5-10	50 -100	20 - 50	5-10	
<b>X</b>	very coarse	>10	> 100	> 50	> 10	

### 74 type of structure (mandatory)

- P** platy particles arranged around a generally horizontal plane
- R** prismatic prisms without rounded upper end
- C** columnar prisms with rounded caps
- A** angular blocky bounded by plains intersecting at largely sharp angles.
- S** subangular blocky mixed rounded and plane faces with vertices mostly rounded
- G** granular spheroidal or polyhedral, relatively non-porous
- B** crumb spheroidal or polyhedral, porous
- M** massive no structure
- N** single grain no structure, individual grains
- W** wedge shaped structure in horizons with slickensides

**COARSE FRAGMENTS**

The presence of any rock or mineral fragments in the horizon is described in items 75 and 76.

**75** *abundance of coarse fragments* (mandatory)

Classes of volume % of rock or mineral fragments (> 2 mm) in soil matrix (FAO, 1990).

<b>N</b>	none	0 %
<b>V</b>	very few	0- 2 %
<b>F</b>	few	2- 5 %
<b>C</b>	common	5-15 %
<b>M</b>	many	15-40 %
<b>A</b>	abundant	40-80 %
<b>D</b>	dominant	80 %

**76** *size of coarse fragments*

Size of dominant rock or mineral fragments in classes (FAO, 1990).

<b>V</b>	very fine	< 2 mm
<b>F</b>	fine	2- 6 mm
<b>M</b>	medium	6-20 mm
<b>C</b>	coarse	20 mm

**77** *very coarse sand*

Weight % of particles 2.0-1.0 mm in fine earth fraction.

**78** *coarse sand*

Weight % of particles 1.0-0.5 mm in fine earth fraction.

**79** *medium sand*


Weight % of particles 0.5-0.25 mm in fine earth fraction.

**80** *fine sand*

Weight % of particles 0.25-0.10 mm in fine earth fraction.

**81** *very fine sand*

Weight % of particles 0.10-0.05 mm in fine earth fraction.



**82** *total sand* (mandatory)

Weight % of particles 2.0-0.05 mm in fine earth fraction. The total sand fraction, either as an absolute value, or as the sum of the sub-fractions.

**83** *silt* (mandatory)

Weight % of particles 0.05-0.002 mm in fine earth fraction.

**84** *clay* (mandatory)

Weight % of particles < 0.002 mm in fine earth fraction.

**85** *particle size class*

The particle size class as derived, with the aid of figure 11, from the particle size analysis results.

**86** *bulk density* (mandatory)

The bulk density in  $\text{kg dm}^{-3}$ .

**87** *moisture content at various tensions*

The database accepts the soil moisture content (%) at 5 different tensions, of which one should be the moisture content at field capacity (-33 KPa) and one the moisture content at wilting point (-1500 KPa).



For example:

KPa	-33	-98	-300	-510	-1500
soil moisture %	41	22	17	12	09

**88** *hydraulic conductivity*

The saturated hydraulic conductivity in  $\text{cm h}^{-1}$ .

**89** *infiltration rate*

The basic infiltration rate in  $\text{cm h}^{-1}$ .

**90** *pH (H<sub>2</sub>O) (mandatory)*

The pH is determined in the supernatant suspension of a 1:2.5 soil-water mixture (mandatory).

**91** *pH (KCl)*

The pH is determined in the supernatant suspension of a 1:2.5 soil-1 M KCl mixture.

**92** *electrical conductivity (EC<sub>e</sub>)*

The electrical conductivity of saturation extract,  $\text{dS m}^{-1}$ , only mandatory if the soil contains salts.

*SOLUBLE SALTS*

The type and amount of soluble salts, only when the  $\text{EC}_e \geq 4 \text{ dS m}^{-1}$ , are described in items 93-100.

**93** *soluble Na<sup>+</sup>*

The soluble Na<sup>+</sup> content of the saturated paste in  $\text{cmol}_c \text{ l}^{-1}$  (= meq/l).

**94** *soluble Ca<sup>++</sup>*

The soluble Ca<sup>++</sup> content of the saturated paste in  $\text{cmol}_c \text{ l}^{-1}$ .

**95** *soluble Mg<sup>++</sup>*

The soluble Mg<sup>++</sup> content of the saturated paste in  $\text{cmol}_c \text{ l}^{-1}$ .

**96** *soluble K<sup>+</sup>*

The soluble K<sup>+</sup> content of the saturated paste in  $\text{cmol}_c \text{ l}^{-1}$ .

**97** *soluble Cl<sup>-</sup>*

The soluble Cl<sup>-</sup> content of the saturated paste in mmol<sub>c</sub> l<sup>-1</sup>.

**98** *soluble SO<sub>4</sub><sup>2-</sup>*

The soluble SO<sub>4</sub><sup>2-</sup> content of the saturated paste in cmol<sub>c</sub> l<sup>-1</sup>.

**99** *soluble HCO<sub>3</sub><sup>-</sup>*

The soluble HCO<sub>3</sub><sup>-</sup> content of the saturated paste in cmol<sub>c</sub> l<sup>-1</sup>.

**100** *soluble CO<sub>3</sub><sup>2-</sup>*

The soluble HCO<sub>3</sub><sup>-</sup> content of the saturated paste in cmol<sub>c</sub> l<sup>-1</sup>.

**101** *exchangeable Ca<sup>++</sup>*

The exchangeable Ca in cmol<sub>c</sub> kg<sup>-1</sup> (= meq/100 g).

**102** *exchangeable Mg<sup>++</sup>*

The exchangeable Mg in cmol<sub>c</sub> kg<sup>-1</sup>.

**103** *exchangeable Na<sup>+</sup>*

The exchangeable Na in cmol<sub>c</sub> kg<sup>-1</sup>.

**104** *exchangeable K<sup>+</sup>*

The exchangeable K in cmol<sub>c</sub> kg<sup>-1</sup>.

**105** *exchangeable Al<sup>+++</sup>*

The exchangeable Al in cmol<sub>c</sub> kg<sup>-1</sup>.

**106** *exchangeable acidity*

The exchangeable acidity, as determined in 1N KCl, in cmol<sub>c</sub> kg<sup>-1</sup>.

**107** *CEC soil (mandatory)*

The cation exchange capacity of the soil at pH 7.0 in cmol<sub>c</sub> kg<sup>-1</sup>.

**108** *total carbonate equivalent*

The content of carbonates in g kg<sup>-1</sup>.

**109** *gypsum*

The gypsum content in  $\text{g kg}^{-1}$ .

**110** *total carbon (mandatory)*

The content of total organic carbon in  $\text{g kg}^{-1}$ , a mandatory attribute for the topsoil (first 25 cm, or A horizon, whichever is deeper).

**111** *total nitrogen*

The content of total N in  $\text{g kg}^{-1}$ .

**112** *P<sub>2</sub>O<sub>5</sub>*

The P<sub>2</sub>O<sub>5</sub> content in  $\text{mg kg}^{-1}$ .

**113** *phosphate retention*

The phosphate retention in %.

**114** *Fe, dithionite extractable*

The Fe fraction, in weight %, extractable in dithionite.

**115** *Fe, pyrophosphate extractable*

The Fe fraction, in weight %, extractable in pyrophosphate at pH 10.

**116** *Al, dithionite extractable*

The Al fraction, in weight %, extractable in dithionite.

**117** *Al, pyrophosphate extractable*

The Al fraction, in weight %, extractable in pyrophosphate at pH 10.

**118** *clay mineralogy*

The dominant type of mineral in the clay fraction.

**AL** allophane

**CH** chloritic

**IL** illitic

**IN** interstratified or mixed

**KA** kaolinitic

**MO** montmorillonitic

**SE** sesquioxidic

**VE** vermiculitic

## **PART II**

# **LAND USE AND VEGETATION**



## Chapter 7

# Land cover

In SOTER, land cover characteristics (vegetation and land use) are stored in two files that are separated from the soil and terrain properties. Attributes of land use and vegetation are displayed in Table 5. In contrast with the more stable attributes of the land which are covered in Part I of this manual, land cover is considered a more dynamic entity which can change quickly in time. Therefore there may be a frequent need for addition of more recent data. Moreover, third parties are working on global databases for land use (FAO) and for vegetation, or are planning to do so. At present, such databases are not available but the need exists for the subsequent incorporation of these data into SOTER.

TABLE 5  
Attributes of land use and vegetation files

LAND USE	VEGETATION
1 SOTER unit_ID	1 SOTER unit_ID
2 date of observation	2 date of observation
3 land use	3 vegetation
4 proportion of SOTER unit	4 proportion of SOTER unit

For interpretative uses of the SOTER database there is a need for land cover data. A provisional system for such data is implemented for the SOTER database. In it, the land cover information is given at the level of the SOTER unit. By doing so, the effort of digitizing separate land cover boundaries is avoided and a simple link is possible between the soil and terrain data and the land cover.

### LAND USE

The land use file contains only four attributes, of which the first two, viz. SOTER unit ID and date of observation, are the key attributes.

#### 1 *SOTER unit\_ID*

Identification code of a SOTER unit (see chapter 6.1 Terrain).

#### 2 *date of observation*

Date of observation for the land use; stored in format MM/YYYY.

#### 3 *land use*

Land use classes are defined in a hierarchical system (Remmelzwaal, 1990). At the highest level, classes are subdivided into subclasses and groups on the basis of the type of land use, and the occurrence of input and/or output (animal products, crops). The codes for land use are given in Table 6 and full descriptions in Annex 3.

TABLE 6  
Hierarchy of land use: land use orders, groups, and systems

Order	Group	System
<b>S</b> SETTLEMENT/ INDUSTRIES	<b>SR</b> residential use	
	<b>SI</b> industrial use	
	<b>ST</b> transport	
	<b>SC</b> recreational	
	<b>SX</b> excavations	
<b>A</b> AGRICULTURE	<b>AA</b> annual field cropping	<b>AA1</b> shifting cultivation
		<b>AA2</b> fallow system cultivation
		<b>AA3</b> ley system cultivation
		<b>AA4</b> rainfed arable cultivation
		<b>AA5</b> wet rice cultivation
		<b>AA6</b> irrigated cultivation
	<b>AP</b> perennial field cropping	<b>AP1</b> non-irrigated
		<b>AP2</b> irrigated
	<b>AT</b> tree & shrub cropping	<b>AT1</b> non-irrigated tree crop cultivation
		<b>AT2</b> irrigated tree crop cultivation
<b>AT3</b> non-irrigated shrub crop cultivation		
<b>AT4</b> non-irrigated shrub crop cultivation		
<b>H</b> ANIMAL HUSBANDRY	<b>HE</b> extensive grazing	<b>HE1</b> nomadism
		<b>HE2</b> semi-nomadism
		<b>HE3</b> ranching
	<b>HI</b> intensive grazing	<b>HI1</b> animal production
		<b>HI2</b> dairying
<b>F</b> FORESTRY	<b>FN</b> exploitation of natural forest and woodland	<b>FN1</b> selective felling
		<b>FN2</b> clear felling
	<b>FP</b> plantation forestry	
<b>M</b> MIXED FARMING	<b>MF</b> agro-forestry	
	<b>MP</b> agro-pastoralism (cropping & livestock systems)	
<b>E</b> EXTRACTION/ COLLECTING	<b>EV</b> exploitation of natural vegetation	
	<b>EH</b> hunting and fishing	
<b>P</b> NATURE PROTECTION	<b>PN</b> nature and game preservation	<b>PN1</b> reserves
		<b>PN2</b> parks
		<b>PN3</b> wildlife management
	<b>PD</b> degradation control	<b>PD1</b> non-interference
	<b>PD2</b> with interference	
<b>U</b> UNUSED		



**4** *proportion of SOTER unit*

Proportion that the land use occupies within the SOTER unit, in %.

**VEGETATION**

The vegetation file contains four attributes, of which the first two, viz. SOTER unit ID and date of observation, are the key attributes.

**1** *SOTER unit\_ID*

Identification code of a SOTER unit (see chapter 6.1 Terrain).

**2** *date of observation*

Date of observation for the native vegetation; stored in format MM/YYYY.

**3** *vegetation*

Generalized description of the physiognomy of the present native vegetation (Unesco, 1973). Table 7 gives the hierarchical classification of the vegetation to apply at the SOTER unit level. A full description of the classes is given in Annex 4. Vegetation should be specified at least on the formation subclass level.

**4** *proportion of SOTER unit*

Proportion that the vegetation occupies within the SOTER unit, in %.

TABLE 7

**Hierarchical vegetation classes**

Class	Subclass	Group	
I	closed forest	<b>IA</b> mainly evergreen forest	<b>IA1</b> tropical ombrophilous forest
			<b>IA2</b> tropical and subtropical evergreen seasonal forest
			<b>IA3</b> tropical and subtropical semi-deciduous forest
			<b>IA4</b> subtropical ombrophilous forest
			<b>IA5</b> mangrove forest
			<b>IA6</b> temperate and subpolar evergreen ombrophilous forest
			<b>IA7</b> temperate evergreen seasonal broad-leaved forest
			<b>IA8</b> winter-rain evergreen broad-leaved sclerophyllous forest
			<b>IA9</b> tropical and subtropical evergreen needle-leaved forest
			<b>IA10</b> temperate and subpolar evergreen needle-leaved forest
..	..	<b>IB</b> mainly deciduous forest	<b>IB1</b> tropical and subtropical drought-forest
			<b>IB2</b> cold-deciduous forest with evergreen trees (or shrubs)
			<b>IB3</b> cold-deciduous forest without evergreen trees

..		<b>IC</b>	extremely xeromorphic forest	<b>IC1</b>	sclerophyllous-dominated extremely xeromorphic forest
				<b>IC2</b>	thorn-forest
				<b>IC3</b>	mainly succulent forest
<b>II</b>	woodland	<b>IIA</b>	mainly evergreen woodland	<b>IIA1</b>	evergreen broad-leaved woodland
				<b>IIA2</b>	evergreen needle-leaved woodland
..		<b>IIB</b>	mainly deciduous woodland	<b>IIB1</b>	drought-deciduous woodland
				<b>IIB2</b>	cold-deciduous woodland with evergreen trees
				<b>IIB3</b>	cold-deciduous woodland without evergreen trees
..		<b>IIC</b>	extremely xeromorphic woodland	subdivisions as extremely xeromorphic forest (IC)	
<b>III</b>	scrub	<b>IIIA</b>	mainly evergreen scrub	<b>IIIA1</b>	evergreen broad-leaved shrubland (or thicket)
				<b>IIIA2</b>	evergreen needle-leaved and microphyllous shrubland
..		<b>IIIB</b>	mainly deciduous scrub	<b>IIIB1</b>	drought-deciduous scrub with evergreen woody plants admixed
				<b>IIIB2</b>	drought-deciduous scrub without evergreen woody plants admixed
				<b>IIIB3</b>	cold-deciduous scrub
..		<b>IIIC</b>	extremely xeromorphic (subdesert) shrubland	<b>IIIC1</b>	mainly evergreen subdesert shrubland
				<b>IIIC2</b>	deciduous subdesert shrubland
<b>IV</b>	dwarf scrub and related communities	<b>IVA</b>	mainly evergreen dwarf-scrub	<b>IVA1</b>	evergreen dwarf-scrub thicket
				<b>IVA2</b>	evergreen dwarf shrubland
				<b>IVA3</b>	mixed evergreen dwarf-shrubland and herbaceous formation
..		<b>IVB</b>	mainly deciduous dwarf-scrub	<b>IVB1</b>	facultatively drought-deciduous dwarf-thicket (or dwarf-shrubland)
				<b>IVB2</b>	obligatory, drought-deciduous dwarf-thicket (or dwarf-shrubland)
				<b>IVB3</b>	cold-deciduous dwarf-thicket (or dwarf-shrubland)
..		<b>IVC</b>	extremely xeromorphic dwarf-shrubland	subdivisions as extremely xeromorphic (subdesert) shrubland (IIIC)	
..		<b>IVD</b>	tundra	<b>IVD1</b>	mainly bryophyte tundra
				<b>IVD2</b>	mainly lichen tundra
..		<b>IVE</b>	mossy bog formations with dwarf-shrub	<b>IVE1</b>	raised bog
				<b>IVE2</b>	non-raised bog
<b>V</b>	herbaceous vegetation	<b>VA</b>	tall graminoid vegetation	<b>VA1</b>	tall grassland with a tree synusia covering 10-40%
				<b>VA2</b>	tall grassland with a tree synusia <10%
				<b>VA3</b>	tall grassland with a synusia of shrubs
				<b>VA4</b>	tall grassland with a woody synusia

..	<b>VB</b>	medium tall grassland		<b>VA5</b>	tall grassland practically without woody synusia
				<b>VB1</b>	medium tall grassland with a tree synusia covering 10-40%
				<b>VB2</b>	medium tall grassland with a synusia <10%
				<b>VB3</b>	medium tall grassland with a synusia of shrubs
				<b>VB4</b>	medium tall grassland with an open synusia of tuft plants (usually palms)
				<b>VB5</b>	medium tall grassland practically without woody synusia
..	<b>VC</b>	short grassland		<b>VC1</b>	short grassland with a tree synusia covering 10-40%
				<b>VC2</b>	short grassland with a tree synusia <10%
				<b>VC3</b>	short grassland with a synusia of shrubs
				<b>VC4</b>	short grassland with an open synusia of tuft plants
				<b>VC5</b>	short grassland practically without woody synusia
				<b>VC6</b>	short to medium tall mesophytic grassland
				<b>VC7</b>	graminoid tundra
..	<b>VD</b>	forb vegetation		<b>VD1</b>	tall forb communities
				<b>VD2</b>	low forb communities
..	<b>VE</b>	hydromorphic vegetation	fresh-water	<b>VE1</b>	rooted fresh-water communities
				<b>VE2</b>	free-floating fresh-water communities

## **PART III**

### **MISCELLANEOUS FILES**



## Chapter 8

### Reference files

Tables containing information on the source materials used for the compilation of the SOTER units, generally soil maps, the laboratories that analyzed the soil samples, the laboratory methods and the organisations responsible for the national profile database are described in this chapter.

#### SOURCE MAP

In this file information on type of map, scale, location and date are stored. As the location in max and min X and Y-coordinates is recorded, the GIS can be used to overlay this information on the SOTER map. There exists a direct link (primary key 'map\_ID') between the terrain table and the source map table. The attributes are shown in Table 8.

**1** *map\_ID*

The source map identification code from which the data were derived for the compilation of the SOTER units. See also map\_ID in Chapter 6, *Terrain*.

**2** *map title*

The citation of the source map title. There is room for 40 characters.

**3** *year*

The year of publication of the source map.

**4** *scale*

The scale of the source map as a representative fraction.

**5** *minimum latitude*

The minimum latitude (Y-coordinate) of the source map, in decimal degrees North. Latitude South is a negative figure.

**6** *minimum longitude*

The minimum longitude (X-coordinate) of the source map, in decimal degrees East. Longitude West gets a negative number.

TABLE 8  
Attributes of related tables

SOURCE MAP	LABORATORY	PROFILE DATABASE
1 map_ID	1 lab_ID	1 soil profile database_ID
2 map title	2 laboratory name	2 name of institute
3 year		
4 scale	LABORATORY METHOD	
5 minimum latitude		
6 minimum longitude	3 lab_ID	
7 maximum latitude	4 date	
8 maximum longitude	5 attribute	
9 type of map	6 method of analysis_ID	
ANALYTICAL METHOD		
7 method of analysis_ID		
8 description		

**7** *maximum latitude*

The maximum latitude (Y-coordinate) of the source map, in decimal degrees North.

**8** *maximum longitude*

The maximum longitude (X-coordinate) of the source map, in decimal degrees East.

**9** *type of source map*

The type of source map:

- S** pure soil map
- M** morpho-pedological map (soil-landscapes)
- O** other map

**LABORATORY INFORMATION**

For every analysis method that has been applied in a particular laboratory separate entries in these tables should be made.

**Laboratory**

**1** *lab\_ID*

Identification code for the laboratory that analyzed the reference soil profile. A country code with a sequential number is given. See list of country codes in Annex 5.

**2** *laboratory name*

Name of the laboratory, in full (up to 40 characters).



**Laboratory method****3** *lab\_ID*

Laboratory code (see attribute 1, lab\_ID).

**4** *date*

Date at which the laboratory introduced a method for a given attribute. Format is MM/YYYY.

**5** *attribute*

Profile layer attribute that was analyzed. The item code preceding the attribute in Table 1 and in the margin is used.

**6** *method of analysis\_ID*

Identification code for the analysis method applied. This code consists of the attribute code (item 5) followed by a sequential number.

**Analytical method****7** *method of analysis\_ID*

Method code (see attribute 6).

**8** *description*

A complete description of the analytical method used. There is room for 256 characters.

**SOIL PROFILE DATABASE**

Information on the (national) soil profile database that has been consulted for the selection of the SOTER profile data can be found as an additional file. A code for the country (ISO code from Annex 5) followed by a sequence number is given. Also the name of the organization can be indicated.

**1** *profile database\_ID*

The identification code for the owner, institute or organisation that holds (part of) the national soil profile database. The code consists of an ISO code for the country (see Annex 5) and a sequence number.

**2** *name*

Name (in full) of the owner, institute or organisation of the national soil profile database and address, up to 40 characters.



## Chapter 9

# Climate

Climatic data forms an inseparable part of the basic inventory of natural resources. Nevertheless, climate is treated separately from the SOTER database as the climate data are not directly linked to the SOTER units. Climate data are based on point observations only and the link with the soils and terrain information exists by means of the geographical location of these points. The SOTER climate files are intended for multiple applications of the soils and terrain database. Monthly data are considered sufficient for most of the (small scale) applications.

At the Workshop on Procedures Manual Revisions (ISRIC, 1990b), it was recommended that the attribute data for the climate database of SOTER should be derived, if possible, from existing computerized databases, e.g. WMO (CLICOM), FAO and CIAT. Data from these databases can be imported through an ASCII file interface. Care should be taken on the units of measure.

Data from point observations are extracted from meteorological data sets and consist of two major groupings: (1) climate station particulars, and (2) monthly climate data.

The files shown in Table 9 are used to store the station particulars and the monthly climatic data as well as the date sources.

### CLIMATE STATION

#### 1 *climate station\_ID*

The climate station\_ID is given as a two-character ISO country code (according to Annex 5) followed by a four digit sequential number.

#### 2 *climate station name*

The name of the climate station is given. Up to 40 characters are permitted.

#### 3 *latitude*

The latitude is stored in decimal degrees North; latitudes in the southern hemisphere are negative.

TABLE 9

**Attributes for climate station, climate data and source tables**

CLIMATE STATION	CLIMATE DATA	DATA SOURCE
1 climate station_ID	6 climate station_ID 25 source_ID	
2 climate station name	7 kind of data	26 source name
3 latitude	8 source_ID	
4 longitude	9 first year	
5 altitude	10 last year	
	11 years of record	
	12 jan	
	..	
	23 dec	
	24 annual	

**4** *longitude*

The longitude is stored in decimal degrees East; longitudes in the western hemisphere are negative.

**5** *altitude*

The altitude above or below (negative) sea level, m.

**CLIMATE DATA****6** *climate station\_ID*

Code for the climate station. See station code under Climate station.

**7** *kind of data*

The various kinds of climatic data are treated in paragraph 9.4

**8** *source\_ID*

Identification code for the main source of the data for each separate kind of data. Codes are to be explained in the data source file (see chapter 9.6).

**9** *first year*

The first year of the observation period.

**10** *last year*

The last year of the observation period.

**11** *years*

The number of years of record in the observation period

**12...23** *jan...dec*

The data values for each individual month. Average monthly value for the numbers of years recorded.

**24** *annual*

The annual value (average or total).

**VARIOUS CLIMATE CHARACTERISTICS**

In this section various climate characteristics (attribute 8: 'kind of data') are arranged in several groups. The importance of the kind of data attribute is indicated by a letter (M = mandatory, D = desirable and O = optional). When a mandatory characteristic is missing, the station should not be included in the database.

**rainfall**

Data on rainfall is recorded in mm. The amount of rainfall is a mandatory attribute; if it is missing, it is considered of no use to include the climate station in the database.

**RAIN** M precipitation total, mm  
**RDAY** D number of rainy days; days with at least 1 mm of precipitation  
**RMAX** O maximum 24-hour rainfall, mm  
**RR75** O rainfall reliability; the amount of rainfall exceeded in 3 out of 4 years, mm

**temperature**

Temperature is stored in degrees centigrade ( $^{\circ}\text{C}$ ). Both minimum and maximum temperatures are mandatory. The average temperature is optional because it can be derived from the minimum and maximum temperatures.

**TEMP** O mean temperature during 24-hour period  
**TMIN** M minimum temperature during a 24 hour period  
**TMAX** M maximum temperature during a 24 hour period

**radiation/sunshine**

Either radiation or sunshine hours is mandatory; the other is then optional. Radiation data is preferred.

**RADI** M/O total radiation,  $\text{MJ}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$

**SUNH** O/M hours of bright sunshine per day  
**CLOU** O degree of cloudiness, octas

### humidity

Either vapour pressure or relative humidity is mandatory. Vapour pressure is preferable to above relative humidity.

**VAPP** M/O vapour pressure, mbar  
**HUMI** O/M average relative humidity during 24 hour period, %  
**HMIN** O minimum relative humidity during 24 hour period, %  
**HMAX** O maximum relative humidity during 24 hour period, %

### wind

Wind velocity in m/s.

**WIND** D mean wind velocity at 2m during 24 hour period  
**WDAY** O wind speed during day at 2m during 24 hour period  
**WNIG** O wind speed during night at 2m during 24 hour period  
**WDIR** O dominant wind direction at 2m during 24 hour period

### risk or occurrence of adverse weather events

**WRIS** O risk or occurrence of adverse weather events like severe hailstorms, hurricanes and nightfrost. Indicated on a scale of 0 (never) to 1 (every year in the month under consideration). Intermediate values are used if the frequency is less than every year (for that month). E.g.: One occurrence every 5 years in the month of March = 0.2

### evaporation

**EPAN** O class A pan evaporation, mm  
**ECOL** O Colorado pan evaporation, mm  
**EPIC** O evaporation, Piche, mm

### evapotranspiration

Because evapotranspiration is a calculated characteristic, it is optional.

**PETP** O Penman potential evapotranspiration, mm  
**PETH** O Hargreaves potential evapotranspiration, mm  
**PETT** O Thornthwaite potential evapotranspiration, mm

TABLE 10

**Example of various kinds of climatic data recorded for a climate station (Posedas, Argentina).**

Stat.	SR	Data	F-yr	L-yr	Yrs	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
AR21	06	Rain	1901	1980	80	141	148	139	146	131	127	97	99	143	189	134	149	1643
AR21	07	Rday	1951	1980	30	9.6	9.3	9.3	8.3	8.3	9.6	9.3	9.3	11.0	10.6	7.6	8.6	110.8
AR21	01	Temp	1951	1980	30	26.2	25.8	24.3	20.7	18.1	16.5	15.6	17.3	18.8	20.9	23.3	25.7	21.1
AR21	01	Tmin	1951	1980	30	19.7	19.4	18.2	14.8	12.5	11.5	10.0	11.0	12.8	14.7	16.5	18.8	15.0
AR21	01	Tmax	1951	1980	30	32.7	32.2	30.4	26.6	23.6	21.5	21.2	23.6	24.8	27.1	30.1	32.6	27.2
AR21	01	Vapp	1951	1980	30	24.2	24.5	32.0	19.3	17.5	15.9	14.2	14.7	16.5	18.5	19.7	21.8	19.2
AR21	01	Wind	1951	1980	30	1.5	1.7	1.5	1.5	1.7	1.7	2.0	2.0	2.0	2.0	1.7	1.7	1.8
AR21	01	PETP	1951	1980	30	149	125	105	69	45	32	41	63	74	104	138	161	1109

**ADDITIONAL CONVENTIONS**

Data can be given for different categories of climate characteristics:

For Penman calculations, mandatory data are minimum and maximum temperature, irradiation, vapour pressure or relative humidity, wind speed, monthly rainfall, and number of rainy days.

When data are missing, some parameters can be estimated from others:

- relative humidity and vapour pressure can be estimated from each other
- radiation, sunshine hours, and cloudiness degree
- minimum and maximum temperature determine average temperature.

**DATA SOURCES**

One related file to the climate database exists: data sources. It contains one key field namely the source\_ID of the climate data file and one attribute: the full name of the source (published report, or name and address of the meteorological organisation holding the complete climate dataset).

**25 source\_ID**

Identification code for the source of data (as item 8).

**26 source name**

The full name of the source from which the climatic data have been taken.





## Annex 1

# Hierarchy of landforms

The term landform as used in this manual, is land with a characteristic slope (see also Remmelzwaal, 1990). Landform separation (first and second level) is thus based on morphometric criteria, chief amongst which is the slope gradient. The relief intensity is the second most important criterion used to subdivide the landscape. Subdivisions of level lands also take into account the position of the landform vis-à-vis the surrounding land. Further separation of the landforms according to hypsometric criteria is different for each 1st level landform (see item 10). Exceptions to this are noted with the description of the 2nd level landforms. The classification as presented here has been tested for a 1:5 million physiographic inventory of South America and Africa (Eschweiler, 1993 and Wen, 1993).

### 1ST LEVEL LANDFORMS

#### LEVEL LAND

Level lands are all lands with dominant slopes between 0 and 8% ( $0^\circ$  and  $4^\circ 40'$ ). Moreover, the relief intensity is such that the difference between the highest and the lowest point within one slope unit is mostly less than 50 m.

#### SLOPING LAND

Sloping land embraces all landforms that have dominant slopes between 8% and 30%, combined with in most cases a relief intensity of more than 50 m per slope unit. In general, sloping land will be more heterogeneous with respect to its slope than level land.

#### STEEP LAND

Steep land is mainly confined to mountainous country, where average slopes are over 30% (the variability of slope gradients may be so much as to make it difficult to recognize a dominant slope) and the relief intensity is more than 600 m/2 km.

#### LANDS WITH COMPOSITE LANDFORMS

Two strongly contrasting landforms, themselves not separable at the scale of mapping, may be combined if they are part of an outstanding landform that as such can be delineated at the scale of mapping. Examples of such landforms associations are valleys, made up of side-slopes and a valley bottom, and narrow plateaux, where a level surface is surrounded by relative steeply sloping land. Not all possible combinations are given here and the user may define others if the need for them arises (e.g. deeply incised plateau, consisting of a plateau

and high-gradient valleys).

## **2ND LEVEL LANDFORMS**

### **L Level lands**

Except for low-gradient footslopes, all types of level lands that can be distinguished meet the same criteria, although they differ in their relationship towards the surrounding land. As the upper slope limit for level land is a gradient of 8%, areas with a perceptible slope may still be considered level land.

#### **LP Plains**

Plains are all level lands that are not enclosed between higher lying lands, that do not protrude above the surrounding country, or that do not rise gently against land with a considerable steeper slope.

#### **LL Plateaux**

Plateaux are level lands that are, compared with the surrounding landscapes, situated at relatively elevated positions. Plateau can be very extensive, but must always on at least one side be bounded by a slope or escarpment (8% or more), connecting it with lower lying land. Many so-called plateaux are in fact elevated plains, and should be classified as such.

#### **LD Depressions**

A depression is an area of level land that is on all sides surrounded by higher lying level or sloping land. The area occupied by the band of sloping land that forms the transition from the higher ground to the floor of the depression is small compared to the area within the depression taken up by level land.

#### **LF Low-gradient footslopes**

Steadily rising level land, abutting strongly sloping or steep lands, are classified as low level footslopes. They merge into other types of level land, including low gradient footslopes that rise in an opposite direction. Pediments, (coalescing) alluvial fans and other similar landforms can all be considered low level footslopes. Footslopes with a higher gradient than 8% are accommodated under hills, as such slopes are usually incised to the extent that they take a hilly character.

#### **LV Valley floors**

Elongated strips of level land, on both sides flanked by areas with sloping or steep land, constitute valley floors. Valley floors normally taper off at one end, where they are embraced by steeper land on three sides. They may connect with other types of level land or sloping land at the other end. In mountainous areas valley floors can be surrounded on all sides by steep lands, and do not necessarily have to be elongated.

**S Sloping land**

Sloping land is land with a gradient of between 8 and 30%. In most cases the relief intensity of sloping land is more than 50 m per slope unit.

**SM Medium-gradient mountains**

Relatively gently sloping (15-30% gradient) mountains with a local relief intensity of more than 600 m. Many volcanoes will fall into this category, as do several foothill zones of major mountain systems.

**SH Medium-gradient hills**

All sloping land with an undulating relief (minimum relief intensity 50 m per slope unit), not elongated, or more than 600 m high, or incorporated in mountainous terrain, are considered hills. This group does not only include hilly landforms, but also accommodates other landforms such as medium-gradient footslopes, etc.

**SE Medium-gradient escarpment zone**

Relatively gently sloping (usually 15-30% gradient) zone that forms a transition between high and low lying country. The local relief intensity of this landform is normally less than 600 m/2 km.

**SR Ridges**

A ridge meets all the qualifications of medium-gradient hills, but has an elongated shape with a single crest, which may have a more or less constant elevation, or may contain a number of peaks. Relatively narrow plateaus are excluded from this landform group.

**SU Mountainous highland**

Land which, although forming part of a mountain range (slopes of more than 30% and relief intensities in excess of 600 m per 2 km), constitute a restricted zone with less steep slopes and subdued relief. Mountainous highland always forms part of a mountain system, and is thus on at least at one side bounded by high-gradient mountains. Hypsometric subdivision of this category is according to the qualifiers for steep lands.

**SP Dissected plains**

Sloping land with a more or less constant crest level, and relief intensities of less than 50 m per slope unit.

**T Steep land**

All land with slopes in excess of 30% is considered steep land. The main landform in this category is mountainous land.

**TM High-gradient mountains**

All steep land with a relief intensity of more than 600 m per 2 km, and surrounding one or more outstanding peaks.

**TH High-gradient hills**

Steep but low relief land (relief intensity of less than 600 m per 2 km). Badlands would be a landform taken care of by this group, which is hypsometrically subdivided according to the qualifiers for sloping land.

**TE High-gradient escarpment zone**

Steep land that forms the transition between high and low lying country and lacks outstanding peaks. The relief intensity is normally more than 600 m per 2 km.

**TV High-gradient valleys**

Very steep valleys, with normally very little valley floor. No height limit is given, as the lack of valley floor and the presence of steep slopes ensure that only deep valleys will cover sufficient area to produce mappable delineations. Mostly incised elevated sedimentary plateaux.

**C Lands with composite landforms**

Landforms, containing both level and steep or sloping land, which cannot be separated at the scale of the mapping, are considered composite landforms. Composite landforms are using hypsometric qualifiers according to the characteristics of their level part.

**CV Valleys**

The valley, made up of sideslopes and a valley bottom, is taken as one landform.

**CL Narrow plateaus**

A narrow strip of level land surrounded on all sides by sloping or steep falling land form together a narrow plateau.

**CD Major depressions**

A large tract of level land, surrounded on all sides by high, rising sloping or

steep land, is characterized as a major depression. Uvalas are typical for this group.



## Annex 2

FAO soil unit codes<sup>1</sup>**FL** FLUVISOLS

<b>FLe</b>	Eutric Fluvisols
<b>FLc</b>	Calcaric Fluvisols
<b>FLd</b>	Dystric Fluvisols
<b>FLm</b>	Mollic Fluvisols
<b>Flu</b>	Umbric Fluvisols
<b>FLt</b>	Thionic Fluvisols
<b>FLs</b>	Salic Fluvisols

**LP** LEPTOSOLS

<b>LPe</b>	Eutric Leptosols
<b>LPd</b>	Dystric Leptosols
<b>LPk</b>	Rendzic Leptosols
<b>LPm</b>	Mollic Leptosols
<b>LPu</b>	Umbric Leptosols
<b>LPq</b>	Lithic Leptosols
<b>LPi</b>	Gelic Leptosols

**GL** GLEYSOLS

<b>GLe</b>	Eutric Gleysols
<b>GLk</b>	Calcic Gleysols
<b>GLd</b>	Dystric Gleysols
<b>GLa</b>	Andic Gleysols
<b>GLm</b>	Mollic Gleysols
<b>GLu</b>	Umbric Gleysols
<b>GLt</b>	Thionic Gleysols
<b>GLi</b>	Gelic Gleysols

**AR** ARENOSOLS

<b>ARh</b>	Haplic Arenosols
<b>ARb</b>	Cambic Arenosols
<b>ARi</b>	Luvic Arenosols
<b>ARo</b>	Ferralic Arenosols
<b>Ara</b>	Albic Arenosols
<b>ARc</b>	Calcaric Arenosols
<b>ARg</b>	Gleyic Arenosols

**RG** REGOSOLS

<b>RGe</b>	Eutric Regosols
<b>RGc</b>	Calcaric Regosols
<b>RGy</b>	Gypsic Regosols
<b>RGd</b>	Dystric Regosols
<b>RGu</b>	Umbric Regosols
<b>Rgi</b>	Gelic Regosols

**AN** ANDOSOLS

<b>ANh</b>	Haplic Andosols
<b>ANm</b>	Mollic Andosols
<b>ANu</b>	Umbric Andosols
<b>ANz</b>	Vitric Andosols
<b>ANg</b>	Gleyic Andosols
<b>ANi</b>	Gelic Andosols

---

<sup>1</sup> After FAO (1988).

**VR** VERTISOLS

<b>VRe</b>	Eutric Vertisols
<b>VRd</b>	Dystric Vertisols
<b>VRk</b>	Calcic Vertisols
<b>VRy</b>	Gypsic Vertisols

**CM** CAMBISOLS

<b>CMe</b>	Eutric Cambisols
<b>CMd</b>	Dystric Cambisols
<b>CMu</b>	Humic Cambisols
<b>CMc</b>	Calcaric Cambisols
<b>CMx</b>	Chromic Cambisols
<b>CMv</b>	Vertic Cambisols
<b>CMo</b>	Ferralic Cambisols
<b>CMg</b>	Gleyic Cambisols
<b>CMi</b>	Gelic Cambisols

**CL** CALCISOLS

<b>CLh</b>	Haplic Calcisols
<b>CLl</b>	Luvic Calcisols
<b>CLp</b>	Petric Calcisols

**GY** GYPSISOLS

<b>GYh</b>	Haplic Gypsisols
<b>GYk</b>	Calcic Gypsisols
<b>GYl</b>	Luvic Gypsisols
<b>GYp</b>	Petric Gypsisols

**SN** SOLONETZ

<b>SNh</b>	Haplic Solonetz
<b>SNm</b>	Mollic Solonetz
<b>SNk</b>	Calcic Solonetz
<b>SNy</b>	Gypsic Solonetz
<b>SNj</b>	Stagnic Solonetz
<b>SNg</b>	Gleyic Solonetz

**SC** SOLONCHAKS

<b>SCh</b>	Haplic Solonchaks
<b>SCm</b>	Mollic Solonchaks
<b>SCK</b>	Calcic Solonchaks
<b>Scy</b>	Gypsic Solonchaks
<b>SCn</b>	Sodic Solonchaks
<b>SCg</b>	Gleyic Solonchaks
<b>SCi</b>	Gelic Solonchaks

**KS** KASTANOZEMS

<b>KSh</b>	Haplic Kastanozems
<b>KSl</b>	Luvic Kastanozems
<b>KSk</b>	Calcic Kastanozems
<b>KSy</b>	Gypsic Kastanozems

**CH** CHERNOZEMS

<b>CHh</b>	Haplic Chernozems
<b>CHk</b>	Calcic Chernozems
<b>CHl</b>	Luvic Chernozems
<b>CHw</b>	Glossic Chernozems
<b>CHg</b>	Gleyic Chernozems

**PH** PHAEOZEMS

<b>PHh</b>	Haplic Phaeozems
<b>PHc</b>	Calcaric Phaeozems
<b>PHl</b>	Luvic Phaeozems
<b>PHj</b>	Stagnic Phaeozems
<b>PHg</b>	Gleyic Phaeozems

**GR** GREYZEMS

<b>GRh</b>	Haplic Greyzems
<b>GRg</b>	Gleyic Greyzems



**LXg** Gleyic Lixisols**LV** LUVISOLS

<b>LVh</b>	Haplic Luvisols
<b>LVf</b>	Ferric Luvisols
<b>LVx</b>	Chromic Luvisols
<b>LVk</b>	Calcic Luvisols
<b>LVv</b>	Vertic Luvisols
<b>LVa</b>	Albic Luvisols
<b>LVj</b>	Stagnic Luvisols
<b>LVg</b>	Gleyic Luvisols

**AC** ACRISOLS

<b>ACH</b>	Haplic Acrisols
<b>ACf</b>	Ferric Acrisols
<b>ACu</b>	Humic Acrisols
<b>ACp</b>	Plinthic Acrisols
<b>ACg</b>	Gleyic Acrisols

**PL** PLANOSOLS

<b>PLe</b>	Eutric Planosols
<b>PLd</b>	Dystric Planosols
<b>PLm</b>	Mollic Planosols
<b>PLu</b>	Umbric Planosols
<b>Pli</b>	Gelic Planosols

**AL** ALISOLS

<b>ALh</b>	Haplic Alisols
<b>ALf</b>	Ferric Alisols
<b>ALu</b>	Humic Alisols
<b>ALp</b>	Plinthic Alisols
<b>Alj</b>	Stagnic Alisols
<b>ALg</b>	Gleyic Alisols

**PD** PODZOLUVISOLS

<b>PDe</b>	Eutric Podzoluvisols
<b>PDD</b>	Dystric Podzoluvisols
<b>PDj</b>	Stagnic Podzoluvisols
<b>PDg</b>	Gleyic Podzoluvisols
<b>PDi</b>	Gelic Podzoluvisols

**NT** NITISOLS

<b>NTh</b>	Haplic Nitisols
<b>NTr</b>	Rhodic Nitisols
<b>NTu</b>	Humic Nitisols

**PZ** PODZOLS

<b>PZh</b>	Haplic Podzols
<b>PZb</b>	Cambic Podzols
<b>PZf</b>	Ferric Podzols
<b>PZc</b>	Carbic Podzols
<b>PZg</b>	Gleyic Podzols
<b>PZi</b>	Gelic Podzols

**FR** FERRALSOLS

<b>FRh</b>	Haplic Ferralsols
<b>FRx</b>	Xanthic Ferralsols
<b>FRr</b>	Rhodic Ferralsols
<b>FRu</b>	Humic Ferralsols
<b>FRg</b>	Geric Ferralsols
<b>FRp</b>	Plinthic Ferralsols

**LX** LIXISOLS

<b>LXh</b>	Haplic Lixisols
<b>LXf</b>	Ferric Lixisols
<b>LXp</b>	Plinthic Lixisols
<b>LXa</b>	Albic Lixisols
<b>LXj</b>	Stagnic Lixisols

**PT** PLINTHOSOLS

<b>PTe</b>	Eutric Plinthosols
<b>PTd</b>	Dystric Plinthosols
<b>PTu</b>	Humic Plinthosols
<b>PTa</b>	Albic Plinthosols

---

		<b>AT</b>	ANTHROSOLS
		<b>ATa</b>	Aric Anthrosols
		<b>ATc</b>	Cumulic Anthrosols
		<b>ATf</b>	Fimic Anthrosols
		<b>ATu</b>	Urbic Anthrosols
<b>HS</b>	HISTOSOLS		
<b>HSI</b>	Folic Histosols		
<b>HSs</b>	Terric Histosols		
<b>HSf</b>	Fibric Histosols		
<b>HSt</b>	Thionic Histosols		
<b>HSi</b>	Gelic Histosols		

## Annex 3

### Hierarchy of land use<sup>1</sup>

**S Settlement/industries:** Residential, industrial use.

**SR** Residential use: Cities.

**SI** Industrial use: Industries.

**ST** Transport: Roads, railways etc.

**SC** Recreation: In use for recreation.

**SX** Excavations: Land used for excavations, quarries.

**A Agriculture:** Land used for cultivation of crops.

**AA** Annual field cropping: One or more crops harvested within one year. Land under temporary crops.

**AA1** Shifting cultivation: Agricultural systems that involve an alternation between cropping for a few years on selected and cleared plots and a lengthy period when the soil is rested. The land is cultivated for less than 33% of the years.

**AA2** Fallow system cultivation: Agricultural systems that involve an alternation of cropping periods and fallow periods. The land is cultivated between 33 and 67% of the growing seasons; bush or grass fallow are typical.

**AA3** Ley system cultivation: Several years of arable cropping are followed by several years of grass and legumes utilized for livestock production.

**AA4** Rainfed arable cultivation: Agricultural systems where the land is cultivated in more than 67% of the growing seasons.

**AA5** Wet rice cultivation: Annual field cropping system for the production of wetland rice. Paddies with or without controlled water supply and drainage system. Plots are inundated during at least some part of the cropping period.

---

<sup>1</sup> Adapted from Remmelzwaal (1990).

**AA6** Irrigated cultivation: Annual field cropping system with an artificial supply of water, in addition to rain.

**AP** Perennial field cropping: Land under perennial crops. Crops harvested more than one year after planting. Examples of perennial field crops are sugar-cane, bananas, pineapples and sisal.

**AP1** Non-irrigated cultivation

**AP2** Irrigated cultivation

**AT** Tree & shrub cropping: Crops harvested annually or perennially; trees or shrubs produce more than one crop. Examples of tree crops are oil-palm, rubber, cacao, coconuts and cloves; typical shrub crops are coffee and tea.

**AT1** Non-irrigated tree crop cultivation

**AT2** Irrigated tree crop cultivation

**AT3** Non-irrigated shrub crop cultivation

**AT4** Irrigated shrub crop cultivation

**H** **Animal husbandry:** Animal products.

**HE** Extensive grazing: Grazing on natural or semi-natural grassland or savanna vegetation.

**HE1** Nomadism: Systems in which the animal owners do not have a permanent place of residence. No regular cultivation practices. People move with herds.

**HE2** Semi-nomadism: Animal owners have a permanent place of residence where supplementary cultivation is practised. Herds are moved to distant grazing areas.

**HE3** Ranching: Grazing within well defined boundaries, movements less distant and higher management level as compared to semi-nomadism.

**HI** Intensive grazing: Stationary animal husbandry. Grazing on permanent/semi-permanent improved grassland systems.

**HI1** Animal production

**HI2** Dairying

**F** **Forestry:** Activities related to the production of wood. Exploitation of forest for wood, with reforestation. A commercial activity.

**FN** Exploitation of natural forest and woodland: Wood is extracted from natural forest and woodland for commercial purpose.

- FN1** selective felling: Only selected species are removed from the natural vegetation.
- FN2** clear felling: All natural vegetation is cleared after which the area is reforested. This land use system develops into a plantation forestry system.
- FP** Plantation forestry: Forested areas. Relatively high management level. Homogeneous tree stands.
- M** **Mixed farming:** Activities concerning cropping and forestry or animal husbandry are mixed.
- MF** Agro-forestry: Combination of agriculture and forestry (with reforestation).
- MP** Agro-pastoralism: Combination of agriculture and animal husbandry, also called transhumance (farmers with a permanent place of residence send their herds, tended by herdsman, for long periods of time to distant grazing areas).
- E** **Extraction/collecting:** Extraction of products from the environment.
- EV** exploitation of natural vegetation: Land used for extraction of wood or other products from the vegetation; for domestic use.
- EH** hunting and fishing: Extraction of animals or fish from ecosystem.
- P** **Nature protection:** No, or low intensity of use, but under management system; low level of interference with natural environment or ecosystem.
- PN** Nature and game preservation
- PN1** Reserves
- PN2** Parks
- PN3** Wildlife management
- PD** Degradation control: Degradation of land, in most cases further degradation, is not desirable and the land is protected.
- PD1** Non-interference: All uses of the land are prohibited.
- PD2** Interference: The land is managed. Works are implemented in order to stop degradation and limit the degradation risk.
- U** **Unused:** Not used and not managed.



## Annex 4

# Hierarchy of vegetation<sup>1</sup>

**I Closed forest:** Formed by trees at least 5 m tall with their crowns interlocking.

**IA** Mainly evergreen forest: The canopy is never without green foliage. However, individual trees may shed their leaves for that period.

**IA1** Tropical ombrophilous forest (tropical rain forest): Consisting mainly of broad-leaved evergreen trees, neither cold nor drought resistant. Truly evergreen, i.e. the forest canopy remains green all year though individual trees may be leafless for a few weeks.

**IA2** Tropical and subtropical evergreen seasonal forest: Consisting mainly of broad-leaved evergreen trees. Foliage reduction during the dry season noticeable, often as partial shedding of leaves.

**IA3** Tropical and subtropical semi-deciduous forest: Most of the upper canopy trees deciduous or drought-resistant; many of the understorey trees and shrubs evergreen and more or less sclerophyllous<sup>2</sup>.

**IA4** Subtropical ombrophilous forest: Forest with a dry season and more pronounced temperature differences between summer and winter than tropical ombrophilous forest.

**IA5** Mangrove forest: Composed almost entirely of evergreen sclerophyllous broad-leaved trees/shrubs with either stilt roots or pneumatophores.

**IA6** Temperate and subpolar evergreen ombrophilous forest: Consisting mostly of truly evergreen hemi-sclerophyllous trees and shrubs. Rich in epiphytes and herbaceous ferns.

**IA7** Temperate evergreen seasonal broad-leaved forest: Consisting mainly of hemi-sclerophyllous evergreen trees and shrubs, rich in herbaceous undergrowth.

---

<sup>1</sup> After UNESCO (1973).

<sup>2</sup> Sclerophyllous: thick, hard leaves.

- IA8** Winter-rain evergreen broad-leaved sclerophyllous forest (Mediterranean forest): Consisting mainly of sclerophyllous evergreen trees and shrubs, most of them showing rough bark. Herbaceous undergrowth almost lacking.
- IA9** Tropical and subtropical evergreen needle-leaved forest: Consisting mainly of needle-leaved evergreen trees. Broad-leaved trees may be present.
- IA10** Temperate and subpolar evergreen needle-leaved forest: Consisting mainly of needle-leaved or scale-leaved evergreen trees, but broad-leaved trees may be admixed.
- IB** Mainly deciduous forest: Majority of trees shed their foliage simultaneously in connection with the unfavourable season.
  - IB1** Tropical and subtropical drought-deciduous forest: Unfavourable season mainly characterized by drought, in most cases winter-drought. Foliage is shed regularly every year. Most trees with relatively thick, fissured bark.
  - IB2** Cold-deciduous forest with evergreen trees (or shrubs): Unfavourable season mainly characterized by winter frost. Deciduous broad-leaved trees dominant, but evergreen species present.
  - IB3** Cold-deciduous forest without evergreen trees: Deciduous trees absolutely dominant.
- IC** Extremely xeromorphic forest: Dense stand of xeromorphic phanerophytes such as bottle trees, tuft trees with succulent leaves and stem succulents. Undergrowth with shrubs of similar xeromorphic adaptations.
  - IC1** Sclerophyllous-dominated extremely xeromorphic forest: Predominance of sclerophyllous trees.
  - IC2** Thorn forest: Species with thorny appendices predominate.
  - IC3** Mainly succulent forest: Tree-formed and shrub-formed succulents
- II** **Woodland:** Composed of trees at least 5 m tall with crowns not usually touching but with a coverage of at least 40%.
  - IIA** Mainly evergreen woodland: The canopy is never without green foliage.
    - IIA1** Evergreen broad-leaved woodland: Mainly sclerophyllous trees and shrubs.
    - IIA2** Evergreen needle-leaved forest: Mainly needle-leaved or scale-leaved.



- IIB** Mainly deciduous woodland: Majority of trees shed their foliage simultaneously in connection with the unfavourable season.
  - IIB1** Drought deciduous woodland: Unfavourable season mainly characterized by winter-drought. Foliage is shed regularly every year. Most trees with relatively thick, fissured bark.
  - IIB2** Cold-deciduous woodland with evergreen trees: Unfavourable season mainly characterized by winter frost. Deciduous broad-leaved trees dominant, but evergreen species present.
  - IIB3** Cold-deciduous woodland without evergreen trees: Deciduous trees absolutely dominant.
- IIC** Extremely xeromorphic woodland: Open stand of xeromorphic phanerophytes such as bottle trees, tuft trees with succulent leaves and stem succulents. Undergrowth with shrubs of similar xeromorphic adaptations.
  - IIC1** Sclerophyllous-dominated extremely xeromorphic woodland: Predominance of sclerophyllous trees.
  - IIC2** Thorn woodland: Species with thorny appendices predominate.
  - IIC3** Mainly succulent woodland: Tree-formed and shrub-formed succulents.
- III Scrub** (shrubland or thicket): Mainly composed of woody plants 0.5 to 5 m tall. Subdivisions: Shrubland: most of the individual shrubs not touching each other; often grass undergrowth; Thicket: individual shrubs interlocked.
  - IIIA** Mainly evergreen scrub: The canopy is never without green foliage. However, individual shrubs may shed their leaves.
    - IIIA1** Evergreen broad-leaved shrubland (or thicket): Mainly sclerophyllous shrubs.
    - IIIA2** Evergreen needle-leaved and microphyllous shrubland (or thicket): Mainly needle-leaved or scale-leaved shrubs.
  - IIIB** Mainly deciduous scrub: Majority of shrubs shed their foliage simultaneously in connection with the unfavourable season.
    - IIIB1** Drought-deciduous scrub with evergreen woody plants admixed
    - IIIB2** Drought-deciduous scrub without evergreen woody plants admixed
    - IIIB3** Cold-deciduous scrub

- IIIC** Extremely xeromorphic (subdesert) shrubland: Very open stands of shrubs with various xerophytic adaptations, such as extremely scleromorphic or strongly reduced leaves, green branches without leaves, or succulents stems, etc., some of them with thorns.
- IIIC1** Mainly evergreen subdesert shrubland: In extremely dry years some leaves and shoot portions may be shed.
- IIIC2** Deciduous subdesert shrubland: Mainly deciduous shrubs, often with a few evergreens.
- IV Dwarf-scrub and related communities:** Rarely exceeding 50 cm in height. Subdivisions: Dwarf-scrub thicket: branches interlocked; Dwarf-shrubland: individual dwarf-shrubs more or less isolated or in clumps.
- IVA** Mainly evergreen dwarf-scrub: Most dwarf-scrubs evergreen.
- IVA1** Evergreen dwarf-scrub thicket: Densely closed dwarf-scrub cover, dominating the landscape.
- IVA2** Evergreen dwarf-shrubland: Open or more loose cover of dwarf-shrubs.
- IVA3** Mixed evergreen dwarf-shrub and herbaceous formation
- IVB** Mainly deciduous dwarf-scrub: Most dwarf-scrubs deciduous.
- IVB1** Facultatively drought-deciduous dwarf-thicket (or dwarf-shrubland): Foliage is shed only in extreme years.
- IVB2** Obligatory, drought-deciduous dwarf-thicket (or dwarf-shrubland): Densely closed dwarf-shrub stands which lose all or at least part of their leaves in the dry season.
- IVB3** Cold-deciduous dwarf-thicket (or dwarf-shrubland): Densely closed dwarf-shrub stands which lose all or at least part of their leaves at the beginning of a cold season.
- IVC** Extremely xeromorphic dwarf-shrubland: More or less open formations of dwarf-shrubs, succulents and other life forms adapted to survive or to avoid a long dry season. Mostly subdesertic.
- IVC1** Mainly evergreen subdesert dwarf-shrubland: In extremely dry years some leaves and shoot portions may be shed.
- IVC2** Deciduous subdesert dwarf-shrubland: Mainly deciduous dwarf-shrubs, often with a few evergreens.
- IVD** Tundra: Slowly growing, low formations, consisting mainly of dwarf-shrubs and

graminoids beyond the subpolar tree line.

**IVD1** Mainly bryophyte tundra: Dominated by mats or small cushions of mosses (bryophytes).

**IVD2** Mainly lichen tundra: Mats of lichen dominating.

**IVE** Mossy bog formations with dwarf-shrub: Oligotrophic peat accumulations formed by *Sphagnum* or other mosses.

**IVE1** Raised bog: By growth of *Sphagnum* species raised above the general groundwater table.

**IVE2** Non-raised bog: Not or not very markedly raised above the mineral-water table of the surrounding landscape.

## **V Herbaceous vegetation**

**VA** Tall graminoid vegetation: Dominant graminoids over 2 m tall. Forb<sup>1</sup> coverage less than 50%.

**VA1** Tall grassland with a tree synusia<sup>2</sup> covering 10-40%: More or less like a very open woodland.

**VA2** Tall grassland with a tree synusia covering less than 10%

**VA3** Tall grassland with a synusia of shrubs

**VA4** Tall grassland with a woody synusia consisting mainly of tuft plants (usually palms)

**VA5** Tall grassland practically without woody synusia

**VB** Medium tall grassland: The dominant graminoid growth forms are 50 cm to 2 m tall. Forbs cover less than 50%.

**VB1** Medium tall grassland with a tree synusia covering 10-40%

**VB2** Medium tall grassland with a tree synusia covering less than 10%

**VB3** Medium tall grassland with a synusia of shrubs

**VB4** Medium tall grassland with an open synusia of tuft plants (usually palms)

**VB5** Medium tall grassland practically without woody synusia

---

<sup>1</sup> Forb: non-graminoid/non-woody vegetation.

<sup>2</sup> Synusia: layer.

- VC** Short grassland: The dominant graminoid growth forms are less than 50 cm tall.  
Forbs cover less than 50%.
  - VC1** Short grassland with a tree synusia covering 10-40%
  - VC2** Short grassland with a tree synusia covering less than 10%
  - VC3** Short grassland with a synusia of shrubs
  - VC4** Short grassland with an open synusia of tuft plants (usually palms)
  - VC5** Short grassland practically without woody synusia
  - VC6** Short to medium tall mesophytic grassland
  - VC7** Graminoid tundra
- VD** Forb vegetation: Mainly forbs, graminoid cover less than 50%.
  - VD1** Tall forb communities: Dominant forb growth forms are more than 1 m tall.
  - VD2** Low forb communities: Dominant forb growth forms are less than 1 m tall.
- VE** Hydromorphic fresh-water vegetation
  - VE1** Rooted fresh-water communities
  - VE2** Free floating fresh-water communities

## Annex 5

## ISO country codes

Country codes according to ISO-3166 of 1994.

AF	Afghanistan	CG	Congo	HU	Hungary
AL	Albania	CK	Cook Islands	IS	Iceland
DZ	Algeria	CR	Costa Rica	IN	India
AS	American Samoa	CI	Côte d'Ivoire	ID	Indonesia
AD	Andorra	HR	Croatia	IR	Iran, Islamic Republic of
AO	Angola	CU	Cuba	IQ	Iraq
AI	Anguilla	CY	Cyprus	IE	Ireland
AQ	Antarctica	CZ	Czech Republic	IL	Israel
AG	Antigua and Barbuda	CS	Czechoslovakia	IT	Italy
AR	Argentina	DK	Denmark	JM	Jamaica
AM	Armenia	DJ	Djibouti	JP	Japan
AW	Aruba	DM	Dominica	JO	Jordan
AU	Australia	DO	Dominican Republic	KZ	Kazakhstan
AT	Austria	TP	East Timor	KE	Kenya
AZ	Azerbaijan	EC	Ecuador	KI	Kiribati
BS	Bahamas	EG	Egypt	KR	Korea, Republic of
BH	Bahrain	SV	El Salvador	KP	Korea, Dem. Peopl. Rep.
BD	Bangladesh	GQ	Equatorial Guinea	KW	Kuwait
BB	Barbados	ER	Eritrea	KG	Kyrgystan
BY	Belarus	EE	Estonia	LA	Lao, People's Democratic Rep.
BE	Belgium	ET	Ethiopia	LV	Latvia
BZ	Belize	FK	Falkland Islands	LB	Lebanon
BJ	Benin	FO	Faroe Islands	LS	Lesotho
BT	Bhutan	FJ	Fiji	LR	Liberia
BO	Bolivia	FI	Finland	LY	Libyan Arab Jamahiri
BA	Bosnia and Herzegovina	FR	France	LI	Liechtenstein
BW	Botswana	GF	French Guiana	LT	Lithuania
BV	Bouvet Island	PF	French Polynesia	LU	Luxembourg
BR	Brazil	TF	French Southern Territories	MO	Macao
IO	British Indian Ocean Territory	GA	Gabon	MG	Madagascar
BN	Brunei Darussalam	GM	Gambia	MW	Malawi
BG	Bulgaria	GE	Georgia	MY	Malaysia
BF	Burkina Faso	DE	Germany	MV	Maldives
BU	Burma	GH	Ghana	ML	Mali
BI	Burundi	GI	Gibraltar	MT	Malta
KH	Cambodia	GR	Greece	MH	Marshall Islands
CM	Cameroon	GL	Greenland	MQ	Martinique
CA	Canada	GD	Grenada	MR	Mauritania
CV	Cape Verde	GP	Guadeloupe	MU	Mauritius
KY	Cayman Islands	GU	Guam	MX	Mexico
CF	Central African Republic	GT	Guatemala	FM	Micronesia
TD	Chad	GN	Guinea	MD	Moldova, Republic of
CL	Chile	GW	Guinea-Bissau	MC	Monaco
CN	China	GY	Guyana	MN	Mongolia
CX	Christmas Island	HT	Haiti	MS	Montserrat
CC	Cocos Islands	HM	Heard and McDonald Islands	MA	Morocco
CO	Colombia	HN	Honduras	MZ	Mozambique
		HK	Hong Kong		

MM	Myanmar	TO	Tonga
NA	Namibia	TT	Trinidad and Tobago
NR	Nauru	TN	Tunisia
NP	Nepal	TR	Turkey
NL	Netherlands	TM	Turkmenistan
AN	Netherlands Antilles	TC	Turks and Caicos Islands
NT	Neutral Zone	TV	Tuvalu
NC	New Caledonia	UM	US, Minor Outlying Islands
NZ	New Zealand	UG	Uganda
NI	Nicaragua	UA	Ukraine
NE	Niger	AE	United Arab Emirates
NG	Nigeria	GB	United Kingdom
NU	Niue	US	United States
NF	Norfolk Island	UY	Uruguay
MP	Northern Mariana Islands	UZ	Uzbekistan
NO	Norway	SU	USSR
OM	Oman	VU	Vanuatu
PK	Pakistan	VA	Vatican City State
PW	Palau	VE	Venezuela
PA	Panama	VN	Viet Nam
PG	Papua New Guinea	VG	Virgin Islands (U.K.)
PY	Paraguay	VI	Virgin Islands (U.S.)
PE	Peru	WF	Wallis and Futuna Islands
PH	Philippines	EH	Western Sahara
PN	Pitcairn	YE	Yemen
PL	Poland	YU	Yugoslavia
PT	Portugal	ZR	Zaire
PR	Puerto Rico	ZM	Zambia
QA	Qatar	ZW	Zimbabwe
RE	Reunion		
RO	Romania		
RU	Russian Federation		
RW	Rwanda		
LC	Saint Lucia		
WS	Samoa		
SM	San Marino		
ST	Sao Tome and Principe		
SA	Saudi Arabia		
SN	Senegal		
SC	Seychelles		
SL	Sierra Leone		
SG	Singapore		
SK	Slovakia		
SI	Slovenia		
SB	Solomon Islands		
SO	Somalia		
ZA	South Africa		
ES	Spain		
LK	Sri Lanka		
SH	St. Helena		
KN	St. Kitts and Nevis		
PM	St. Pierre and Miquelon		
VC	St. Vincent and the Grenadines		
SD	Sudan		
SR	Suriname		
SJ	Svalbard and Jan Mayen		
SZ	Swaziland		
SE	Sweden		
CH	Switzerland		
SY	Syrian Arab Republic		
TW	Taiwan, Province of China		
TJ	Tajikistan		
TZ	Tanzania, United Republic of		
TH	Thailand		
TG	Togo		
TK	Tokelau		

## Annex 6

# **SOTER data entry forms**







form 3

## SOTER data entry

## SOIL COMPONENT

33 SOTER unit_ID	LLLLL
34 terrain component number	L
35 soil component number	L
36 proportion of SOTER unit	LLL
37 profile_ID	LLLLLLLLLLLLL
38 number of reference profiles	LLL
39 position in terrain component	L
40 surface rockiness	L
41 surface stoniness	L
42 types of erosion/deposition	L
43 area affected	L
44 degree of erosion	L
45 sensitivity to capping	L
46 rootable depth	L
47 relation with other soil components	

form 4

SOTER data entry

PROFILE	
48 profile_ID	LLLLLLLLLLLLLLLL
49 profile database_ID	LLLLL
50 latitude	LLLLLLL
51 longitude	LLLLLLLLL
52 elevation	LLLLL
53 sampling date	LLLLL
54 lab_ID	LLLLLL
55 drainage	L
56 infiltration rate	L
57 surface organic matter	L
58 classification FAO	LLLL
59 classification version	LLLL
60 national classification	LLLLLLLLLLLLL
61 Soil Taxonomy	LLLLLLLLLLLLL
62 phase	LL
HORIZON representative profile (* = mandatory)	
64 horizon number*	L
65 diagnostic horizon*	LL
66 diagnostic property*	LL
67 horizon designation	LLLLL
68 lower depth*	LLL
69 distinctness transition	L
70 moist colour*	LLLLLLLLL
71 dry colour	LLLLLLLLL
72 grade of structure	L
73 size of structure elements	L
74 type of structure*	L

form 5

SOTER data entry

HORIZON representative profile (continued) (* = mandatory)		profile_ID
64 horizon number*	U	U
75 abundance coarse fragments*	U	U
76 size of coarse fragments	U	U
77 very coarse sand	UUU	UUU
78 coarse sand	UUU	UUU
79 medium sand	UUU	UUU
80 fine sand	UUU	UUU
81 very fine sand	UUU	UUU
82 total sand*	UUU	UUU
83 silt*	UUU	UUU
84 clay*	UUU	UUU
85 particle size class	UUUU	UUUU
86 bulk density*	UUUU	UUUU
87 moisture content at various tensions	UUUU UUUU UUUU UUUU UUUU	UUUU UUUU UUUU UUUU UUUU
88 hydraulic conductivity	UUUU	UUUU
89 infiltration rate	UUUU	UUUU
90 pH H <sub>2</sub> O*	UUU	UUU
91 pH KCl	UUU	UUU

Note: Estimated mandatory (numeric) attributes should be preceded by the (-) sign.

SOTER data entry form 6

HORIZON	representative profile (continued) (* = mandatory)	profile_ID
64 horizon number*	U	U
92 electrical conductivity	UUU	UUU
93 soluble Na*	UUU	UUU
94 soluble Ca**	UUU	UUU
95 soluble Mg**	UUU	UUU
96 soluble K*	UUU	UUU
97 soluble Cl	UUU	UUU
98 soluble SO <sub>4</sub> **	UUU	UUU
99 soluble HCO <sub>3</sub>	UUU	UUU
100 soluble CO <sub>3</sub> .	UUU	UUU
101 exchangeable Ca**	UUU	UUU
102 exchangeable Mg**	UUU	UUU
103 exchangeable Na*	UUU	UUU
104 exchangeable K*	UUU	UUU
105 exchangeable Al***	UUU	UUU
106 exchangeable acidity	UUU	UUU
107 CEC soil*	UUUU	UUUU
108 total carbonate equivalent	UUU	UUU
109 gypsum	UUU	UUU
110 total carbon*	UUU	UUU

Note: Estimated mandatory (numeric) attributes should be preceded by the (-) sign.

form 7

SOTER data entry

HORIZON representative profile (continued) (* = mandatory)		profile_ID
64 horizon number*	LL	LL
111 total nitrogen	LLLLLL	LLLLLL
112 P <sub>2</sub> O <sub>5</sub>	LLLJ	LLLJ
113 phosphate retention	LLLJ	LLLJ
114 Fe dithionite	LLLJ	LLLJ
115 Al dithionite	LLLJ	LLLJ
116 Fe pyrophosphate	LLLJ	LLLJ
117 Al pyrophosphate	LLLJ	LLLJ
118 clay mineralogy	LLJ	LLJ

Note: Estimated mandatory (numeric) attributes should be preceded by the (.) sign.

form 8

SOTER data entry

HORIZON minimum values		profile_ID
64 horizon number	U	U
75 abundance coarse fragments	U	U
76 size of coarse fragments	U	U
77 very coarse sand	UUU	UUU
78 coarse sand	UUU	UUU
79 medium sand	UUU	UUU
80 fine sand	UUU	UUU
81 very fine sand	UUU	UUU
82 total sand	UUU	UUU
83 silt	UUU	UUU
84 clay	UUU	UUU
85 particle size class	UUUU	UUUU
86 bulk density	UUUU	UUUU
87 moisture content at various tensions	UUUU UUUU UUUU UUUU	UUUU UUUU UUUU UUUU
88 hydraulic conductivity	UUUU	UUUU
89 infiltration rate	UUUU	UUUU
90 pH H <sub>2</sub> O	UUU	UUU
91 pH KCl	UUU	UUU
92 electrical conductivity	UUU	UUU

form 9

SOTER data entry

HORIZON minimum values (continued)		profile_ID
64 horizon number	LJ	LJ
93 soluble Na*	LLLU	LLLU
94 soluble Ca**	LLLU	LLLU
95 soluble Mg**	LLLU	LLLU
96 soluble K*	LLLU	LLLU
97 soluble Cl	LLLU	LLLU
98 soluble SO <sub>4</sub> ***	LLLU	LLLU
99 soluble HCO <sub>3</sub>	LLLU	LLLU
100 soluble CO <sub>3</sub> .	LLLU	LLLU
101 exchangeable Ca**	LLLU	LLLU
102 exchangeable Mg**	LLLU	LLLU
103 exchangeable Na*	LLLU	LLLU
104 exchangeable K*	LLLU	LLLU
105 exchangeable Al***	LLLU	LLLU
106 exchangeable acidity	LLLU	LLLU
107 CEC soil*	LLLU	LLLU
108 total carbonate equivalent	LLLU	LLLU
109 gypsum	LLLU	LLLU
110 total carbon*	LLLU	LLLU
111 total nitrogen	LLLU	LLLU
112 P <sub>2</sub> O <sub>5</sub>	LLLU	LLLU





form 11

SOTER data entry

	HORIZON maximum values				profile_ID
64 horizon number	U	U	U	U	U
75 abundance coarse fragments	U	U	U	U	U
76 size of coarse fragments	U	U	U	U	U
77 very coarse sand	UUUU	UUUU	UUUU	UUUU	UUUU
78 coarse sand	UUUU	UUUU	UUUU	UUUU	UUUU
79 medium sand	UUUU	UUUU	UUUU	UUUU	UUUU
80 fine sand	UUUU	UUUU	UUUU	UUUU	UUUU
81 very fine sand	UUUU	UUUU	UUUU	UUUU	UUUU
82 total sand	UUUU	UUUU	UUUU	UUUU	UUUU
83 silt	UUUU	UUUU	UUUU	UUUU	UUUU
84 clay	UUUU	UUUU	UUUU	UUUU	UUUU
85 particle size class	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
86 bulk density	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
87 moisture content at various tensions	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
88 hydraulic conductivity	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
89 infiltration rate	UUUUU	UUUUU	UUUUU	UUUUU	UUUUU
90 pH H <sub>2</sub> O	UUUU	UUUU	UUUU	UUUU	UUUU
91 pH KCl	UUUU	UUUU	UUUU	UUUU	UUUU
92 electrical conductivity	UUUU	UUUU	UUUU	UUUU	UUUU

form 12

SOTER data entry

HORIZON maximum values (continued)		profile_ID
64 horizon number	U	UUUUUUUUUUUUUUUUUUUU
93 soluble Na <sup>+</sup>	UUUJ	UU UU
94 soluble Ca <sup>++</sup>	UUUJ	UUUU UU
95 soluble Mg <sup>++</sup>	UUUJ	UUUU UU
96 soluble K <sup>+</sup>	UUUJ	UUUU UU
97 soluble Cl <sup>-</sup>	UUUJ	UUUU UU
98 soluble SO <sub>4</sub>	UUUJ	UUUU UU
99 soluble HCO <sub>3</sub>	UUUJ	UUUU UU
100 soluble CO <sub>3</sub>	UUUJ	UUUU UU
101 exchangeable Ca <sup>++</sup>	UUUJ	UUUU UU
102 exchangeable Mg <sup>++</sup>	UUUJ	UUUU UU
103 exchangeable Na <sup>+</sup>	UUUJ	UUUU UU
104 exchangeable K <sup>+</sup>	UUUJ	UUUU UU
105 exchangeable Al <sup>+++</sup>	UUUJ	UUUU UU
106 exchangeable acidity	UUUJ	UUUU UU
107 CEC soil <sup>*</sup>	UUUJ	UUUU UU
108 total carbonate equivalent	UUUJ	UUUU UU
109 gypsum	UUUJ	UUUU UU
110 total carbon <sup>*</sup>	UUUJ	UUUU UU
111 total nitrogen	UUUJ	UUUU UU
112 P <sub>2</sub> O <sub>5</sub>	UUUJ	UUUU UU



SOTER data entry		form 14	
<b>LAND USE</b>		<b>VEGETATION</b>	
1 SOTER unit_ID	LLLLL	1 SOTER unit_ID	LLLLL
2 date of observation	LL/LL/LL	2 date of observation	LL/LL/LL
3 land use	LLL	3 vegetation	LLLLL
4 proportion of SOTER unit	LLL	4 proportion of SOTER unit	LLL

form 15

SOTER data entry

SOURCE MAP

1 map_ID	LLLLLLLLLLLLLLLL
2 map title	
3 year	LLLL
4 scale	1:LLLLLLLL
5 minimum latitude	LLLLLLLL
6 minimum longitude	LLLLLLLL
7 maximum latitude	LLLLLLLL
8 maximum longitude	LLLLLLLL
9 type of map	L

form 15

SOTER data entry

SOURCE MAP

1 map_ID	LLLLLLLLLLLLLLLL
2 map title	
3 year	LLLL
4 scale	1:LLLLLLLL
5 minimum latitude	LLLL.LLL
6 minimum longitude	LLLL.LLL
7 maximum latitude	LLLL.LLL
8 maximum longitude	LLLL.LLL
9 type of map	L

## form 17

## SOTER data entry

## CLIMATE STATION

1 climate station\_ID     |          |  
 2 climate station name   |          |  
 3 latitude                |          |  
 4 longitude               |          |  
 5 altitude                |          |

## CLIMATE DATA

6 climate station_ID		12 January		18 July	
7 kind of data		13 February		19 August	
8 source_ID		14 March		20 September	
9 first year		15 April		21 October	
10 last year		16 May		22 November	
11 years		17 June		23 December	
				24 annual	

## DATA SOURCE

25 source\_ID             |          |  
 26 source name



## References

- BAI. 1991. Booker tropical soil manual. Booker/Tate, London.
- Christian, C.S. and Stewart, G.A. 1953. General report on survey of Katharina-Darwin region, 1946. CSIRO Austr. Land Res. Ser. No. 1.
- Cochrane, T.T., de Castro, N.F. and Netto, J.M. 1981. An explanatory manual for CIAT's computerized land resource study of tropical America. CIAT, Cali.
- Cochrane, T.T., Sanchez, L.G., de Azevedo, L.G., Porras, J.H., and Garves, C.L. 1985. Land in tropical America. CIAT, Cali.
- Day, J.H. (ed.). 1983. The Canada soil information system (CanSIS), Manual for describing soils in the field. Agriculture Canada, Research Branch, Ottawa.
- DMA. 1992. Digital chart of the world. U.S. Defence Mapping Agency.
- Eschweiler, J.A. 1993. A draft physiographic map of Africa (at scale of 1:5 million). FAO internal public., Rome, p.46.
- FAO. 1977. Guidelines for the description of soils. FAO, Rome.
- FAO. 1986. Guidelines for the coding of soil data. Proposals for an international soil data bank, reprint 1977. FAO, Rome.
- FAO. 1988. Soil map of the world, Revised legend. World Soil Resources Report 60, FAO, Rome.
- FAO. 1989. FAO-ISRIC soil database. World Soil Resources Report 64. FAO, Rome.
- FAO. 1990. Guidelines for soil description. 3rd Edition (Revised). FAO, Rome.
- FAO-UNESCO. 1974. Soil map of the world. Vol 1, legend. Unesco, Paris.
- Gunn, R.H., Beattie, J.A., Reid, R.E. and van de Graaff, R.H.M. 1990. Australian soil and land survey handbook. Inkata Press, Melbourne. 300 p.
- Holmes. 1968. Physical geology. John Wiley, New York.
- ISRIC. 1990a. SOTER Procedures manual for small-scale map and database compilation. 3rd version. V.W.P. van Engelen and J.H.M. Pulles (eds.). Working paper and preprint 90/2, ISRIC, Wageningen, 69p.
- ISRIC. 1990b. Proceedings of the international workshop on procedures manual revisions for the Global Soils and Terrain Digital Database, Wageningen 24-26 April 1990. N.H. Batjes (ed.). Working paper and preprint 90/5, ISRIC, Wageningen, 25p.

- ISRIC. 1991. The SOTER manual. Procedures for small scale digital map and database compilation of soil and terrain conditions. V.W.P. van Engelen and J.H.M. Pulles (eds.). Working paper and preprint 91/3, ISRIC, Wageningen, 92p.
- ISRIC-UNEP, 1988. Guidelines for general assessment on the status of human-induced soil degradation. Working paper and preprint 88/4. ISRIC, Wageningen, 11p.
- ISSS. 1986a. Proceedings of an international workshop on the structure of a digital international soil resources map annex database. M.F. Baumgardner and L.R. Oldeman (eds.). SOTER Report 1, ISSS, Wageningen, 138p.
- ISSS. 1986b. Project proposal "World soils & terrain digital database at a scale 1:1 M (SOTER)". M.F. Baumgardner (ed.). ISSS, Wageningen, 23p.
- ISSS. 1987. Proceedings of the second international workshop on a global soils and terrain digital database. 18-22 May 1987, at UNEP Nairobi. R.F. van de Weg (ed.). SOTER Report 2, ISSS, Wageningen, 47p.
- ISSS. 1988a. International Reference Base Group. ISSS Commission V, Leuven.
- ISSS. 1988b. Proceedings of the first regional workshop on a global soils and terrain digital database and global assessment of soil degradation. 20-25 March 1988, Montevideo, Uruguay. W.L. Peters (ed.). SOTER Report 3, ISSS, Wageningen, 81p English/Spanish.
- ISSS. 1989. Proceedings of the second regional workshop on a global soils and terrain digital database and global assessment of soil degradation. 12-16 December 1988, Porto Alegre. W.L. Peters (ed.). SOTER Report 4, ISSS, Wageningen, 105p English/Spanish.
- Kenya Soil Survey Staff. 1987. Soil survey manual. Kenya Soil Survey, Nairobi.
- McDonald, R.C., Isbell, R.F., Speight, J.G., Walker, J. and Hopkins M.S. 1990. Australian soil and land survey. Field handbook. Second edition. Inkata Press, Melbourne. 198 p.
- Meijerink, A.M.J. and Valenzuela, C.R. 1987. ILWIS: Integrated Land and Watershed Management System. Methodology, terrain soil module, input. Publication M2, 2nd version. ITC, Enschede.
- National Bureau of Soil Survey and Land Use Planning. 1987. Field manual for soil mapping of India and of different states. Nagpur.
- Oliveira, J.B. and van de Berg, M. 1992. Application of the SOTER methodology for a semi-detailed survey (1:100,000) in the Piracicaba region (Sao Paulo, Brazil). SOTER Report 6. ISSS, Wageningen.
- Pulles, J.H.M. 1988. A model for a soils and terrain digital database. Working paper and preprint 88/8. ISRIC, Wageningen.
- Rommelzwaal, A. 1990. Classification of land and land use, first approach. FAO, Rome (unpublished).
- Rommelzwaal, A. 1991. Draft physiographic system. FAO, Rome (unpublished).
- Ruthenberg, H. 1980. Farming systems in the tropics. Third edition. Clarendon, Oxford.
- Shields, J.A. and Coote, D.R. 1988. SOTER Procedures manual for small-scale map and database

- compilation. Working paper and preprint 88/2. ISRIC, Wageningen, 144p.
- Shields, J.A. and Coote, D.R. 1989. SOTER Procedures manual for small-scale map and database compilation including proposed procedures for interpretation of soil degradation status and risk. Internal report, ISRIC, Wageningen.
- Soil Conservation Service. 1979. Pedon coding system for the national cooperative soil survey. US Dept. of Agric., Washington DC.
- Soil Conservation Service. 1981. Soil survey manual: revised chapter 4. US Dept. of Agric., Washington DC.
- Soil Conservation Service. 1986. National soils handbook, 1986 amendment. Glossary of landform and geologic terms. US Dept. of Agric., Washington DC.
- Soil Survey Staff. 1951. Soil survey manual. US Dept. of Agric. Handbook No. 18. Government Printer, Washington DC.
- Soil Survey Staff. 1975. Soil taxonomy. US Dept. of Agric. Handbook No. 438, Government Printer, Washington DC.
- Sombroek, W.G. 1984. Towards a global soil resource inventory at scale 1:1 M. Working paper and preprint 84/4, ISRIC, Wageningen.
- Strahler, A.N. 1969. Physical geology. John Wiley, New York.
- Tempel, P. 1994a. Global and national soils and terrain databases (SOTER). Attribute database user manual. Working paper and preprint 94/4, ISRIC, Wageningen, p. 34.
- Tempel, P. 1994b. Global and national soils and terrain databases (SOTER). Database structure. Working paper and preprint 94/5, ISRIC, Wageningen, p. 68.
- Touber, L., Smaling, E.M.A., Andriessse, W. and Hakkeling, R.T. 1989. Inventory and evaluation of tropical forest land. Guidelines for a common methodology. Tropenbos Foundation, Ede.
- UNESCO. 1973. International classification and mapping of vegetation. Ecology conservation 6. UNESCO, Paris.
- Van den Berg, M. 1992. SWEAP, a computer program for water erosion assessment applied to SOTER. SOTER Report 7. ISSS-UNEP-ISRIC, Wageningen. p. 37.
- Van de Weg, R.F. (ed.). 1987 Proceedings of the second international workshop on a global soils and terrain digital database. SOTER Report 2. ISSS, Wageningen.
- Van Waveren, E.J. and Bos, A.B. 1988a. Guidelines for the description and coding of soil data. Technical paper 14, ISRIC, Wageningen. 43p.
- Van Waveren, E.J. and Bos, A.B. 1988b. ISRIC Soil Information System; User manual, Technical manual. Technical paper 15, ISRIC, Wageningen. 63p.
- Verhasselt, Y. 1961. Essai de mise au point d'une mesure rapide de mesure de longueur et de surface. Bull. Soc. Royale Belge Géogr. Vol. 85, 1-4, pp. 131-144.
- Vogel, A.W. 1986. Class limits for land and soil properties. A comparative study for use at the establishment of a World Soil and Terrain Digital Database (SOTER). Working paper and

preprint 86/3, ISRIC, Wageningen. 154 p.

Wen, T.T. 1993. Draft physiographic map of South America. Internal public. FAO, Rome.

## Glossary

attribute data	Non-graphic information on elements in a GIS. In this manual: associated with SOTER units.
database	A computerized recordkeeping system.
database structure	The way in which data are organized in a database.
backup	A copy of a file or of a whole disk in case the original is lost/ damaged.
DBMS	Database Management System; a system for management and manipulation a database.
geo-referenced data	Information that has a precise location (coordinates).
GIS	Geographic(al) Information System = a system of hardware, software and procedures designated to support the capture, management, manipulation, analysis, modelling and display of spatially referenced data.
input	The process of entering data.
mapping unit	A set of areas (polygons) on a map that represent a well-defined feature or set of features; mapping units are described by the map legend.
polygon	Delineated area on a map
primary key	Attribute or combination of attributes that uniquely identify a record in a table/file.
RDBMS	Relational Database Management System; a computerized recordkeeping system in which the data are structured in sets of records so that relationships between data can be used for the management and manipulation. The data files are perceived as tables.
SOTER unit	Special type of mapping unit; a set of areas (polygons) on a map that have a distinctive, often repetitive pattern of landform, surface form, parent material and soil.

- topology            The way in which geographic elements are linked together (neighbouring elements, enclosed elements).
- uvala                depression in limestone terrain, generally elongated, comprising a series of joined dolinas or collapsed sinkholes.

## Related publications

- ISSS. 1986. Project proposal "World Soils and Terrain Digital Database at a scale 1:1M (SOTER)". M.F. Baumgardner (ed.). ISSS, Wageningen. 23 p.
- ISSS. 1986. Proceedings of an International Workshop on the Structure of a Digital International Soil Resources Map annex Data Base. (20-24 January 1986, ISRIC, Wageningen). M.F. Baumgardner and L.R. Oldeman (eds.). SOTER Report 1, ISSS, Wageningen. 138 p.
- ISSS. 1987. Proceedings of the Second International Workshop on a Global Soils and Terrain Digital Database (18-22 May 1987, UNEP, Nairobi). R.F. van de Weg (ed.). SOTER Report 2. ISSS, Wageningen. 47 p.
- ISSS. 1988. Proceedings of the First Regional Workshop on a Global Soils and Terrain Digital Database and Global Assessment of Soil Degradation. SOTER Report 3. ISSS, Wageningen. (81 p. in English, 86 p. in Spanish).
- ISSS. 1989. Proceedings of the Second Regional Workshop on a Global Soils and Terrain Digital Database (12-16 December 1988, Porto Alegre). W.L. Peters (ed.). SOTER Report 4. ISSS, Wageningen (97 p. in English, 104 p. in Spanish).
- ISRIC. 1990. Proceedings of the International Workshop on Procedures Manual Revisions for the Global Soils and Terrain Digital Database. 24-26 April 1990, Wageningen. N.H. Batjes (ed.). Working Paper & Preprint 90/5. ISRIC, Wageningen. 25 p.
- Batjes, N.H. 1990. Macro-scale land evaluation using the 1:1 M world soils and terrain digital database. SOTER Report 5. ISSS, Wageningen. 45 p.
- Oliveira, J.B. de and M. van den Berg. 1992. Application of the SOTER methodology to a semi-detailed survey (1:100,000) in the Piracicaba region (Sao Paulo State, Brazil). SOTER Report 6, ISSS, Wageningen. 28 p.
- Van den Berg, M. 1992. SWEAP, a computer program for water erosion assessment applied to SOTER. SOTER Report 7. ISSS-UNEP-ISRIC, Wageningen. 37. p.
- Tempel, P. 1994. Global and National Soils and Terrain Digital Databases (SOTER), Attribute Database User Manual. Working Paper & Preprint 94/04. ISRIC, Wageningen. p. 34.
- Tempel, P. 1994. Global and National Soils and Terrain Digital Databases (SOTER), Database Structure. Working Paper & Preprint 94/05. ISRIC, Wageningen. 68 p.





This publication describes the procedures for the Global and National **Soils and Terrain Digital Databases (SOTER)** to produce digitized map units and their attribute data. It explains how to delineate areas with a specific set of soil and terrain characteristics and to construct an attribute database related to the mapping units. SOTER is a land resource database with specific information on landform, terrain and soil components that can be complemented by data on land-related characteristics such as land use, natural vegetation and climate. The main function of the SOTER approach is to store data at national and global scales in an easily accessible format for improved thematic mapping and monitoring of changes of soil and terrain resources useful to scientists, planners, decision-makers and policy-makers.