

International Society for Environmental Information Sciences 2010 Annual Conference (ISEIS)

Biomonitoring and Bioindicators Used for River Ecosystems: Definitions, Approaches and Trends

Li Li, Binghui Zheng, Lusan Liu*

*Research Centre of River and Coastal Environmental, Institute of Water Environments, Chinese Research Academy of Environmental Sciences,
Beijing, 100012, PR China*

Abstract

In this paper, we present a review on concepts, current use and anticipated future directions of biomonitoring approaches and bioindicators used for river ecosystems. Periphyton, benthic macroinvertebrates and fish are the most common indicators in river biomonitoring, which can be used separately or contemporaneously. Their importances in the ecosystems and advantages for biomonitoring have been described in detail. Commonly used biomonitoring approaches include diversity, biotic indices, multimetric approaches, multivariate approaches, functional feeding groups (FFGs) and multiple biological traits. Among these techniques, biotic indices and multimetric approaches are most frequently used to evaluate the environment health of streams and rivers. However, functional measures have been increasingly applied as a complementary approach to reflecting ecological integrity. Furthermore, recent researches have demonstrated the efficiency of molecular techniques on enhancing the taxonomic resolutions and detecting the genetic diversity in river biomonitoring.

© 2009 Published by Elsevier Ltd.

Keywords: River Ecosystem; Biomonitoring; Bioindicators; Review

1. Introduction

1.1. Background

Since streams and rivers are among the most endangered ecosystems worldwide [1-3], there are urgent demands for comprehensive methodological approaches to evaluate the actual state of these ecosystems and to monitor their rate of changes [4]. Physical, chemical and bacteriological measurements commonly form the basis of monitoring,

* Corresponding author. Tel.: +86-10-8491-5316; fax: +86-10-8491-5302.
E-mail address: liuls@craes.org.cn.

because they provide complete spectrum of information for proper water management [5]. However, in running waters, where changes in hydrology are rapid and difficult to estimate, they cannot reflect the integration of numerous environment factors and long-term sustainability of river ecosystems for their instantaneous nature. Biomonitoring has been proven to be necessary supplementary to those traditional monitoring techniques [6]. Aquatic organisms, such as diatoms [7-10] and benthic macroinvertebrates [4-5], can serve as bioindicators to integrate their total environment and their responses to complex sets of environmental conditions [11]. They offer the possibility to obtain an ecological overview of the current status of streams or rivers.

1.2. Conceptual Issues

Biomonitoring, or biological monitoring, is generally defined as “the systematic use of living organisms or their responses to determine the condition or changes of the environment” [12-14]. Indeed, measurements (endpoints) used for river ecosystems may be selected from any level of biological organization (suborganismal, organismal, population, community, and ecosystem). However, the historical focus has been on ecological methods and higher levels of organization, e.g. populations, communities, and ecosystems. Therefore, the term of biomonitoring used in this paper tends to follow Markert et al. [15]: “Biomonitoring is a method of observing the impact of external factors on ecosystems and their development over a period, or of ascertaining differences between one location and another.” Compared to the former definition, the latter is considered to reflect the ecological content of biomonitoring better.

According to Markert et al. [15-17], a bioindicator is “an organism (or part of an organism or a community of organisms) that contains information on the quality of the environment (or a part of the environment)”. An “ideal” indicator at least should have the characteristics as follows: (a) taxonomic soundness (easy to be recognized by nonspecialist); (b) wide or cosmopolitan distribution; (c) low mobility (local indication); (d) well-known ecological characteristics; (e) Numerical abundance; (f) suitability for laboratory experiments; (g) high sensitivity to environmental stressor (s); (h) high ability for quantification and standardization [4, 18-19].

2. Bioindicators Used for River Ecosystems

Bioindicators need to not only indicate the long-term interaction of several environmental conditions, but also react to a sudden change of the important factor(s). There are several alternations for indicators of biomonitoring in streams and rivers, however benthic macroinvertebrates, periphytons and fishes are the most frequently utilized. Their efficacy when used separately has been demonstrated by many studies, e.g., Whitton and Rott [20], Vis et al. [21], Prygiel et al. [22], and Coste et al. [23] for periphyton, Rosenberg and Resh [4], Lenat and Barbour [24], Statzner et al. [25], and Buffagni et al. [26] for benthic macroinvertebrates, Fausch et al. [27], Joy and Death [28], Oberdorff et al. [29], and Pont et al. [30] for fish. In other studies, nevertheless, two or more assemblages have been used contemporaneously for monitoring river ecosystems, such as in Soinen and Könönen [6], Scuri et al. [31], Carlisle et al. [32], Birk and Hering [33], and Torrisi et al. [34].

2.1. Periphyton


Periphytons are valuable indicators of environmental conditions in streams and rivers. As primary producers, periphytons act as important foundation of food webs in river ecosystems [9, 35-36]. Periphytons generally have rapid reproduction rates and very short life cycles and therefore can be expected to reflect short-term impacts and sudden changes in the environment [37-38]. Because the assemblages usually attach to substrate, their growing and prospering can respond directly and sensitively to many physical, chemical and biological variation occurring in the stream (or river) reach, including temperature, nutrient levels, current regimes and grazing etc.[37, 39-45].

Periphytons, especially diatoms, have been preferred for river biomonitoring purposes by many authors [20, 46-55]. Taxa richness and diversity [56-59], assemblage similarity [60-61], taxonomic composition [62], Chlorophyll a [63-65] and biomass [46] have all been reported as measures to indicate the environmental stress. Furthermore, many biotic indices based on species-specific sensitivities and tolerances have been developed to infer specific or general environmental conditions in streams and rivers. Most of them are indicators of organic pollution ([66], see review [67]). Several biotic indices also have been successfully applied in many studies to estimate the status of river ecosystems, mainly in central and northern European rivers [55, 68-70].

2.2. Benthic Macroinvertebrate

Many countries have a long history of using macroinvertebrates to monitor the ecological status of river ecosystems [71]. Benthic macroinvertebrates are key components of aquatic food webs that link organic matter and nutrient resources (e.g., leaf litter, algae and detritus) with higher trophic levels [72]. These organisms have mostly sedentary habits [73] and are therefore representative of site specific ecological conditions. With the sensitive life stage [74] and relatively long life span [75], they have the ability to integrate the effects of short-term environmental variations. Besides, these assemblages are made up of many species among which there is a wide range of trophic levels and pollution tolerances [73, 75-76], therefore providing strong information for interpreting cumulative effects. Community structure of the assemblages frequently changes in response to environmental disturbances in predictable ways, which is the basis for development of biocriteria to evaluate anthropogenic influences [77]. These responses have been summarized by Gray [78] into three categories, including reduction in diversity, retrogression to dominance by opportunistic (e.g. shorter life-cycle, faster reproducing) species and reduction in individual size of dominating species. For example, in streams and rivers polluted by organic matters [5, 79-80] or heavy metals [78, 81-86], species richness and diversity of the macroinvertebrate community strongly reduces for the direct and indirect impact of contaminants; and, Chironomidae commonly possesses the dominant status at the expense of other more sensitive groups, such as stoneflies (Ephemeroptera), caddisflies (Plecoptera) and mayflies (Trichoptera).

Studies on the potential use of benthic macroinvertebrates as bioindicators for river ecosystems have been widely reported in literatures [4, 87-94]. Benthic macroinvertebrates, especially aquatic insects, have been traditionally used in the biomonitoring of stream and river ecosystems for various environmental stress types, such as organic pollution [95-98], heavy metals [82, 86, 99], hydromorphological degradation [26, 100-101], nutrient enrichment [71, 102-106], acidification [107-110] and general stressors [38, 111-112]. Indeed, the assemblages constitute the basis of most biomonitoring program currently in Europe and North America. Many countries (or states or water authorities) even have developed their own biotic indices (e.g. Netherlands [113], France [114], Belgium [115], Denmark [116], U.K [117], Switzerland [118] and U.S.A. [119]; see Fig.1).



2.3. Fish

As highly visible and valuable components of the freshwater ecosystems, fish communities have been applied to monitor river ecosystem health for a long time [27, 120-121]. Fish are the top of the aquatic food web and are consumed by humans, which makes them important for assessing contamination [38]. Due to their relatively long life cycle and mobility, they can be good indicators of long-term (several years) effects and broad habitat conditions [38]. In addition, with wide range of trophic level, including the highest level occupied by top predators, community structure of fish assemblage is reflective of integrated aquatic environment health [122-125].

Fish communities respond significantly and predictably to almost all kinds of anthropogenic disturbances, including eutrophication, acidification, chemical pollution, flow regulation, physical habitat alteration and fragmentation, human exploitation and introduced species [122-123, 126-130]. Their sensitivities to the health of surrounding aquatic environments form the basis for using fishes to monitor environmental degradation [27]. Over the last 30 years, a variety of fish-based biotic indices have been widely used to assess river quality, and the use of multimetric indices, inspired by the index of biotic integrity (IBI) [122-123], has grown rapidly [131].

3. Common Approaches Used for Biomonitoring of River Ecosystems

There are several different biomonitoring techniques currently employed in river ecosystems. The selection of an appropriate technique depends on the issues being addressed and available resources. Potential biomonitoring methods include diversity indices, biotic indices, multimetric approaches, multivariate approaches, functional feeding groups (FFGs) and multiple biological traits.

Bioaccumulation and toxicity of contaminants in indicator species also remain an important component of several river monitoring programs. However, they are not described here since our focus has been on the ecological approaches to measure ecosystem status. In addition, *the saprobic systems* have been once used (in Europe) primarily to indicate oxygen deficits caused by biologically decomposable, organic pollution in running waters, on the basis of Saprobic values of indicator species (mainly bacteria, algae, protozoans and rotifers, but also some macroinvertebrates and fish). However, by the mid-1970s, these indices have been rejected by most European countries for their limits [132-134].

3.1. Diversity Indices

As traditional biomonitoring approaches, many diversity indices have been developed to describe responses of a community to environment variation, combining the three components of community structure, namely *richness* (number of species present), *evenness* (uniformity in the distribution of individuals among the species) and *abundance* (total number of individuals present) (e.g., Shannon-Wiener Index [135], Simpson Index [136], Margalef Index [137]; see review [5]). The assumption is that undisturbed environments are characterized by high diversity or richness, an even distribution of individuals among the species, and moderate to high counts of individuals. The best use of diversity-related indices in river and stream monitoring is probably as an indicator of changes in species composition when comparing impacted and reference assemblages [57]. Many criticisms have been made against the usefulness of diversity indices when employed separately in assessment of river systems [5], and now these indices are preferred to be used together with other metrics (see 3.3 *Multimetric Approaches* below).

3.2. Biotic Indices

Biotic approach, as defined by Tolkamp [138], combines the relative abundance on the basis of certain taxonomic groups with their sensitivities or tolerances into a single index or score. The sensitivity and tolerance of indicator assemblages to a number of environmental characteristics, such as organic pollution, heavy metals, pesticides, eutrophication and pH, are known to differ among species. Therefore, these species-specific pollution indications can be used to infer environmental conditions in a habitat. Biotic indices of macroinvertebrate and periphyton are widely used in European countries. To take benthic macroinvertebrate for an example, numerous biotic index and score systems have been developed [5] (as illustrated in Fig.1). Commonly used biotic indices for macroinvertebrates include Trent Biotic Index (TBI) and Extended Biotic Index (EBI), Chandler's Score System, Biological Monitoring Working Party Score System (BMWP) and ASPT (Average Score per Taxon), Hilsenhoff's Biotic Index (HBI) etc. Among these indices, BMWP and its derivative, IBMWP, are recommended by the Water Framework Directive and widely used in the European Union.

3.3. *Multimetric Approaches*

Multimetric indices represent a means to integrate a set of variable or metrics, which represent various structural and functional attributes of an ecosystem (such as taxa richness, relative abundance, dominance, functional feeding groups, pollution tolerance, life history strategies, disease, and density), therefore provide robust and sensitive insights into the responses of an assemblage to natural and anthropogenic stressors [27, 112, 122, 139-140]. Since Karr [122] first introduced Index of Biotic Integrity (IBI) on the basis of fish assemblages, similar indices have been developed for benthic macroinvertebrates [141-143], fish [144-146], periphyton [50, 147]. By now, multimetric approaches for benthic macroinvertebrates have been the most widely used approach for river biomonitoring in USA [140] and recently used in other parts of the world as well [148-150].

3.4. *Multivariate Approaches*

Multivariate approaches have been initially introduced to assess the biological status of rivers within the UK, with the development of RIVPACS (River Invertebrate Prediction and Classification System) [151]. Multivariate approaches adopt statistical analyses to predict site-specific fauna patterns, which are expected in the absence of major environmental stress; and, the biological evaluations are then performed by comparing the observed fauna at the site with the expected fauna [152-153]. Multivariate approaches have been proven to be effective for biomonitoring. Several predictive models using multivariate techniques are widely used, such as RIVPACS and its derivative, AusRivAS (Australian Rivers Assessment System) [154], BEAST (Benthic Assessment Sediment) [155-156], or the recent ANNA (Assessment by Nearest Neighbor Analysis) [157]. In recent studies, except for macroinvertebrate, multivariate approaches have been developed for periphytons and fishes (e.g., Joy and Death [28]).

3.5. *Functional Approaches*

It is generally recognized that adequate characterization of ecosystems requires information on both structure (pattern) and function (process) [158]. Thus, although assemblage structure and composition has been successfully used in studies of impairment, there has been a recent renaissance in the use of function analyses as a complementary approach to reflecting ecological integrity.

3.5.1 *Functional Feeding Groups (FFGs)*

Analyses of Functional feeding groups (FFGs) are the key components of river continuum concept (RCC) [159] and have been applied to assess ecosystem-level processes in rivers and wetlands [160-162]. In river biomonitoring,

FFGs measures have been used in the forms of single feeding groups (as absolute or relative abundance), ratios between two groups, or composite index that includes several trophic aspects [e.g., the Index of Trophic Completeness (ITC) [163]. In recent years, these measures have been combination with other metrics and applied in biomonitoring approaches (see 3.3 *Multimetric Approaches* above, e.g. “Benthic Index of Biotic Integrity” [141], “Florida Stream Condition Index” [142]). These evaluation have been performed on the basis of easily observed morphological and behavioral attributes, which are associated with feeding and modes of attachment, concealment, and locomotion, together with life-history patterns (voltinism) and drift propensity [164].

3.5.2 Multiple Biological Traits

Biological traits are related to habitat characteristics and the biological and ecological functions of species, thus permit a view into the function structure of bioconosis [165]. Multiple biological traits of aquatic and terrestrial organisms (e.g., size, body form, life cycle, food and feeding habits, reproductive and other traits) in the context of environmental constraints [25] have been recently developed for freshwater biomonitoring. The utilization of multiple traits generally has been combined with *Multivariate Approaches* (see 3.4 above). Currently, multiple biological traits are mainly used for aquatic invertebrates in the running waters of Europe, and relevant researches proposed a multitude of traits that are weighted by the abundance or occurrence of the taxa [165]. Similar attempts also have been performed on fish assemblages [166]. Several applications of trait based methods (e.g., in relation to river pollution [167-168], anthropogenic influences in general [111, 162] or ecological assessment theories [25]) demonstrate the potentialities of investigating trait structures.

4. Trends in Biomonitoring of River Ecosystems

4.1. Increasing Application of Functional Measures

By now, diverse biomonitoring techniques have already been developed to quantify the human impact on the environment of streams and rivers. However, because of the new trends in environmental policies, ecologists are currently facing new demands of effective tools to correlate the current status of ecosystems and the management for conservation and restoration [169-171]. Therefore, there are increasing applications of functional measures in river biomonitoring, including microbial enzyme activity [50, 61], bacterial luminescence [172], photosynthesis [173], respiration [174-175], locomotory activity [176], fluctuating asymmetry [177], community metabolism (primary productivity and respiration) [173], nutrient uptake and spiraling [178], and secondary production [179-181], except for FFGs and multiple biological traits mentioned above.

4.2. Molecular Techniques

In recent years, some efforts have been attempted to apply molecular techniques as biomonitoring tools. Molecular approaches used in biomonitoring mainly focus on the species identification and genetic diversity.

It is no doubt that finer taxonomic resolutions are ideal to obtain the most complete analysis of ecosystem health. Unfortunately, the acquisition of genus or species-level information for macroinvertebrates and periphytons is time consuming; and even with high levels of taxonomic skill, misidentifications of species may still result. However, genera or species can be rapidly identified at any life stage by molecular markers. Recent researches demonstrate the accuracy and effectiveness of DNA-based methods as biomonitoring tools, such as PCR-RFLP, T-RFLP and COI sequence, which have been used for Chironomids and periphytons in aquatic systems [182-184].

Genetic diversity is fundamentally a trait of biological populations, and significant changes in genetic diversity reflect important population-level changes. Since data of genetic diversity offer powerful tools for examining the current status of populations, inferring the history of population changes, and anticipating future population directions, molecular approaches provide a logical extension of previously described approaches to measure the

variation of environmental status. Such attempts to relate the variability of molecular genetic markers to specific aquatic stressors date back more than 30 years. These studies include both field surveys and controlled laboratory experiments of fish populations, and have evaluated the effects of metals, acidity, pesticides, radionuclides, and complex effluents (see review [185]). Moreover, USEPA has carried out a series of researches to assess the utility of incorporating a genetic diversity indicator into large-scale assessment and monitoring efforts [186]. Although the application of molecular genetic diversity in river monitoring are still in their infancy, there are a number of compelling reasons to believe that molecular genetic measures will ultimately provide highly useful bioindicators.

Acknowledgements

We would like to thank the support of National Major Project of Water Pollution Control and Prevention Technology (2008ZX07528-002-003).

References

- [1] World Resources Institute, 2001. Part 2: Taking stock of ecosystem. In: World Resources Institute (Ed) World Resources 2000-2001. Elsevier, Washington, DC, pp. 43-145.
- [2] Malmqvist, B. and Rundle, S., 2002. Threats to the running water ecosystems of the world. *Environment Conservation*, 29:134-153.
- [3] Saunders, D.L., Meeuwig, J.J and Vincent, A.C.J, 2002. Freshwater protected areas: Strategies for Conservation. *Conservation Biology*, 16(1): 30-41.
- [4] Rosenberg, D.M. and Resh, V.H. (Eds.), 1993. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York.
- [5] Metcalfe, J.L., 1989. Biological Water Quality Assessment of Running Waters Based on Macroinvertebrate Communities: History and Present Status in Europe. *Environmental Pollution*, 60: 101-139.
- [6] Soininen, J. and Könönen, K., 2004. Comparative study of monitoring South-Finnish rivers and streams using macroinvertebrates and benthic diatom community structure. *Aquatic Ecology*, 38: 63-75.
- [7] Patrick, R., 1973. Use of algae, especially diatoms, in the assessment of water quality. In: Cairns, J. Jr., Dickson, K.L. (Eds.). *Biological Methods for the Assessment of Water Quality*. Journal of the American Society for Testing and Materials, Philadelphia, pp. 76-95.
- [8] Stevenson, R.J., Lowe, R.L., 1986. Sampling and interpretation of algal patterns for water quality assessments, In: Isorn, B.G. (Ed.), *Rationale for Sampling and Interpretation of Ecological Data in the Assessment of Freshwater Ecosystems*. American Society for Testing and Materials, Philadelphia, pp. 118-149.
- [9] Lowe, R.L. and Pan, Y., 1996. Benthic algal communities as biological indicators. In: Stevenson, R.J., Bothwell, M.L., Lowe, R.L. (Eds.). *Algal Ecology: Freshwater Benthic Ecosystems*. Academic Press, San Diego, pp. 705-739.
- [10] Stevenson, R.J. and Pan, Y.D., 2001. Assessing environmental conditions in rivers and streams with diatoms. In: Stoermer, E.F. and Smol, J.P. (Eds.). *The Diatoms: Applications for the Environmental and Earth Sciences*. Cambridge University Press, Cambridge, pp. 11-40.
- [11] Worf, D.L., 1980. *Biological monitoring for environmental effects*. Lexington Books, Toronto.
- [12] Rosenberg, D.M. 1998. A National Aquatic Ecosystem Health Program for Canada: We should go against the flow. *Bull. Entomol. Soc. Can.*, 30(4):144-152.
- [13] Gerhardt, A. (Ed), 1999. *Biomonitoring of Polluted Water - Reviews on Actual Topics*. Environmental Research Forum, Vol. 9. Trans Tech Publications - Scitech Publications. Uetikon-Zuerich, Switzerland, pp. 1-13.
- [14] Oertel, N. and Salánki, J., 2003. *Biomonitoring and Bioindicators in Aquatic Ecosystems*. In: Ambasht RS, Ambasht NK (Eds.) *Modern trends in applied aquatic ecology*. Kluwer Academic/Plenum Publishers, New York, pp. 219-246.
- [15] Markert, B., Wappelhorst, O., Weckert, V., Herpin, U., Siewers, U. and Friese, K., 1999. The use of bioindicators for monitoring the heavy-metal status of the environment. *Journal of Radioanalytical Nuclear Chemistry*, 240(2): 425-429.

- [16] Markert, B., Oehlmann, J. and Roth, M., 1997. General aspects of heavy metal monitoring by plants and animals. In: Subramanian G, Iyengar V, editors. *Environmental biomonitoring - exposure assessment and specimen banking*. ACS Symposium Series, Vol. 654. Washington DC: American Chemical Society.
- [17] Markert B, Breure T, Zechmeister H (Eds.), 2003. *Bioindicators and biomonitors - principles, concepts and applications*. Amsterdam: Elsevier.
- [18] Hilty, J. and Merenlender, A., 2000. Faunal indicator taxa selection for monitoring ecosystem health. *Biological Conservation*, 92: 185-197.
- [19] Füreder, L. and Reynolds, J.D., 2003. Is *Austropotamobius pallipes* a good bioindicator?. *Bulletin Français de la Pêche et de la Pisciculture*, 370-371: 157-163.
- [20] Whitton, B.A., Rott, E. (Eds.), 1996. *Use of Algae for Monitoring Rivers II*. Institut für Botanik. Universität Innsbruck, Innsbruck, Austria, 196 pp.
- [21] Vis, C., Hudon, C., Cattaneo, A., Pinel-Alloul, B., 1998. Periphyton as an indicator of water quality in the St Lawrence River (Quebec, Canada). *Environmental Pollution*, 101(1): 13-24.
- [22] Prygiel, J., Whitton, B.A., Bukowska, J. (Eds.), 1999. *Use of algae for monitoring rivers III*. Agence de l'Eau Artois-Picardie, Douai, France, 271 pp.
- [23] Coste, M., Botry, S., Tison-Rosebery, J., Delmas, F., 2008. Improvements of the Biological Diatom Index (BDI): description and efficiency of the new version (BDI-2006). *Ecological Indicators*, 9: 621-650.
- [24] Lenat, D.R. and Barbour, M.T., 1994. Using benthic macroinvertebrate community structure for rapid, cost-effective, water quality monitoring: rapid bioassessment. In: Loeb, S.L., Spacie, A. (Eds.), *Biological Monitoring of Aquatic Systems*. Lewis Publishers, Boca Raton, FL, pp. 187-215.
- [25] Statzner, B., Bis, B., Dolédec, S. and Usseglio-Polatera, P., 2001. Perspectives for biomonitoring at large scales: a unified measure for the functional composition of invertebrate communities in European running waters. *Basic and Applied Ecology*, 2: 73-85.
- [26] Buffagni, A., Erba, S., Cazzola, M. and Kemp, J.L., 2004. The AQEM multimetric system for the southern Italian Apennines: assessing the impact of water quality and habitat degradation on pool macroinvertebrates in Mediterranean rivers. *Hydrobiologia*, 516: 313-329.
- [27] Fausch, K.D., Lyons, J., Karr, J.R. and Angermeier, P.L., 1990. Fish communities as indicators of environmental degradation. *American Fisheries Society Symposium*, 8: 123-144.
- [28] Joy, M.K. and Death, R.G., 2002. Predictive modeling of freshwater fish as a biomonitoring tool in New Zealand. *Freshwater Biology*, 47(11): 2261-2275.
- [29] Oberdorff, T., Pont, D., Hugué, B. and Porcher, J.P., 2002. Development and validation of a fish-based index (FBI) for the assessment of 'river health' in France. *Freshwater Biology*, 47: 1720-1734.
- [30] Pont, D., Hugué, B., Beier, U., Goffaux, D., Melcher, A., Noble, R., Rogers, C., Roset, N., and Schmutz, S., 2006. Assessing river biotic condition at a continental scale: a European approach using functional metrics and fish assemblages. *Journal of Applied Ecology*, 43: 70-80.
- [31] Scuri, S., Torrisi, M., Cocchioni, M. and Dell'Uomo, A., 2006. The European Water Framework Directive 2000/60/EC in the evaluation of the ecological status of watercourses. Case study: the river Chienti (central Apennines, Italy). *Acta Hydrochimica et Hydrobiologica*, 34: 498-505.
- [32] Carlisle, D.M., Hawkins, C.P., Meador, M.R., Potapova, M., and Falcone, J., 2008. Biological assessments of Appalachian streams based on predictive models for fish, macroinvertebrate, and diatom assemblages. *Journal of the North American Benthological Society*, 27(1): 16-37.
- [33] Birk, S., Hering, D., 2009. A new procedure for comparing class boundaries of biological assessment methods: a case study from the Danube Basin. *Ecological Indicators* 9, 528-539.
- [34] Torrisi, M., Scuri, S., Dell'Uomo, A., Cocchioni, M., 2010. Comparative monitoring by means of diatoms, macroinvertebrates and chemical parameters of an Apennine watercourse of central Italy: The river Tenna. *Ecological Indicators*, 10: 910-913.
- [35] Mayer, M.S. and Likens, G.E., 1987. The importance of algae in a shaded headwater stream as food for an abundant caddisfly (Trichoptera). *Journal of the North American Benthological Society*, 6(4): 262-269.
- [36] Lamberti, G. A., 1996. The role of periphyton in benthic food webs. Pages 533–572 in R. J. Stevenson, M. L. Bothwell, and R. L. Lowe, editors. *Algal ecology: freshwater benthic ecosystems*. Academic Press, San Diego, California, USA.

- [37] McCormick, P., Stevenson, R., 1998. Periphyton as a tool for ecological assessment and management in the Florida Everglades. *Journal of Phycology*, 34: 726-733.
- [38] Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. *Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish*, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency; Office of Water; Washington, D.C.
- [39] Squires, L. E., Rushforth, S. R. and Brotherson, J. D., 1979. Algal response to a thermal effluent: Study of a power station on the Provo River, Utah, USA. *Hydrobiologia*, 63: 17-32.
- [40] Horner, R.R., Welch, E.B., 1981. Stream periphyton development in relation to current velocity and nutrients. *Canadian Journal of Fisheries and Aquatic Sciences* 38,449-457.
- [41] Descy, J.P. and Mouvet, C., 1984. Impact of the Tihange nuclear power plant on the periphyton and the phytoplankton of the Meuse River (Belgium). *Hydrobiologia*, 119: 119-128.
- [42] Pringle, C.M. and Bowers, J.A., 1984. An in situ substratum fertilization technique: Diatom colonization on nutrient-enriched, sand substrata. *Canadian Journal of Fisheries and Aquatic Sciences*, 41: 1247-1251.
- [43] Steinman, A.D., McIntire, C.D., 1986. Effects of current velocity and light energy on the structure of periphyton assemblages in laboratory streams. *Journal of Phycology*, 22: 352-361.
- [44] Steiman, A.D., McIntire, C.D., Gregory, S.V., Lamberti, G.A. and Ashkenas, L.R., 1987. Effects of herbivore type and density on taxonomic structure and physiology of algal assemblages in laboratory streams. *Journal of the North American Benthological Society*, 6: 175-188.
- [45] Pan, Y., Stevenson, R.J., Hill, B.H., Herlihy, A.T., Collins, G.B., 1996. Using diatoms as indicators of ecological conditions in lotic systems: a regional assessment. *Journal of the North American Benthological Society*, 15: 481-495.
- [46] Biggs, B.J.F., 1989. Biomonitoring of organic pollution using periphyton, South Branch, Canterbury, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 23: 263-274.
- [47] Kutka, F. and Richards, C., 1996. Relating diatom assemblage structure to stream habitat quality. *Journal of the North American Benthological Society*, 15: 469-480.
- [48] Mattila, J. and Räsänen, R., 1998. Periphyton growth as an indicator of eutrophication: an experimental approach. *Hydrobiologia*, 377(1):15-23.
- [49] Rott, E., Duthie, H. and Pipp, E., 1998. Monitoring organic pollution and eutrophication in the Grand River, Ontario, by means of diatoms. *Canadian Journal of Fisheries and Aquatic Sciences*, 55: 1443-1453.
- [50] Hill, B., Herlihy, A., Kaufmann, P., Stevenson, R., McCormick, F., Johnson, C., 2000a. Use of periphyton assemblage data as an index of biotic integrity. *Journal of the North American Benthological Society*, 19: 50-67.
- [51] Winter, J., and Duthie, H., 2000a. Epilithic diatoms as indicators of stream total N and total P concentration. *Journal of the North American Benthological Society*, 19: 32-49.
- [52] Winter, J. and Duthie, H., 2000b. Stream biomonitoring at an agricultural test site using benthic algae. *Can. J. Bot.*, 78: 1319-1325.
- [53] Munn, M., Black, R. and Gruber, S., 2002. Response of benthic algae to environmental gradients in an agriculturally dominated landscape. *Journal of the North American Benthological Society*, 21: 221-237.
- [54] Potapova, M., Charles, D., 2003. Distribution of benthic diatoms in U.S. rivers in relation to conductivity and ionic composition. *Freshwater Biology*, 48: 1311-1328.
- [55] Gomà, J., et al., 2005. Diatom communities and water quality assessment in Mountain Rivers of the upper Segre basin (La Cerdanya, Oriental Pyrenees). *Hydrobiologia*, 551, 209-225.
- [56] Weitzel, R.L., Bates, J.M., 1981. Assessment of effluent impacts through evaluation of periphyton diatom community structure. In: Bates, J.M., Weber, C.I. (Eds.). *Ecological Assessments of Effluent Impacts on Communities of Indigenous Aquatic Organisms*, American Society for Testing and Materials, Philadelphia, pp. 142-165.
- [57] Stevenson, R.J., 1984. Epilithic and epipellic diatoms in the Sandusky River, with emphasis on species diversity and water pollution. *Hydrobiologia*, 114: 161-175.
- [58] Khan, I.S.A.N., 1991. Effect of urban and industrial wastes on species diversity of the diatom community in a tropical river. Malaysia. *Hydrobiologia*, 224: 175-184.
- [59] Whitton, B.A., Rott, E., Friedrich, G. (Eds), 1991. *Use of Algae for Monitoring Rivers*. Institut für Botanik, Universität Innsbruck, Innsbruck, Austria.

- [60] Medley, C.N. and Clements, W.H., 1998. Responses of diatom communities to heavy metals in streams: the influence of longitudinal variation. *Ecological Applications*, 8: 633-644.
- [61] Hill, B.H., Willingham, W.T., Parrish, L.P. and McFarland, B.H., 2000b. Periphyton assemblage responses to elevated metal concentrations in a Rocky Mountain stream. *Hydrobiologia*, 428: 161-169.
- [62] Archibald, R.E.M., 1972. Diversity in some South African diatom associations and its relation to water quality. *Water Research*, 6: 1229-1238.
- [63] Leland, H.V., 1995. Distribution of phytobenthos in the Yakima River basin, Washington, in relation to geology, land use, and other environmental factors. *Canadian Journal of Fisheries and Aquatic Sciences*, 52: 1108-1129.
- [64] Pan, Y., Stevenson, R.J., Hill, B.H., Kaufmann, P.R. and Herlihy, A.T., 1999. Spatial patterns and ecological determinants of benthic algal assemblages in mid-Atlantic streams, USA. *Journal of Phycology*, 35: 460-468.
- [65] Pan, Y., Stevenson, R.J., Hill, B.H. and Herlihy, A.T., 2000. Ecoregions and benthic diatom assemblages in mid-Atlantic Highland streams, USA. *Journal of the North American Benthological Society*, 19: 518-540.
- [66] Whitton, B.A. and Kelly, M.G., 1995. Use of algae and other plants for monitoring. *Austral Ecology*, 20(1): 45-56.
- [67] Coste, M., Boca, C. and Dauta, A., 1991. Use of algae for monitoring rivers in France. In: Whitton, B.A., Rott, E. and Friedrich, G. (Eds). *Use of algae for monitoring rivers*. Institut für Botanik, Universität Innsbruck, Innsbruck, Austria. pp 75-88.
- [68] Prygiel, J., and Coste, M., 1993. The assessment of water quality in the Artois-Picardie water basin (France) by the use of diatom indices. *Hydrobiologia*, 269/270: 343-349.
- [69] Kelly, M. G. and Whitton, B. A., 1995. The trophic diatom index: A new index for monitoring eutrophication in rivers. *Journal of Applied Phycology*, 7: 433-444.
- [70] Eloranta, P., and Soininen, J., 2002. Ecological status of some Finnish rivers evaluated using benthic diatom communities. *Journal of Applied Phycology*, 14: 1-7.
- [71] Hellawell, J.M., 1986. *Biological indicators of Freshwater Pollution and Environmental Management*. Elsevier Applied Science, London, UK, 546 pp.
- [72] Wallace, J.B. and Webster, J.R., 1996. The role of macroinvertebrates in stream ecosystem function. *Annual Review of Entomology*, 41: 115-139.
- [73] Cook, S.E.K., 1976. Quest for an index of community structure sensitive to water pollution. *Environmental Pollution*, 11: 269-288.
- [74] Hutchinson, T.H., Solbe, J. and Klopper-Sams, P.J., 1998. Analysis of the ecetoc aquatic toxicity (EAT) database III-Comparative toxicity of chemical substances to different life stages of aquatic organisms. *Chemosphere*, 36(1): 129-142.
- [75] Pratt, J.M. and Coler, R.A., 1976. A procedure for the routine biological evaluation of urban runoff in small rivers. *Water Research*, 10: 1019-1025.
- [76] France, R.L., 1990. Theoretical framework for developing and operationalizing an index of zoobenthos community integrity: Application to biomonitoring with zoobenthos communities in the Great Lakes. In: Edwards, C.J. and Regier, H.A. (Eds.), *An Ecosystem Approach to the Integrity of the Great Lakes in Turbulent Times*, Great Lakes Fishery Commission Special Publication, Ann Arbor, MI. pp. 169-193.
- [77] Boyle, T.P. and Fraleigh Jr., H.D., 2003. Natural and anthropogenic factors affecting the structure of the benthic macroinvertebrate community in an effluent-dominated reach of the Santa Cruz River, AZ. *Ecological Indicators*, 3: 93-117.
- [78] Gray, L.J., 1989. Emergence production and export of aquatic insects from a tallgrass prairie stream. *Southwestern Naturalist*, 34: 313-318.
- [79] Hawkes, H.A. and Davies, L.J., 1971. Some effects of organic enrichment on benthic invertebrate communities in stream riffles. In *The scientific management of animal and plant communities for conservation*, Duffey E. and Watt, A.S. Symposium of the British Ecological Society, 11th, 271-93. Oxford, Blackwell Scientific Publications.
- [80] Whitehurst, I.T. and Lindsey, B.I., 1990. The impact of organic enrichment on the benthic macroinvertebrate communities of a lowland river. *Water Research*, 24(5): 625-630.
- [81] Winner, R. W., Scott Van Dyke, J., Caris, N. and Farrell, M. P., 1975. Responses of the macroinvertebrate fauna to a copper gradient in an experimentally-polluted stream. *Verh. Int. Verein. Limnol.*, 19: 2121-2127.
- [82] Winner, R.W., Boesel, M.W. and Farrell, M.P., 1980. Insect community structure as an index of heavy-metal pollution in lotic ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(4): 647-655.
- [83] Clements, W.H., 1994. Benthic invertebrate community responses to heavy metals in the Upper Arkansas River Basin, Colorado. *Journal of the North American Benthological Society*, 13(1): 30-44.

- [84] Groenendijk, D., Postma, J.F., Kraak, M.H.S. and Admiraal, W., 1998. Seasonal dynamics and larval drift of *Chironomus riparius* (Diptera) in a metal contaminated lowland river. *Aquatic Ecology*, 32: 341-351.
- [85] Hickey, C.W. and Clements, W.H., 1998. Effects of heavy metals on benthic macroinvertebrate communities in New Zealand streams. *Environmental Toxicology and Chemistry*, 17(11): 2238-2346.
- [86] Smolders, A.J.P., Lock, R.A.C, Van der Velde, G., Medina Hoyos, R.I. and Roelofs, 2003. Effects of mining activities on heavy metal concentrations in water, sediment and macroinvertebrates in different reaches of the Pilcomayo River, South America. *Archives of Environmental Contamination and Toxicology*, 44(3): 314-323.
- [87] Cairns Jr., J. and Der Schalie, W.H.V.D., 1980. Review paper: Biological monitoring. Part I- Early warning systems. *Water Research*, 14: 1179-1196.
- [88] Cairns Jr., J., 1981. Review paper: Biological monitoring. Part VI-Future needs. *Water Research*, 15: 941-952.
- [89] Matthews, R., Buikema, A.L., Cairns Jr., J., and Rodgers Jr., J.H., 1982. Review paper: Biological monitoring. Part IIA-Receiving system functional methods, relationships and indices. *Water Research*, 16: 129-139.
- [90] Herricks, E. and Cairns Jr., J., 1982. Biological monitoring. Part III Receiving system methodology based on community structure. *Water Research*, 16: 141-153.
- [91] Buikema Jr., A.L., Niederlehner, B.R. and Cairn Jr., J., 1982. Biological monitoring. Part IV- Toxicity testing. *Water Research*, 16: 239-262.
- [92] Cherry, D.S. and Cairns Jr., J., 1982. Biological monitoring. Part VPreference and avoidance studies. *Water Research*, 16: 263-301.
- [93] Ogeibu, A.E. and Oribhabor, B.J., 2002. Ecological impact of river impoundment using benthic macro-invertebrates as indicators. *Water Research*, 36: 2427-2436.
- [94] Mason, C.F. and Parr, B.L., 2003. Long-term trends in water quality and their impact on macroinvertebrate assemblages in eutrophic lowland rivers. *Water Research*, 37: 2969-2979.
- [95] Zelinka, V. and Marvan, P., 1961. Zur Prazisierung der biologischen Klassifikation der Reinheit fließender Gewässer. *Archive fur Hydrobiologia*, 57: 389-407.
- [96] Armitage, P.D., Moss, D., Wright, J.F. and Furse, M.T., 1983. The performance of a new biological water quality score based on macroinvertebrates over a wide range of unpolluted running-water sites. *Water Research*, 17: 333-347.
- [97] Rae, J.G., 1989. Chironomid midges as indicators of organic pollution in the Scioto River Basin, Ohio. *The Ohio Journal of Science*, 89(1): 5-9.
- [98] Zamora-Muñoz, C. and Alba-Tercedor, J., 1996. Bioassessment of organically polluted Spanish rivers, using a biotic index and multivariate methods. *Journal of the North American Benthological Society*, 15(3): 332-352.
- [99] Poulton B.C., Monda, D.P., Woodward, D.F., Wildhaber, M.L. and Brumbaugh, W.G., 1995. Relations between benthic community structure and metals concentrations in aquatic macroinvertebrates: Clark Fork River, Montana. *Journal of Freshwater Ecology*, 10(3): 277-293.
- [100] Lorenz, A., Hering, D., Feld, C.K. and Rolauffs, P., 2004. A new method for assessing the impact of hydromorphological degradation on the macroinvertebrate fauna of five German stream types. *Hydrobiologia*, 516: 107-127.
- [101] Friberg, N., Sandin, L. and Pedersen, M.L., 2009. Assessing the effects of hydromorphological degradation on macroinvertebrate indicators in rivers: examples, constraints, and outlook. *Integrated Environmental Assessment and Management*, 5(1): 86-96.
- [102] Hynes, H.B.N., 1960. *The biology of polluted water*. Liverpool University Press, Liverpool, England.
- [103] Hellawell, J.M., 1978. *Biological surveillance of rivers, a biological monitoring hand- book*. Water Research Centre, Medmenham and Stevenage, UK, 331 pp.
- [104] Mason, C.F., 1996. *Biology of Freshwater Pollution*, 3rd edition. Longman, London.
- [105] Hering, D., Feld, C.K., Moog, O., Ofenböck, T., 2006. Cook book for the development of a Multimetric-Index for biological condition of aquatic ecosystems: experiences from the European AQEM and STAR projects and related initiatives. *Hydrobiologia*, 566:311-324.
- [106] Johnson, R.K., Hering, D., Furse, M.T., Verdonschot, P.F.M., 2006. Indicators of ecological change: comparison of the early response of four organism groups to stress gradients. *Hydrobiologia*, 566: 139-152.
- [107] Sandin, L. and Johnson, R.K., 2000. The statistical power of selected indicator metrics using macroinvertebrates for assessing acidification and eutrophication of running waters. *Hydrobiologia*, 422/423: 233-243.

- [108] Braukmann, U., 2001. Stream acidification in South Germany-chemical and biological assessment methods and trends. *Aquatic Ecology*, 35: 207-232.
- [109] Sandin, L., Dahl, J., and Johnson, R.K., 2004. Assessing acid stress in Swedish boreal and alpine streams using benthic macroinvertebrates. In: Hering, D., Verdonshot, P.F.M., Moog, O., Sandin, L. (Eds.), *Integrated Assessment of Running Waters in Europe*. *Hydrobiologia*, 516: 129-148.
- [110] Davy-Bowker, J., Murphy, J.F.; Rutt, G.P., Steel, J.E.C. and Furse, M.T., 2005. The development and testing of a macroinvertebrate biotic index for detecting the impact of acidity on streams. *Archiv für Hydrobiologie*, 163(3): 383-403.
- [111] Dolédec, S., Statzner, B. and Bournard, M., 1999. Species traits for future biomonitoring across ecoregions: patterns along a human-impacted river. *Freshwater Biology*, 42(4): 737-758.
- [112] Karr, J.R. and Chu, E.W., 1999. *Restoring Life in Running Waters: Better Biological Monitoring*. Island Press, Washington, DC.
- [113] Tolkamp, H.H. and Gardiener, J.J.P., 1977. Hydrobiological survey of lowland streams in the Achterhoek (The Netherlands) by means of a system for the assessment of water quality and stream character based on macroinvertebrates. *Mitteilungen des Institutes für Wasserwirtschaft, Hydrologie und landwirtschaftlichen Wasserbau, T.U. Hannover*, 41:215-235.
- [114] Verneaux, J., Galmiche, P., Janier, F. and Monnot, A., 1982. Une nouvelle méthode pratique d'évaluation de la qualité des eaux courantes. Un indice biologique de qualité générale (BIG). *Annales Scientifiques de l'Université de Franche-Comté Besançon, Biologie animale 4 ème série, fasc. 3, Note No. 2*: 11-21.
- [115] De Pauw, N. and Vanhooren, G., 1983. Method for biological quality assessment of watercourses in Belgium. *Hydrobiologia*, 100: 153-168.
- [116] Andersen, M.M., Riget, F.F. and Sparholt H., 1984. A modification of the Trent Index for use in Denmark. *Water Research*, 18: 141-151.
- [117] Extence, C.A. and Ferguson, J.D., 1989. Aquatic invertebrate surveys as a water quality management tool in the Anglian water region. *Regulated Rivers: Research and Management*, 4: 139-146.
- [118] Lang, C. and Reymond, O., 1995. An improved index of environmental quality of Swiss rivers based on benthic invertebrates. *Aquatic Sciences*, 57: 172-180.
- [119] Hilsenhoff, W.L., 1987. An improved biotic index of organic stream pollution. *Great Lakes Entomologist*, 20: 31-39.
- [120] Plafkin, J. L., M. T. Barbour, K. D. Porter, S. K. Gross & R. M. Hughes, 1989. *Rapid Bioassessment Protocols for Use in Streams and Rivers: Benthic Macroinvertebrates and Fish*. EPA/444/4 - 89 - 001 U.S. Environmental Protection Agency, Washington, DC.
- [121] Simon, T.P. 1991. Development of ecoregion expectations for the index of biotic integrity (IBI) Central Corn Belt Plain. U.S. Environmental Protection Agency, Region V, Chicago, Illinois. EPA 905/9-91/025.
- [122] Karr, J.R., 1981. Assessment of biotic integrity using fish communities. *Fisheries*, 6(6): 21-27.
- [123] Karr, J.R., Fausch, K.D., Angermeier, P.L., Yant, P.R. and Schlosser, I.J., 1986. *Assessing Biological Integrity in Running Waters: A Method and its Rationale*. Special Publication No. 5. Illinois Natural History Survey, Champaign, IL.
- [124] Harris, J.H., 1995. The use of fish in ecological assessments, *Aust. J. Ecol.*, 20: 65-80.
- [125] Shields, F.D., Knight Jr., S.S., and Cooper, C.M., 1995. Use of the index of biotic integrity to assess physical habitat degradation in warmwater streams. *Hydrobiologia*, 312(3):191-208.
- [126] Hartman, W.L., 1973. Effects of exploitation, environmental changes, and new species on the fish habitats and resources of Lake Eire. *Great Lakes Fishery Commission Technical Report 22*, Ann Arbor, Michigan.
- [127] Hughes, R.M., 1985. Use of watershed characteristics to select control streams for estimating effects of metal mining wastes on extensively disturbed streams. *Environmental Management*, 9: 253-262.
- [128] Leonard, P.M. and Orth, D.J., 1986. Application and testing of an index of biotic integrity in small, coolwater streams. *Transactions of the American Fisheries Society*, 115: 401-414.
- [129] Berkman, H.E. and Rabeni, C.F., 1987. Effect of siltation on fish communities. *Environmental Biology of Fishes*, 18: 285-294.
- [130] Ormerod, S.J., 2003. Current issues with fish and fisheries: editor's overview and introduction. *Journal of Applied Ecology*, 40: 204-213.
- [131] Simon, T.P., 1999. *Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities*. Lewis Publishers, Boca Raton, FL.
- [132] Chutter, F. M., 1972. An empirical biotic index of the quality of water in South African streams and rivers. *Water Research*, 6: 19-30.

- [133] Persoone, G. and De Pauw, N., 1979. Systems of biological indicators for water quality assessment. In: *Biological Aspects of Freshwater Pollution*, Ravera, O. (ed) Oxford, Pergamon Press, pp. 39-75.
- [134] Jones, J.R., Tracy, B.H., Sebaugh, J.L., Hazelwood, D.H. and Smart, M.M., 1981. Biotic index testing for ability to assess water quality of Missouri Ozark streams. *Transactions of the American Fisheries Society*, 110: 627-637.
- [135] Shannon, C.E. and Weaver, W., 1949. *The Mathematical Theory of Communication*. University of Illinois Press, Urbana.
- [136] Simpson, E.H., 1949. Measurement of Diversity. *Nature*, 163: 688.
- [137] Margalef, R., 1951. Diversidad de especies en las comunidades naturales. *Publnes Inst. Biol. apl.*, Barcelona, 9:5-27.
- [138] Tolkamp, HH, 1985. Biological Assessment of Water Quality in Running Water Using Macroinvertebrates: A Case Study for Limburg, The Netherlands. *Water Science and Technology*, 17(17): 867-878.
- [139] Barbour, M.T., Stribling, J.B. and Karr, J.R., 1995. Multimetric approaches for establishing biocriteria and measuring biological condition. In *Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making*, ed. WS Davis, TP Simon, Boca Raton, FL: Lewis. pp. 63-77.
- [140] Barbour, M.T. and Yoder, C.O., 2000. The multimetric approach to bioassessment as used in the United States of America. See Ref. 149a, pp. 281-292.
- [141] Kerans, B.L., and Karr, J.R., 1994. A benthic index of biotic integrity (B-IBI) for rivers of the Tennes- see valley. *Ecological Applications*, 4:768-785.
- [142] Barbour, M.T., Gerritsen, J., Griffith, G.E., Frydenborg, R., Mccarron, E., White, J.S., Bastian, M.L., 1996. A framework for biological criteria for Florida streams using benthic macroinvertebrates. *Journal of the North American Benthological Society*, 15: 185-211.
- [143] Fore, L.S., Karr, J.R., and Wisseman, R.W, 1996. Assessing invertebrate responses to human activities: Evaluating alternative approaches. *Journal of the North American Benthological Society*, 15(2):212-231.
- [144] Miller, D.L., Leonard, P.M., Hughes, R.M., Karr, J.R., Moyle, P.B., Schrader, L.H., Thompson, B.A., Daniels, R.A., Fausch, K.D., Fitzhugh, G.A., Gammon, J.R., Halliwell, D.B., Angermeier, P.L. and Orth, D.J., 1988. Regional applications of an index of biotic integrity for use in water resource management. *Fisheries*, 13: 12-20.
- [145] Simon T.P. and Emery E.B., 1995. Modification and assessment of an index of biotic integrity to quantify water resource quality in Great Rivers. *Regulated Rivers: Research and Management*, 11: 283-298.
- [146] Hughes, R.M. and Oberdorff, T., 1999. Applications of IBI concepts and metrics to waters outside the United States and Canada. In: T.P. Simon (ed.) *Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities*. Boca Raton, FL: CRC Press, pp. 79-93.
- [147] Hill, B.H., Herlihy, A.T., Kaufmann, P.R., Decelles, S.J. and Vander Borgh, M.A., 2003. Assessment of streams of the eastern United States using a periphyton index of biotic integrity. *Ecological Indicators*, 2: 325-338.
- [148] Sivaramakrishnan, K.G., Hannaford, M.J., and Resh, V.H., 1996. Biological assessment of the Kaveri River catchment, South India, using benthic macroinvertebrates: applicability of water quality monitoring approaches developed in other countries. *Int. J. Ecol. Environ. Sci.*, 22:113-132.
- [149] Thorne, R.S.J. and Williams, W.P., 1997. The response of benthic macroinvertebrates to pollution in developing countries: a multimetric system of bioassessment. *Freshwater Biology*, 37:671-686.
- [150] Vlek HE, Verdonschot PFM, Nijboer RC. 2004. Towards a multimetric index for the assessment of Dutch streams using benthic macroinvertebrates. *Hydrobiologia*, 516:173-189.
- [151] Wright, J.F., 2000. An introduction to RIVPACS. See Ref. 149a, pp. 1-24.
- [152] Niemi, G.J. and McDonald, M.E., 2004. Application of ecological indicators. *Annu. Rev. Ecol. Evol. Syst.* 35:89-111.
- [153] Norris, R.H. and Hawkins, C.P., 2000. Monitoring river health. *Hydrobiologia*, 435:5-17.
- [154] Simpson, J.C. and Norris, R.H., 2000. Biological assessment of river quality: development of AUSRIVAS models and outputs. See Ref. 149a, pp. 125-142.
- [155] Reynoldson, T.B., Bailey, R.C., Day, K.E. and Norris, R.H., 1995. Biological guidelines for freshwater sediment based on Benthic Assessment of Sediment (the BEAST) using a multivariate approach for predicting biological state. *Aust. J. Ecol.* 20:198-219.
- [156] Rosenberg, D.M., Reynoldson, T.B. and Resh, V.H., 2000. Establishing reference conditions in the Fraser River catchment, British Columbia, Canada, using the BEAST (Benthic Assessment of Sediment) predictive model. See Ref. 149a, pp. 181-194.

- [157] Linke, S., Norris, R.H., Faith, D.P. and Stockwell, D., 2005. ANNA: a new prediction method for bioassessment programs. *Freshwater Biology*, 50:147-158.
- [158] Gessner, M.O. and Chauvet, E., 2002. A case for using litter breakdown to assess functional stream integrity. *Ecological Applications*, 12: 498-510.
- [159] Vannote, R.L., Minshall, G.W., Cummins, K.W., Sedell, J.R. and Cushing, C.E., 1980. The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37:130-37.
- [160] Merritt, R.W., Higgins, M.J., Cummins, K.W. and Vandeneeden, B., 1999. The Kissimmee River–Riparian Marsh Ecosystem, Florida. Seasonal differences in invertebrate functional feeding group relationship. In: Batzer DP, Rader R, Wissinger SA (eds) *Invertebrates in freshwater wetlands of North America: ecology and management*. Wiley, New York, pp 55-79.
- [161] Rawer-Jost, C., Böhmer, J., Blank, J., Rahmann, H., 2000. Macroinvertebrate functional feeding group methods in ecological assessment. *Hydrobiologia*, 422/423:225-232.
- [162] Gayraud, S., Statzner, B., Bady, P., Haybachp, A., Schöll, F., Usseglio-Polatera, P., and Bacchi, M., 2003. Invertebrate traits for the biomonitoring of large European rivers: an initial assessment of alternative metrics. *Freshwater Biology*, 48: 2045-2064.
- [163] Pavluk, T.I., bij de Vaate, A. and Leslie, H.A., 2000. Development of an Index of Trophic Completeness for benthic macroinvertebrate communities in flowingwaters. *Hydrobiologia*, 427:135-41.
- [164] Merritt, R.W., Cummins, K.W., Berg, M.B., Novak, J.A., Higgins, M.J., Wessell, K.J. and Lessard J.L., 2002. Development and application of a macroinvertebrate functional-group approach in the bioassessment of remnant river oxbows in southwest Florida.
- [165] Haybach, A., Schöll, F., König, B. and Kohmann, 2004. Use of biological traits for interpreting functional relationships in large rivers. *Limnologia-Ecology and Management of Inland Waters*, 34(4): 451-459.
- [166] Nobel, R.A.A., Cowx, I.G., Goffaux, D. and Kestmont, P., 2007. Assessing the health of European rivers using functional ecological guilds of fish communities: standardising species classification and approaches to metric selection. *Fisheries Management and Ecology*, 14: 381-392.
- [167] Charvet, S., Kosmala, A. and Statzner, B., 1998. Biomonitoring through biological traits of benthic macroinvertebrates : perspectives for a general tool in stream management. *Archiv für Hydrobiologie*, 142: 415-432.
- [168] Usseglio-Polatera P, Beisel JN. 2002. Longitudinal changes in macroinvertebrate assemblages in the Meuse River: anthropogenic effects versus natural change. *River Research and Application*, 18:197-211.
- [169] McGrady-Steed, J., Harris, P.M. and Morin, P.J., 1997. Biodiversity regulates ecosystem predictability. *Nature*, 390: 162-165.
- [170] Van der Heijden, M.G.A., Klironomos, J.N., Ursic, M., Moutoglis, P., Streitwolf-Engel, R., Boller, T., Wiemken, A. and Sanders, I.R., 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*, 396: 69-72.
- [171] Ghilarov, A.M., 2000. Ecosystem functioning and intrinsic value of biodiversity. *Oikos*, 90: 408-412.
- [172] Schwedt, G., Reiter, C., Uthemann, R. and Grabert, E., 1997. Development of an automated bacterial luminescence test for biomonitoring of environmental contaminants. *Fresenius' Journal of Analytical Chemistry*, 359: 155-160.
- [173] Hill, B.H., Lazorchak, J.M., McCormick, F.H., Willingham, W.T., 1997. The effects of elevated metals on benthic community metabolism in a Rocky Mountain stream. *Environmental Pollution*, 95: 183-190.
- [174] Hill, B.H., Herlihy, A.T., Kaufmann, P.R. and Sinsabaugh, R.L., 1998. Sediment microbial respiration in a synoptic survey of mid-Atlantic region streams. *Freshwater Biology*, 39: 493-501.
- [175] Hill, B.H., Hall, R.K., Husby, P., Herlihy, A.T., Dunne, M., 2000c. Interregional comparisons of sediment microbial respiration in streams. *Freshwater Biology*, 44: 213-222.
- [176] Gerhardt, A., Carlsson, A., Ressemann, C. and Stich, K.P., 1998. New online biomonitoring system for *Gammarus pulex* (L.) (Crustacea): in situ test below a copper effluent in South Sweden. *Environmental Science and Technology*, 32: 150-156.
- [177] Hardersen, S., 2000. The role of behavioural ecology of damselflies in the use of fluctuating asymmetry as a bioindicator of water pollution. *Ecological Entomology*, 25: 45-53.
- [178] Hill, B.H., McCormick, F.H., 2000. Nutrient uptake and community metabolism in streams draining harvested and old growth watersheds: a preliminary assessment. In: Guldin, J. (Ed.), *Proceedings of the Symposium on Ecosystem Management Research in the Ouachita and Ozark Mountains*, General Technical Report No. XX, Southern Research Station, USDA Forest Service, Oxford, MS, pp. xx-xx.

- [179] Buffagni, A. and Comin, E., 2000. Secondary production of benthic communities at the habitat scale as a tool to assess ecological integrity in mountain streams. *Hydrobiologia*, 422/423:183-95.
- [180] Buffagni, A., 2001. The use of benthic invertebrate production for definition of Ecologically Acceptable Flows in mountain rivers. In: *Hydro-ecology: Linking Hydrology and Aquatic Ecology*. Proceedings of Workshop HW2, Birmingham, UK, July 1999. IAHS Publication No. 266, 2001: 31-41.
- [181] Entrekin, S.A., Rosi-Marshall, E.J., Tank, J.L., Hoellein, T.J. and Lamberti, G.A., 2007. Macroinvertebrate secondary production in 3 forested streams of the upper Midwest, USA. *Journal of the North American Benthological Society*, 26(3): 472-490.
- [182] Carew, M.E., Pettigrove, V. and Hoffmann, A.A., 2003. Identifying chironomids (Diptera: Chironomidae) for biological monitoring with PCR-RFLP. *Bulletin of Entomological Research*, 93: 483-490.
- [183] Sharley, D.J., Pettigrove, V. and Parsons, Y.M., 2004. Molecular identification of *Chironomus* spp. (Diptera) for biomonitoring of aquatic ecosystems.
- [184] Szabó, K.É., Ács, É., Kiss, K.T., Eiler, A., Makk, J., Plenković-Moraj, A., Tóth, B. and Bertilsson, S., 2007. Periphyton-based water quality analysis of a large river (River Danube, Hungary): exploring the potential of molecular fingerprinting for biomonitoring. *Arch. Hydrobiol. Suppl.*, 161/3-4: 365-382.
- [185] Gillespie, R.B., Guttman, S.I., 1999. Chemical-induced changes in the genetic structure of populations: effects on allozymes. In: Forbes, V.E. (Ed.), *Genetics and Ecotoxicology*. Taylor & Francis, Philadelphia, pp. 55-77.
- [186] Lazorchak, J.M., Hill, B.H., Brown, B.S., McCormick, F.H., Engle, V., Lattier, D.J., Bagley, M.J., Griffith, M.B., Maciorowski, A.F. and Toth, G.P., 2002. USEPA biomonitoring and bioindicator concepts needed to evaluate the biological integrity of aquatic systems. *Trace Metals and other Contaminants in the Environment*, 6: 831-874.