# **Research & Reviews: Journal of Ecology**



ISSN: 2278-2230 (Online) Volume 8, Issue 1 www.stmjournals.com

# Sacred Groves—Repository of Medicinal Plant Resources: A Review

## Sayantani Chanda\*, T.V. Ramachandra

Energy and Wetlands Research Group, Centre for Ecological Sciences, Indian Institute of Science, Bangalore, Karnataka, India

#### Abstract

Sacred groves are traditionally protected forest or forest patches, with the rich repository of diverse flora with the medicinal plants. Ethno-medicinal knowledge on sacred grove plants have enormous prospects either in the form of natural resources and bio-prospecting. Now a days these small patches suffers from multiple disturbances grazing, encroachment, changing social perspective and hence medicinal plant resources. This study reviews medicinal plant resources in the sacred groves on distribution and usage pattern in five zones of Northern part of India. The study reported 827 medicinal plants where Leguminosae are the dominant family. Dominants plants include herbs with leaves having major use values. The endeavour emphasise the need to RET species of medicinal values.

Keywords: Medicinal plants, informant consensus factor (FIC), Fidelity level (FL). Threat, Conservation

\*Author for Correspondence E-mail: isayantani.chanda@gmail.com

#### INTRODUCTION

Historically indigenous people with the inheritance of traditional and cultural legacy were protecting the sacred groves with the practice of nature worship. It is well known that rural inhabitants of the developing countries have been using herbal medicine as healthcare to an extent of 80% global population (WHO 2001). It is also indicated that the usage of medicinal plant in regards to primary health care is still occupying the prominent position in developing countries (WHO 2002) [1]. Sacred groves are rich in rare and endemic species which are unharmed and protected by the local peoples due to the belief of the deities in the forests [2].

Tribal people in these grove areas depend on traditional medicine system for healthcare [3]. A sacred grove creates a cultural space among communities to affirm the identity and solidarity [4]. Sacred groves have been the repositories of valuable medicinal plants and wild relatives of cultivated species which have potential to address the countries food and medicine [5]. In India, studies on sacred groves with regard to ethno-medicinal plants

emphasized the local utilization of this age old tradition. Based on the review of contemporary works such as published literature, effort has been made to maintain and manage the collected records and information in consolidated form towards reviewing various aspects of ethnomedicinal plants. Objectives of the current study are (i) to develop a base line information on medicinal plant resources from sacred grove in five zones of Northern part of India, (ii) to understand the plant distribution and usage pattern, (iii) information on threats will play role in conservation and management planning.

#### METHODOLOGY

The current study reviews published literatures to understand the relevance of sacred groves and the goods and services.

#### Study Area

India has six broad geographical regions namely North-East India, Northern India, Eastern India, Western India, Central India and South India. Among these, the study covers: **North-East India:** Sikkim, Assam, Arunachal Pradesh, Nagaland, Manipur, Mizoram, Tripura, Meghalaya.

**Eastern India:** West Bengal, Bihar, Jharkhand, Orissa.

**Northern India:** Uttar Pradesh, Himachal Pradesh, Uttarakhand, Punjab, Haryana, Jammu and Kashmir.

Western India: Rajasthan, Gujarat,

Central India: Madhya Pradesh, Chhattisgarh.

The study documented 489 sacred groves distributed across the study region (Figure 1a and b).

#### **Data Collection**

Data on ethno-medicinal plants in the sacred groves has been compiled from published papers. Effort has been made to analyse the collected data family wise medicinal plants, spatial distribution with the potential uses.

#### **Data Analysis**

Ethno-medicinal data were analysed to identify proportions of plant families with the known medicinal values, habit, plant parts, frequency and popularly used plants.

# Statistical Analysis Fidelity level (FL)

Fidelity level (FL) was computed (Friedmanetal, 1986) as per equation 1, to determine the most preferred ethnomedicinal plant species used in the treatment of a particular ailment [1].

$$FL (\%) = (Ip/Iu) \times 100 \tag{1}$$

Where, Ip is the number of informants who independently indicated the use of a species for the same major ailment and If the total number of informants who mentioned the plant for any major ailment (Friedman et al. 1986).

#### **Informant's Consensus Factor (ICF)**

Informant consensus factor (ICF) given in equation 2 [1], Heinrich et al. 1998, Trotter and Logan, 1986] provides insights to the use value of particular species among the large proportion of people based on the level of homogeneity among information provided by different informants.

$$ICF = (Nur - Nt)/(Nur - 1)$$

Where, Nur = number of use reports from informants for a particular plant-use category; Nt = number of taxa or species that are used for that plant use category for all informants.

ICF Values range between 0 and 1, where '1' indicates the highest level of informant consent.

#### **Determination of Threat**

International Union of Conservation of Nature and Natural Resources (IUCN) categories threat status of rare, endangered plant species. The objective creating awareness about the importance of threatened species and conservation priorities at the local level.

#### RESULTS AND DISCUSSION

A total number of 827 medicinal plant species under 544 genera and 136 families have been recorded in this study based on the published literatures. Highest number of plant species found in Eastern zone (225 spp.) followed by Northeast (207 spp.), Northern zone (150 spp.), Central zone (125 spp.) and Western zone (125 spp.), as shown in Figure 2.

The dominant plant form is herbs (35.91 %) as represented in Figure 3 followed by trees (30.83 %), shrubs (20.07%), climbers (12.93%) and orchid (0.24%). Different plant parts have been used for curing many ailments.

It was also observed that some medicinally important plants have more than one part with the medicinal uses. Leaves have major use (25%), along with root (16.13%) and stem bark (11.66%), as per Figure 4.

It appears from the study that the most dominant families are Leguminosae (89 spp.) followed by Asteraceae (45 spp.) and Rubiaceae (39 spp.), depicted in Figure 5. *Ficus* with 11 species, *Dioscoreae* and *Phyllanthus* with 8 species are the most dominant genera found in the study region being depicted in Figure 6. Here 590 members of plant species have more than one use for remedial measure.

Aegle marmelos has wide use in 18 locations of different region followed by Cynodon dactylon (17 locations), Azadirachta indica (16 locations), Ocimum tenuiflorum (15 locations) and many are reported from at least 2 locations shown on Figure 7. The five zones of India are inhabitants by local tribes like



Kinnara, Amchi, Bodo, Bhil, Bhilala and Pataya, Meitei, Rawal, Gond, Bhumij, Kurmi, Santhal etc depends on local healthcare system. The plant species which are used in different five zones of India for the treatment are listed for 33 ailments.



Fig. 1: (a) Spatial Distribution of Sacred Groves.




Fig. 1: (b) Five Zones of Study Area.




Fig. 2: Number of Plant Species in Different Zones.




Fig. 3: Plant Forms are Reported in Different Zones.




Fig. 4: Plants Part used for Curing Different Diseases.




Fig. 5: Dominant Families among Different Parts of India.




Fig. 6: Dominant Genera among Different Parts of India.




Fig. 7: Species Presence in Studied Area.

Among these, 157 plant species are noted for abdominal problem (diarrhoea, dysentery, dyspepsia, stomach ache, stomach problem), skin diseases with 141 plants species and respiratory problems (fever, cold, cough) with 138 species. Ninety plant species act as a drug (aphrodisiac, anthelmintic, laxative, febriguge, antiseptic). 52 plant species act as remedy for the persons bitten by animals.

It was also reported that the plant species have medicinal benefits for animals. The ethnoveterinary medicine used from 12 plants against 9 disease (galactagogue, ulcer, wounds, eye disease, urination, loose evacuation, stomach ache, typhoid fever and for removing parasites from their skin.). Stem bark and seeds are most used plants parts for veterinary medicine than root, whole plants, leaves and fruits. The route of administration for animals is oral by using plant parts in the form of decoction, externally as a paste and powder. The ethno medicine for humans are used mostly paste followed by juice, powder, decoction as shown in Figure 8.

### **Fidelity Level**

Table 1 lists the FL values which highlight the most important medicinal species among the five zones of India. Fidelity Level (FL) of plants has been calculated based on most dominant plant and showed the corresponding plant against a particular ailment. Higher values of Fidelity Level (%) indicate the most preferred dominant plant species for each particular ailment category. The analysis

showed that the highest FL value found in 3 plant species among the five study region for different diseases are *Gymmema sylvestre* (diabetes, 100%) followed by *Biden pilosa* (tooth problem, 100%) and *Valernia jantamansi* (stomach ache, 100%). The least FL values for *Oxalis corniculata* (23%) which indicates less preferred plant species in regards to constancy.

#### **Informant Consensus Factor (ICF)**

ICF is considered to know the traditional values of plants among the five zones of India to treat certain ailment categories (Table 2). From the study literature, out of the 489 sacred groves, 829 species with medicinal values, treating 33 different ailments. In our study, for the sake of simplicity out of 33 different ailments only 10 ailments are categorized. It is clear from the study that the ICF values varied from 0.04 to 0.64. Cold has the highest ICF value 0.64 with 40 use-reports for 15 plant species followed by diarrhoea have 0.24 with 76 use-reports for 58 plant species, wounds and cuts have 0.24 with 103 use-reports for 78 plant species.

The high ICF value for cold possibly showed that this ailment is common in the study area. High ICF values also helps searching for bioactive compounds and subsequently considered as more preferred traditionally to treat ailments. The low ICF value is for asthma due to lack of less constancy about the ailments.

#### **Threats**

These small patches have multiple disturbance factors like anthropogenic pressure, pollution, urbanization, soil erosion, logging, agriculture conversion of forest into land and road construction, invasion, over grazing, encroachment, developing industries, shifting social and cultural perspectives which put these medicinal plant resources under threat

category. Out of 829 medicinal plant species, herbs species have 3 spp. endangered, 1 spp. critically endangered, 27 spp. under least concern, where tree species, have 1 spp. near threatened, 2 spp. vulnerable, 18 spp. least concern, shrub have 8 spp. least concern, climber have 1 spp. vulnerable and 2 spp. least concern and even orchid also 1 spp. under least concern.




Fig. 8: Route of Administration used in Health Care System.

| <b>Table 1</b> : Dominant I | Plant Species for a I | Particular Diseases and | their Fidelity Level. |
|-----------------------------|-----------------------|-------------------------|-----------------------|
|                             |                       |                         |                       |

| Disease       | Dominant plants         | Fidelity level |
|---------------|-------------------------|----------------|
| Diarrhoea     | Aegle marmelos          | 37.5%          |
| Cold          | Ocimum sanctum          | 41.17%         |
| Wounds & cuts | Tridax procumbens       | 78%            |
| Skin problem  | Azadirachta indica      | 70%            |
| Malaria       | Andrographis paniculata | 50%            |
| Diabetes      | Gymmema sylvestre       | 100%           |
| Tooth problem | Bidens pilosa           | 100%           |
| Piles         | Oxalis corniculata      | 23.07%         |
| Stomach ache  | Valernia jantamansi     | 100%           |
| Asthma        | Emblica offcinales      | 25%            |

Table 2 Category of Different Diseases and their Informant Consensus Factor (ICF).

| Disease         | No of Plant Species | Used Report | ICF  |
|-----------------|---------------------|-------------|------|
| Diarrhoea       | 58                  | 76          | 0.24 |
| Cold            | 15                  | 40          | 0.64 |
| Wounds and cuts | 78                  | 103         | 0.24 |
| Skin problem    | 141                 | 185         | 0.23 |
| Malaria         | 23                  | 29          | 0.21 |
| Diabetes        | 39                  | 49          | 0.2  |
| Tooth problem   | 42                  | 54          | 0.12 |
| Piles           | 45                  | 52          | 0.07 |
| Stomach ache    | 17                  | 20          | 0.15 |
| Asthma          | 30                  | 34          | 0.04 |



#### CONCLUSION

This study highlights the importance of sacred in rural informal healthcare management. Apart from its socio-religious importance, medicinal plants are one of the active benefits which people can connect with easily. Plants with multiple uses require attention restoration special in conservation planning. The usefulness and protection of all the reported ethno medicinal plants need to be evaluated by phyto-chemical and pharmacological studies. Plants with high informant consensus factor, use report and fidelity level should be given main concern to carry out bioassay and toxicity studies. generation, participatory Awareness management planning, local awareness and thorough evaluation of these plant resources are key factors for long term survival of these groves.

#### REFERENCES

- An ethnobotanical study of medicinal plants used in sacred groves of Kumaon Himalaya, Uttarakhand, India, Harsh Singh Journal of Ethnopharmacology, Vol 154, Date 2014, Page No-98–108.
- 2. Inventory of ethnobotanicals and other systematic procedures for regional conservation of medicinal and sacred plants, Vijay V. Wagh, Ashok K. Jain, Environment Systems and Decisions, Volume 35, Page No-143–156, 2015.
- 3. Sacred and Medicinal Plant Diversity of Vandiol Sacred Grove of Sabarkantha District (N.G.) D.N. Parmar and N.K. Patel Life Sciences Leaflets, Volume-5, Page No-34 to 49.
- 4. Ethnomedicinal Plant Conservation Through Sacred Groves, R. K. Bhakat and U.K. Sen, Tribes And Tribals, Vol 2, 2008, Page No. 55-58.
- 5. Role of sacred groves in the conservation and management of medicinal plants Manoj Kumar Behera, Tapas Ranjan Pradhan and Jangyeswar Sahoo, Journal Of Medicinal Plants Research, Vol 9 No 29,3 August 2015, Page No. 792-798.
- 6. Sacred plants and their Ethno-botanical importance in central India: By Pankaj K. Sahu, A. Kumari, Shweta Sao, M. Singh and P. Pandey, International Journal of Pharmacy & Life Science, Vol 4, Issue 8, Aug 2013, Page No. 2910-2914.

- 7. Phytodiversity of Ethnomedicinal plants in Sacred groves and its Traditional Uses in Kabirdham District of Chhattisgarh, C.P. Rahangdale, R.K. Patley and K.C. Yadav, Indian Forester, Vol-140 No-1, 2014,86-92.
- 8. Assessment of Status and role of sacred grove in conservation of Biodiversity at Different Level of Madhya Pradesh District Chindwara, Dr. J. L Shrivastava, Thesis
- 9. Deogudi' Sacred Grove. A Tribal Concept of Conservation of Plants In Bastar. Rajiv Rai and Satya PrakashTripathi, The Indian Forester, Volume 134, Page No-1686-1695, 2008 December.
- Observation of Medicinal Importance of Sacred Plants of Chitrakoot Region Satna (M.P.) Lipika Devi Bala, Ravindra Singh International Journal of Science and Research (IJSR) Volume 4(8).
- 11. Role of Sacred Grove in Religion and Health Care System of local people of Almora district of Uttarkhand State, Vijay Sharma Academia Arena, Vol 2 No 6, 2010, Page No 19-22.
- 12. Biodiversity conservation through a traditional beliefs system: a case study from Kumaon Himalaya sacred grove Harsh Singh, International Journal of Conservation Science Vol 3, Issue 1, January-March 2012, Pg No 33-40.
- 13. Sacred Grove in conservation of plant biodiversity in Mahendagarh district of Haryana Sanjay Yadav, J.P Jadav, Vedpriya Arya and Manju Panghal, Indian Journal of Traditional Knowledge, Vol 9 No 4,October 2010,Page No-693-700.
- 14. Local Deities in Conservation. A conservation Practice In Nbanju Nami Tok Sacred Grove In Tehri Garhwal, Uttarkhand, Nazir A Pala Indian Forester Vol 138 No 8, 2012, Page 710-713.
- Patalbhuvneshwar: a new sacred grove from Kumaon Himalaya, Priyanka Agnihotri, Harsh Singh and Tariq Husain, Current Science, Vol 102 No 6, 25 March 2012.
- 16. Role of traditional conservation practice: highlighting the importance of Shivbari sacred grove in biodiversity conservation, Vikrant Jaryan, Sanjay Kr. Uniyal, Gopichand, R.D. Singh, Brij Lal, Amit Kumar, Varun Sharma Environmentalist Volume 30, Page No-101–110, 2010.

- 17. Ethnobotanical Studies of Sada Shiv Sacred Grove, District Kangra, Himachal Pradesh, Pooja Sharma, Virendar K. Santavan. Arti Sharma and Bhagwati P. Sharma, Life Sciences Leaflets, Volume No-53, Page No-89 to 96, 2014, 10th June.
- 18. Sacred Grove As Ethnobotanical Gene Pool On Tribal Area of Western Himalaya, Vandana Thaplyal, Sanjeev Sharma, Ajay Ballabh Bhatt The Indian Forester, Volume 138, Page 70-77, 2012, January.
- 19. Sociocultural and ethnobotanical value of Sacred Forest, Thal ke Dhar, Central Himalaya, Chandra Singh negi.
- 20. Medicinal Plant Diversity in Newly Reported Sacred Grove of Pithoragarh District, Uttarakhand Harsh Singh, Pushpi Singh and Tariq Husain, Indian Forester, Page No-1005-100.
- 21. An assessment of ethnomedicinal plant diversity in Haat Kali Sacred Grove in Kumaun Himalaya, Uttarakhand, India Harsh Singh, Priyanka Agnihotri, Veena Dixit, P.C. Pande and Tariq Husain East Himalayan Society for Spermatophyte Taxonomy, Volume-9 (2),Page No-456 464
- 22. Traditional knowledge and biodiversity conservation in the sacred groves of Meghalaya, S Jeeva, BP Mishra, N Venugopal, L Kharlukhi, RC Laloo, Indian Journal of Traditional Knowledge Vol 5 No 4,October 2006, Page 563-568.
- 23. Studies on Plant Diversity and regeneration of a Few Tree Species in the Sacred Grove of Manipur, Khumbongmayum Ashalata Devi, Thesis.
- 24. Ethnomedicinal Plant in the Sacred Grove of Manipur, Khumbongmayum Ashalata Devi, Indian Journal of Traditional Knowledge, Vol 4 No 1, January 2005, Page No 21-32.
- 25. Distribution and conservation Status of Sacred Grove in Garo Hills, Meghalaya, R.K Mohanta, The Indian Forester December 2009, Page No-1627-1649.
- 26. Plants Of Ethnobotanical Importance In The Sacred Groves Of Jaintia Hills Of Meghalaya, K. Upadhya, H.N. Pandey, P.S Law and R.S.Tri [pathi, The Indian Forester, Volume 131, Page No-812-828, 2005.

- 27. Sacred Groves. An analysis made in the cultural perspective with BTC Assam, India
- 28. Jahnovi Brahma, Binay Singh and Tribeni Mandal, Journal of Biological & Scientific Opinion, Vol 2 No 4,2014.
- 29. Status of medicinal plants in the disturbed and the undisturbed sacred forests of Meghalaya, northeast India: population structure and regeneration efficacy of some important species, R. C. Laloo, L. Kharlukhi, S. Jeeva and B. P. Mishra, Current Science, Vol 90, No 2, 25 January 2006, Page No-225-231.
- 30. Geo environmental appraisal of sacred groves and its releated traditional practice in West Bengal
- 31. Mr Tapas Pal, Department of Geography, Visva-Bharati Santiniketan, Thesis, Page 53-58.
- 32. Sacred Groves: traditional Plant conservation through deities, RK Bhakat and UK Sen, The article published in this book among the paper presented UGC sponsered national Seminar organized by the department of Botany, Sonamukhi college, Page No-212-223, 2016, October.
- 33. Sacred Grove of West Midnapore District West Bengal, R.k Bhakat and U.K Sen, Recent Advances in Plant Science, Diversity and Conservation(Proceeding of UGC National Seminar 2011), 2011 January.
- 34. An inventory of Medicinal Plants of Some Sacred Groves Of Purulia District, West Bengal, R.K Bhakat and P.K. Pandit, The Indian Forester, Volume 130, Page 37-44, 2004, January.
- 35. Biodiversity and Ethnobotany Of Sacred Groves. In Bakura District, West Bengal, Ramsankar Basu, The Indian Forester, Volume 135, Page 765-777, 2009, June.
- 36. Role Of Sacred Grove In Conservation Of Medicinal Plants, R.K Bhakat and P.K. Pandit, The Indian Forester, Volume-129, 2003, Feburary.
- 37. Floristic and ethnobotany of sacred groves of Kheda District (Gujrat) and their significance in conserving biodiversity, Dr. Kaushik C. Patel, Hemchandracharya North Gujrat University, Patan, From 106.



- 38. Ethnobotanical study of Sacred groves of Poshina forest of Sabarkanth district, North Gujarat, P.K. Mehta and B.K. Jain, International Journal of Plant Sciences, Volume 6, Page No. 362-366.
- 39. An Ethnobotanical Study of Medicinal Plants Used In Sacred Groves Of Ambaji Forest Gujarat, India Dr. R. S. Patel, International Journal of Advanced Technology in Engineering and Science, Volume No 03, Special Issue No. 01, March 2015.

# **Cite this Article**

Sayantani Chanda, T.V. Ramachandra. Sacred Groves – Repository of Medicinal Plant Resources: A Review. *Research & Reviews: Journal of Ecology*. 2019; 8(1): 12–20p.